2018

PULSES IN INDIA: RETROSPECT AND PROSPECTS

Compiled and Edited by Dr. A.K. Tiwari Director

Dr. A.K. Shivhare Assistant Director



Directorate of Pulses Development Government of India

Ministry of Agriculture & Farmers Welfare

Department of Agriculture Cooperation & Farmers Welfare

Vindhyachal Bhawan, Bhopal – 462004, Madhya Pradesh

Web: www.dpd.gov.in

Compiled and Edited by

A.K. Tiwari, and A.K. Shivhare

Year of Publication: 2019

Correct Citation

A.K. Tiwari and A.K. Shivhare – 2019. *Pulses in India: Retrospect and Prospects* – 2018. Directorate of Pulses Development, GoI, Bhopal – 462004 (M.P), India pp-1-346.

Published by

Director

Directorate of Pulses Development, Government of India, Ministry of Agriculture & Farmers Welfare Department of Agriculture Cooperation & Farmers Welfare Bhopal – 462004, Madhya Pradesh, India

Web: www.dpd.gov.in

© 2019

All rights reserved

No part of this publication may be reproduced in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without the prior permission of the copyright owners.

ISBN 978-81-949512-3-0

Printed by

Delhi Sales Corporation

BS – 103A, Shalimar Bagh, Delhi-110088

Tel: 011-27476373, M-9312219848

Email – dsc1987@yahoo.com, mukeshbansal07@yahoo.com

PULSES IN INDIA: RETROSPECT AND PROSPECTS

Compiled & Edited by
Dr. A.K. Tiwari
Director
Dr. A.K. Shivhare
Assistant Director



DIRECTORATE OF PULSES DEVELOPMENT

GOVERNMENT OF INDIA

MINISTRY OF AGRICULTURE & FARMERS WELFARE
DEPARTMENT OF AGRICULTURE, COOPERATION & FARMERS WELFARE
VINDHYACHAL BHAVAN, BHOPAL-462004 (M.P.)



D.O.No. 1815/AM



कृषि एवं किसान कल्याण मंत्री भारत सरकार

A FARMERS WELFARE
GOVERNMENT OF INDIA
2 9 AUG 2018



पाधा मोहन सिंह RADHA MOHAN SINGH

संदेश

भारतीय अधेकावस्था में सुषि की महत्वपूर्ण धूमिका है। सुषि क्षेत्र को राष्ट्रीय सकत वरेतू बत्यादन में 16 प्रविशत से अधिक एवं रोजगार क्षेत्र में 49 प्रतिशत से अधिक बांगदान है और वर्तमान सरकार है 2022 एक खूबकों की आमदनी को यो गुना करने व विश्वत विभागन व्यवस्था हेतु सहत्वपूर्ण कार्म किसे हैं।

राष्ट्रीय काब सुरक्षा अधिनिमम-2013 के दुव्यिगत एवं मानव विकास सूचकांका की चुनौती को स्वीकार करते हुने पीवण मुक्त खाख की उपलब्धता दुवं पर्यावरण हितेशी सम्बन्धती विकास की प्रतिबद्धता, वर्तमान सरकार की सभी और प्रदेशक एवं केंद्र समर्थित सोजनकों में परिलक्षित है। व्हर्षि खेल में भीतियों, सार्यक्रमों एवं पूंजी निवेश का महत्वपूर्ण सोग्रहान है, तथापि क्रमीनी स्वर पर प्रधानी कार्यक्रम क्रियान्वयन हो सफलता की वहंगी है।

खुबको से सरात् सम्मर्क, गरमंग्रात इतन व उन्तर तकनीकी इस्तांतरण के साथ ही साथ समन्वय पर आधारित विकास कार्यक्रम आंत आवश्यक है। इस पर सरकार स्वत् कार्य कर एही है। विगत तीन वयों में इसहण उत्पादन में आशासीन प्रगति हुई है। वर्ष 2017-18 में इसहण के सर्वोच्च 252 साख दन उत्पादन के साथ इसहण क्षेत्र में देश के आवारिनंत्ता प्राप्त की है। नारत सरकार, मूर्वि सहकारिता एवं कृषि कल्याण विभाग के कसल विकास निवेशालयों, एक्य कृषि विभाग एवं अन्य सहनागी संस्थाओं के समन्वय, आवंक्रमों के प्रभावी क्रियान्यवन व करात अनुश्रवण एवं खुबकों के अवाह परिश्रक का इस उपलब्धि में अहम मुक्तिका रही है।

मुझे प्रसानका है कि राष्ट्रीय खाध सुरक्षा मिशन, फासक अनुवाग-वलक्षण विकास निवेशालय, कृषि एवं किसान कल्याण मंत्रालय, भारत सरकार, भोषास हारा "पल्सेस इन इण्डिया–रिहोस्पेकट एवं प्रोसपेक्ट्स" बहुउपमोगी दलकन से संबंधित जानकारी का संकलन कर प्रकाशन किया गया है, जो कि आहे प्रशंसनीय है।

आरम है कि यह प्रकारण पराहत क्षेत्र में काचेरत नीति निर्भारको, विकास कार्यक्रम क्रियान्वयन संस्थाओं, विचार्वियों, कोश्रार्वियों, इच्चमियों, कृषक अत्यादक समृहों व स्वसहाधता समृहों आदि को लिये अस्पन्त अपयोगी किन्न होगा।

्रांधा मोहन सिंह

Office: Room No.128, Krishi Bhawan, New Delhi-110 001 Tel.: 23383370, 23782891 Fax: 23384129

Pulses in India: Retrospect and Prospects -2018

DPD, Govt. of India, Bhopal\



S.K. PATTANAYAK SECRETARY



Ministry of Agriculture & Farmers Welfare Department of Agriculture, Cooperation & Farmers Welfare.

FOREWORD

Our world today faces a tough challenge, for ensuring food security while providing a balanced diet for everyone. Overcoming hungar and mainutrition in the 21st century means increasing food quantity and quality, while making sure we produce food sustainably and efficiently. Pulses have been an essential part of the human diet for centuries. Yet their nutritional value is not generally recognized and their consumption is frequently under appraciated. Pulses in India have long been considered as the poor man's only source of protein, thus play a crucial role in healthy diets, sustainable food production and above all, in food security.

An improvement in pulses production technology can reduce the cost of production and ensure higher productivity resulting in affordable prices to consumers. This will create soons for further increase in demand for pulse crops by replacing some portion of the disproportionably high level of cereals in the consumption basket for a balanced diet. Considering the importance of pulses, Department of Agriculture. Cooperation and Farmers Welfare in the Ministry of Agriculture & Farmers Welfare has made sincere efforts through National Food Security Mission, which resulted in appreciable increase in the area, production and productivity of pulses in India. Recent policy initiatives under National Food Security Mission (NFSM). Bringing Green Revolution in the Eastern Region (BGREI), Crop Diversification Program (CDP) involving conduct of large scale cluster demonstrations, creation of 160 Seed-hubs for pulses, seed minikit distribution of high yielding new varieties, strengthening seed production infrastructures, seed village programme, creation of FPO's and enhanced MSP coupled with favourable EXIM policy has earned us a place of pride to achieve the highest ever production of 25.23 million tonnes of pulses during 2017-18 (4^{ft} estimates) making the nation self-sufficient in pulses. All these efforts have ushared in a mini green revolution in the pulses sector:

Any information on pulses is highly valuable since the crop is considered. as incredible to mankind in terms of cultivation, food, nutritional and income security. I appreciate the sincere efforts of the authors in bringing out this valuable publication on "Pulses in India: Retrospect and Prospects". I hope this book will be of Immense use to the farmers, students, extension workers, policy makers. and researchers.

New Delhi 13th September, 2018 Schillowayte

Office: Krishi Bhawan, New Dehi-110 001 天明4 / Phone: 23382651, 23388644 常明 机/ Fax No.: 23386004 Pulses in India: Retrospect and Prospects -2018

DPD, Govt. of India, Bhopal\



Dr. S. K. Malhotra
Agriculture Commissioner
Phone: +91-11-23383649, 23381012
E-mail: ag.comm@nic.in
agricommissioner@gmail.com



भारत सरकार कृषि एवं किसान कल्याण मंत्रालय कृषि, सहकारिता एवं किसान कल्याण विभाग कृषि भवन, नई विल्ली-110001

Government of India
Ministry of Agriculture & Farmers Welfare
Department of Agriculture, Cooperation
& Farmers Welfare
Krishi Bhawan, New Deibl-110001

Foreword

Pelses have long been considered as good source of protein, thus play a crucial role in healthy diess, sustainable food production and above all in food security. Food legumes are next to cercule in terms of agricultural importance and are the best options for diversification and intensification of contemporary agriculture on sustainable basis. India appreciably has 35% share in global area and production and is largest producer and consumer of pulses.

In India, pulses are produced with minimum use of resources and honce become less costly than uniqual protoin. Besides, pulses are the continguacy crops for the small and marginal farmers where they suit well as an inter trop and mixed exop in the vulnerable arid and semi-arid agre sensystem. Palsa erepe are low cost protein source, could replace proportionately high cereals consumption for a balanced diet. It was felt that with increasing availability of short duration pulse varieties, coupled with critical intercentions at developmental from through NFSM/ RKVY, the logistic extension approach would definitely enhance the production. Accordingly, National Food Scourty Mission - Pulses, implemented breeder seed production program, created 150 Seed-habs, distributed need minikits of high yielding new varieties, strengthened seed production infragructures. conducted large scale observed emonstrations, started seed village programs, created EPO's and enhanced MSP and thus it has become possible to achieve ever highest production of 25.23 million tennes of pulses during 2017-18 (4th estimates). The agriculture accounts for just 16 per cent of India's economic output, but provides a livelihood to more than 49 per cent of its total population. The inserventions under NESM-Puises have realized the positive impact in onbancing income of farmers. During 3017-18 an approximate increase in pulses has been to the tune of 18% new and 34% production and 13% yield, consequently per eapita aveilability of pulses has also increased during last 4 years.

Present publication on "Padeus in India: Retrospect and Prespects" provides a comprehensive review on pulses production trends; global and national securite; production, poetection and post harvest technologies of chickpan, pigeon pen, black gram, green gram, lentil, field pen, mothbean, horsegram, lathyrus, french bean and cowpen. I compliment the authors for their sincere offeres in bringing out this updated publication. I am sure the publication will be of high use to policy planners, researchers, farmers, students and industry.

New Delbi 12th Suptember, 2018 (Dr. S. K. Malhotra)



Dr. B. Rajender, IAS Joint Secretary



भारत सरकार कृषि एवं किसान कल्याण मंत्रालय कृषि, सहकारिता एवं किसान कल्याण विभाग Government of India Ministry of Agriculture & Farmers Welfare Department of Agriculture, Cooperation & Farmers Welfare

Foreword

India is the world's largest producer and consumer of pulses. Pulses have their significance in ensuring nutritional security of the vegetarian common man owing to higher protein, carbohydrates, fibers, some B' vitamins, trinerals and low in fat. The protein content of pulses is twice that of wheat and three times that of rice. The impact of pulses on human health as also on sustainability of production system is well established by research.

Pulses programme were kept to centre stage owing to their nutritional importance to contain the protein malnutrition vis-à-vis their role in the economy of >40 per cent total human population of the country carning their livelihood in the minfed regions where 80 per cent of total pulses are grown.

For effective developmental planning and its implementation/governance, information and knowledge management of the sector is very important.

In compiling the commodity specific document on "Pulses in India: Retrespect and Prospects" Dr. A.K. Tiwari, Director, Directorate of Pulses Development, Bhopal has made a significant contribution to the understanding of the subject.

Plan-wise analysis on critical development interventions, pulse scenario, individual crop's status, their feasibility and scope, agro-climatic zone-wise constraints and suggestive strategy, issues on seed, post harvest/processing, demand-supply analysis etc., makes this compilation more comprehensive and informative.

I hope the gublication, "Pulses in India: Retrospect and Prospects" will go a long way in providing much needed information to students, researchers and policy makers in the field.

New Delhi

13th September, 2018

Dre (Dr. B.Rajender)

Office: Krishi Bhawan, New Delhi-110 001 पूरभाष / Phone : 23381176 प्रेमिस सं./ Fax No.: 23382417 Nich.: 9599428809 E-moil: b.rajender@los.nic.in



Dr. A. K. Tiwari Director



भारतं सरकार कृषि एवं कियान करूपाण गंत्रालय कृषि, सडकारिता एव कियान करुपाण विभाग यलहन विकास निर्मेशालय भोपाल-462 004 (ग.प्र.) Government of India Ministry of Agriculture & Formers Welfare lepts of Agriculture, Cooperation & Formers Welfare

Ministry of Agriculture & Formers Welfare (Depts of Agriculture, Cooperation & Formers Welfare) Directorate of Pulses Development Biopal 462004 (M.P.)

PREFACE

The existing population of India at 1.36 billion which is 18% of world's population is likely to cross 1.31 billion by 2030. The demographic dividend of the country provides an inherent edge and potential to its economy to grow much faster:

The country has made rapid strides in improving rates of under and makestrition, yet it continues to have a serious challenge to meet out the aims and objectives of National Food Security Act (NFSA)-2013, to ensuring food and autritional security to the most vulnerable. The zero hunger Challenge (ZHC) is extremely relevant for India.

Pulses are the chief source of vegetable protein for large section of vegetarian society containing 20-25 per cent of protein which is twice the content of wheat and thrice that of rice. Pulses are rich source of other minerals, fibers and some form of vitamin B'.

During 2017-18, the nation harvested a record production of 25.24 Million tennes of pulses in 29.99 Million ha at a national yield of 841 kg/ha. The per capita per day availability has enhanced to about 53 g t.e. >19 kg/head/annum. The share of pulses to total food grain basket has increased to about 5 per cent from the stagmated level of 6-7 per centup till 2015-16.

In its resolve to double the farmers' income by 2022 and implement NPSA-2013 to solitore the global target of zero hunger by 2030, the DAC & PW has initiated many programmes like NPSM, RKVY, PMKSY, PMFBY, c-merket place (c-NAM), MIDII, formation of FPO and sustained increase in MSFs. The NPSM-Pulses is implemented in all 29 states/558 districts, especially targeting >73 Mhs rainfed area, 80 per cent pulse area of the country.

Pulses have received appoints attention of the contral government since 2015-16 in terms of meeting-out the consumers demand at affordable rates and adequate programmatic support at the production front including regular field meditoring. The gume changing efforts under the 'Prima Minister's Krisht Sinchayi Yujana', IT initiatives in exposion/apps to access market, Soil Health Cards, INM, aron advisories and B-NAM, involvement of KVKs and creation of 150 seed hubs, enhancing breeder seed production (BBSP) by form infrastructure at ICAR/SAUs, formation of FPOs, creation of buffer stock, imposition of stock limits and offseling pulses at low cost through mobile vans, imposition of import duties and exim policy and announging Foreign Direct investment (FDI) in food processing etc., have been the major policy interventions.

The document, Pulses in India: Retrospect & Prospects-2018, is the IV* in the series of the publication after 2008, 2016 and 2017. The first such publication was brought out in 2006 under the able guidance/forward of Dr. P. K. Miahra, the then Secretary DAC&FW, Govt. of India, (Presently the Additional Principal Secretary to Hon'ble Prime Minister).

The document consist of the overview of the pulse sector, domand and availability status, import-export, impact of plan period interventions on its growth, market scenario, constraints based identified recommendations for sustained growth of the pulses to address the redritional security vis-à-vis doubling the farmers' income by 2022. It also has a vision-2030 analysis. To further enrich the publication, all 12 adials pulses, including the minor pulses have been incorporated in terms of their economic impostance agreement, varietal diversification and the potential regions/districts etc.

I hope, the publication "Pulses in India: Retrospect and Prospects-2018" would benefit the intelligentain Plancers, farmers, FPOs, State deptt of agriculture, development organizations, processors/indexs, researchers and all the readers in sustaining the growth and soff-reliant to pulses.

I am grainful to Dr. S. K. Pattanayak, Hou'ble Scoretary, Govt. of India, Ministry of Agriculture & Farmers Welfare, DAC&FW, Dr. S. K. Malhotra, Agriculture Commissioner and Dr. B. Rajender, John Secretary (Crops/Oilseeds) for their support and guidance in bringing out the volume.

I acknowledge the association of S. Pallewar, SI and Ms. Ashwioi Tikle, Technical Assistant. Ms. Tikle deserve special mention for her sincore involvement, dedication and hard work, beyond office hours, to accomplishing the task of bringing the publication.

Bhopal

14^a September, 2018

(A.K.Tiwari)

यूरभाष /Ikiaphone: 0786-2890383 देलीफीवस:/Iwienx: 0788-2871678



है-मेल /kmail: dpd.mp@nio.in वेदरगाइट /Website: dpd.gev.In

राष्ट्रीय साध सुरक्षा मित्रान - सतत् सूचि हेट्र।

EXPLANATION TO ABBREVATIONS

ADO Agriculture Development Officer

AES Agro-ecological situations

AFC Agriculture Finance Commission
AICRP All India Coordinated Research Project

:Aølines Male sterile lines ALP Aluminium Phosphate

AMDP Accelerated Maize Development Programme

A,P,Y Area, Production, Yield

A.P Andhra Pradesh

ATARI Agriculture Technology Application Research Institute

ATMA Agriculture Technology Management Agency

∃Bølines Maintainer lines

BCMV Bean Curl Mosaic Virus
BSP Breeder Seed Production
BNF Biological nitrogen fixation
BT Bacillus thuringenesis

COPP Change over previous plan periods
CAGR Compound Annual Growth Rate

CZ Central Zone C.G. Chhattisgarh

CGMS Cytoplasmic Genetic Male Sterility

CEC Cation Exchange Capacity

CAZRI Central Arid Zone Research Institute, Jodhpur (RJ)

CZ Central Zone

CPWD Central Public Work Department

CIAE Central Institute of Agriculture Engineering, Bhopal (M.P.)

CCL Cash Credit Limit

CFTRI Central Food Technology Research Institute, Mysore (Karnataka)

CWC Central Warehousing Corporation

DAC Department of Agriculture, Cooperation & Farmers Welfare

DAP Di-ammonium Phosphate

DAS Days after sowing

DFSMEC District Food Security Mission Executive Committee
DGCI&S Director General of Commerce Intelligence and Statistics

EC Emulsifying Concentrate
ETL Economic Threshold Level
EC Empowered Committee
FAQ Fair Average Quality
FFS Farmer Field School
FOs Farmers Organizations

FPOs Farmers Producers Organizations

FIGs Farmers Interest Group FLD Front Line Demonstration

FAO Food and Agriculture Organization

FYM Farm Yard Manure
GOI Government of India
GMS Genetic Male Sterility
HDPE High Density Poly Ethylene

HI Harvest Index

HRD Human Resource Development

HP Horse Power

ITD Innovations in Technology Dissemination IFFCO Indian Farmers Fertilizer Co-operative Ltd.

IPM Integrated Pest Management

ISOPOM Integrated Scheme of Oilseeds, Pulses, Oil palm and Maize

IIPR Indian Institute of Pulse Research, Kanpur (UP)
ICAR Indian Council of Agriculture Research, New Delhi

INM Integrated Nutrient Management ICMR Indian Council of Medical Research

KCl Potassium Chloride

K Potassium

KVK Krishi Vigyan Kendra

KVIC Khadi Village Industries Commission

KW Kilo Watt

KRIBHCO Krishak Bharti Co-operative Ltd.

LE Larval Extract MM-1 Mini-Mission 1

MSP Minimum Support Price

Min. Minimum
Max. Maximum
Mo Molybdenum
MP Madhya Pradesh
MS Maharashtra

NAT New Agriculture Technology

NEPZ North East Plain Zone NWPZ North West Plain Zone NHZ North Hilly Zone

NPV Nuclear Polyhedrosis Virus

NATP National Agriculture Technology Project

NCDC National Co-operative Development Cooperation

NGOs Non-Government organization

NABARD National Bank for Agriculture and Rural Development

NPDP National Pulses Development Project NLMT National Level Monitoring Team

NUE Nutrient Use Efficiency

NFSM National Food Security Mission

NAFED National Agriculture Marketing Federation Ltd

NBSS&LUP National Bureau of Soil Survey and Land Utilisation Planning, Nagpur

NWDPRA National Watershed Development project for Rural Agriculture

NSC National Seed Corporation

NAEP National Agriculture Extension Project

NE North East

NWP North Western Parts

OPDP Oil palm Development Project

OILFED Oil Federation

OPP Oilseed Production Programme

PSHG Pulses Self Help Group
PHT Post Harvest Technology
PSB Phosphate Solubilising Bacteria

PWD Public Work Department

PGPR Plant Growth Promoting Rhizobacteria

PC Project Coordinator

PAU Punjab Agriculture University, Ludhiana (PB)

PDKV Panjabrao Deshmukh Krishi Vidyapeeth, Akola (MS)

PFA Prevention of Food Adulteration

PSS Price Support Scheme RGK Rural Gram Kendra RBI Reserve Bank of India

R Rabi

∹Rølines Restorer Lines

RAEO Rural Agriculture Extension Officer

R&D Research Development

SPPP Strategic Pulses Production Programme

SVS Seed Village Scheme
SES Socio-economic Status
SBI State Bank of India
SSP Single Super Phosphate

SDA State Department of Agriculture

SL Solubilite

SRR Seed Replacement Rate

SHGs Self Help Group

SWCs State Warehousing Corporation

SZ South Zone

SFPP Special Food Grain Production Programme

SSC State Seed Corporation

SFCI State Farm Corporation of India
SLMT State Level Monitoring Team
Taggree of Taglorals are

ToT Transfer of Technology

TMC Technology Mission on Cotton

T&V Training and Visit

TAC Technical Advisory Committee

TE Triennium Ending

TMO Technology Mission on Oilseed

TN Tamil Nadu UK United Kingdom

UNDP United Nations Development Programme

USA United States of America

UTs Union Territories
UP Uttar Pradesh
W.B. West Bengal

WSC Water Soluble Concentrate

YI Yield Index

YMV Yellow Mosaic Virus ZRS Zonal Research Station

@ At the rateCa CalciumCm Centimetre

⁰C Degree Centigrade

Hr Hour

Kg/ha Kilogram/ hectare

Ml Milli litre mg Milli gram

pH Potential Hydrogen

% Percent Square Meter

Zn Zinc

Mg Magnesium

INDEX

Message

Foreword

Preface

About the Directorate

CONTENT	PAGE NO.
Executive Summary	1-10
1. A giant leap in production	1
2. Equilibrium Production and demand	2
3. Significant growth with additional return	2-3
4. Pulses import falls by 03 million tonnes in financial year- 2018	5-6
(Saving of foreign currency)	
5. Export of pulses allowed	6
6. Interventions which proved the trigger	6-7
Conclusion	8-10
Chapter -1: Pulses in Indian Context: 2017-18	11 to 22
1.1 Background	11-12
1.2 Indiags status of pulse production	12
1.3 Pulses share to total foodgrain basket	12-13
1.4 Season & crop contribution in total pulse production (2017-18)	14
1.5 States@Contribution \(\phi(2017-18) \)	15
1.5.1 Total Pulses	15
1.5.2 Kharif Pulses	16
1.5.3 Rabi Pulses	17
1.5.4 Gram (Chickpea)	18
1.5.5 Arhar (Pigeonpea)	19
1.5.6 Mungbean (Greengram)	20
1.5.7 Urdbean (Blackgram)	21
1.5.8 Masoor (Lentil)	22
Chapter-2 : Plan-wise Pulse Overview	23 to 48
2.1 Background	23-25
2.2 Pulses share to total foodgrain basket	26
2.3 Per capita availability of pulses in India	28
2.3 Projected Demand: (XI th & XII th Plan)	29
2.5 Import/Export and Availability	30
2.6 Exim policies in favour of pulses have paid	32 to 35
2.6.1 Import	32
2.6.2 Export of Pulses	32
2.6.3 Buffer Stock	32-33

CONTENT	PAGE NO.
2.7 Growth Rate	34-35
2.8 Production Trends	36 to 48
2.8.1 Global Scenario: Crop-Wise	36
2.8.2 Global Scenario: Total Pulses	37
2.8.3 National Scenario: Plan Periods (Total Pulses)	38
2.8.3 StatesøScenario: Plan Analysis (X th óXII th)	39
2.8.5 District Scenario (2016-17) ó Potential Districts	42
2.8.6 National Scenario: Plan Periods (Kharif Pulses)	43
2.8.7 StatesøScenario: Plan Analysis (X th óXII th)	44
2.8.8 National Scenario: Plan Periods (Rabi Pulses)	47
2.8.9 StatesøScenario: Plan Analysis (X th óXII th)	48
Chapter – 3 : Chickpea (Gram)	51 to 67
3.1 Economic Importance	51
3.2 Nutritive Value	51
3.3 Growth Rate	52
3.4 Import/Export	53-54
3.5 Production Trends	55 to 67
3.5.1 Global Scenario	55
3.5.2 National Scenario : Plan Periods	56
3.5.3 Stateøs Scenario: Plan Analysis (X th óXII th)	57
3.5.4 District scenario (2016-17) ó Potential districts	60
3.6 Chickpea Agronomy	61 to
3.6.1 Economic Classification	61
3.6.2 Botanical Description	61
3.6.3 Production Technology	62-64
3.6.4 Plant Protection Measures	65-67
3.7 Recommendation to achieve higher production	67
Chapter -4 : Pigeonpea	79 to 102
4.1 Economic Importance	79
4.2 Nutritive Value	79
4.3 Agronomic Significance	79
4.4 Growth Rate	80
4.5 Import/Export	81
4.6 Production Trends	82 to 88
4.6.1 Global Scenario	82
4.6.2 National Scenario: Plan Period	84
4.6.3 StatesøScenario: Plan Analysis (X th óXII th)	85

CONTENT	PAGE NO.
4.7 Pigeonpea Agronomy	89
4.7.1 Economic Classification	89
4.7.2 Botanical Decription	89-90
4.7.3 Production Technology	90-93
4.7.4 Plant Protection Measures	94-96
4.8 Recommendation to achieved higher production	96
4.9 Hybrid Pigeonpea	97-99
4.9.1 Chronology of R&D work in development of hybrid pigeonpea	100 to
4.9.2 Scope	100
4.10 Situations/Season of Cultivation	100-102
4.10.1 Rabi Pigeonpea	101
4.10.2 Summer Pigeonpea	101
4.10.3 Production Technology	102
Chapter-5 : Greengram (Mungbean)	111 to 125
5.1 Economic Importance	111
5.2 Nutritive Value	111
5.3 Agronomic Importance	111
5.4 Growth Rate	111
5.5 Production Trends	113 to
5.5.1 National Scenario: Plan Periods	113
5.5.2 StatesøScenario: Plan Analysis (X th óXII th)	114
5.5.3 District Scenario (2016-17) ó Potential Districts	117
5.6 Mungbean Agronomy	118 to122
5.6.1 Botanical Description	118
5.6.2 Production Technology	119-122
5.7 Plant Protection Measures	122-125
5.8 Recommendation to achieved higher production	125
Chapter-6 : Black Gram (Urdbean)	133 to 145
6.1 Economic Importance:	133
6.2 Nutritive Value	133
6.3 Agronomic Significance	133
6.4 Growth Rate	133
6.5 Production Trends	135 to 139
6.5.1 National Scenario: Plan Periods	135
6.5.2 StatesøScenario: Plan Analysis (X th óXII th)	136
6.5.3 District Scenario (2016-17) ó Potential Districts	139
6.6 Urdbean Agronomy	140-145
6.6.1 Botanical Description	140

CONTENT	PAGE NO.
6.6.2 Production Technology	141-143
6.6.3 Plant Protection Measures	143-145
6.7 Recommendation to achieved higher production	145
Chapter -7 : Lentil	151 to 166
7.1 Economic Importance	151
7.2 Nutritive Value	151
7.3 Agronomic Significance	151
7.4 Growth rate of lentil	152
7.5 Import & Export	153-154
7.6 Production Trends	155 to
7.6.1 Global Scenario	155
7.6.2 National Scenario: Plan Periods	156
7.6.3 StatesøScenario: Plan Analysis (X th óXII th)	157
7.6.4 District Scenario (2016-17) ó Potential Districts	160
7.7 Lentil Agronomy	161-166
7.7.1 Botanical Description	161
7.7.2 Production Technology	162-164
7.7.3 Plant Protection Measure	164-166
7.8 Recommendation to achieved higher production	166
Chapter – 8 : Fieldpea	170 to 183
8.1 Economic Importance	170
8.2 Nutritive Value	170
8.3 Agronomic Significance	170
8.4 Growth Rate	170
8.5 Import/Export	172
8.6 Production Trends	173 to 178
8.6.1 Global Scenario: Fieldpea	173
8.6.2 National Scenario: Plan Periods	174
8.6.3 StatesøScenario: Plan Analysis (X th óXII th)	175
8.6.4 District Scenario (2016-17) ó Potential Districts	178
8.7 Fieldpea Agronomy	179 to 183
8.7.1 Botanical Description	179
8.7.2 Production Technology	179-181
8.7.3 Plant Protection Measures	181-182
8.8 Recommendation to achieved higher production	183

CONTENT	PAGE NO.
Chapter-9: Mothbean	187 to 194
9.1 Economic Importance	187
9.2 Agronomic Significance	187-188
9.3 Nutritive Value	188
9.4 Production Trends	188 to
9.4.1 StatesøScenario: Plan Analysis (X th óXII th)	188
9.4.2 District Scenario (2012-13)óPotential Districts	191
9.5 Mothbean Agronomy	192-194
9.5.1 Major Constraint in Production	192
9.5.2 Production Technology	192-193
9.5.3 Plant Protection Measures	194
9.6 Recommendation To Achieved Higher Production	194
Chapter-10 : Horsegram (Kulthi)	195 to 200
10.1 Economic Importance	195
10.2 Nutritive Value	195
10.3 Production Trends	195
10.3.1 StatesøScenario: Plan Analysis (X th óXII th)	195
10.3.2 District Scenario (2016-17) ó Potential Districts	198
10.4 Horsegram Agronomy	199 to 200
10.4.1 Botanical Description	199
10.4.2 Production Technology	199-200
10.4.3 Plant Protection Measures	200
10.5 Recommendation To Achieved Higher Production	200
Chapter-11 : Lathyrus (Khesari)	203 to 211
11.1 Economic Importance	203
11.2 Nutritive Value	203
11.3 Background: Policy on Consumption/Sale of Lathyrus	203
11.3.1 High power committee on lathyrus	204-205
11.4 Production Trend	206
11.4.1 StatesøScenario: Plan Analysis (X th óXII th)	206
11.4.2 District Scenario (2016-17) ó Potential Districts	208
11.5 Lathyrus Agronomy	209 to 210
11.5.1 Botanical Description	209
11.5.2 Production Technology	209-210
11.5.3 Plant Protection Measures	210
11.6 Recommendation to achieved higher production	211

CONTENT	PAGE NO.
Chapter-12: Rajmash	212 to 215
12.1 Economic Importance	212
12.2 Nutritive Value	212
12.3 Rajmash Agronomy	212
12.3.1 Botanical Description	212
12.3.2 Production Technology	212-214
12.3.3 Plant Protection Measures	214-215
12.4 Recommendation to achieved higher production	215
Chapter-13 : Cowpea	217 to 226
13.1 Economic Importance	217
13.2 Nutritive Value	217
13.3 Agronomic Importance	217
13.4 Crop Status	217-218
13.5 Cowpea Agronomy	218 to
13.5.1 Production Technology	218-220
13.5.2 Plant Protection Measures	220-222
13.6 Recommendation to achieve higher production	222
A. Broad bean	225
B. Rice-Bean	226
Chapter-14: Market Scenario: 2016-17	227 to 236
14.1 Production Scenario- 2016-17: An Analysis	227-228
14.2 Availability Status: Total Pulses & Crop-Wise (2013-14 To 2016-17)	231
14.3 Market Prices/Rates and Arrival	231
14.4 Factors attributing to lower market prices (Below MSP)	235-236
Chapter-15 : Seed Production	237 to 247
15.1 Importance	237
15.2 Class of Seed and their production	237-239
15.3 Seed Replacement:	239
15.4 Seed Requirement	240
15.5 Breeder seed production under new varieties	240-242
15.6 New Initiatives under NFSM-Pulses	243-247
15.6.1 Enhancing Breeder Seed Production	243-244
15.6.2 Creation of Seed-Hubs	245-246
15.6.3 Seed Village Programme	247
Chapter-16: Production and Sustainability Constraints Identified	250 to 263
16.1 Constraints related to production	250
16.2 Constraints related to inputs	255-256
16.3 Constraints related to marketing	259
16.4 Constraints related to extension and their interventions	259

CONTENT	PAGE NO.
16.5 Suggestions	259 to 263
16.5.1 Input Related Interventions	259-260
16.5.2 Production Related Intervention	260-261
16.5.3 Marketing Related Interventions	262
16.5.4 Extension Related Intervention	262-263
Chapter-17: Post-Harvest Technology and Management	
17.1Harvesting Precautions	264
17.2 Grading	264
17.3 Storage/Packaging	264
17.4Major Stored Grain Pests	265
17.5 Storage Structures	265
17.5.1 Storage Infrastructure/Programmes/Facilities	266-267
17.6 Marketing Channel	267
17.7 Processing and Value Addition	268
17.7.1 Domestic/Small Scale Pulse Milling in rural sector : Scope	268-269
17.7.2 Benefits of mini/small scale mills	269
17.8 Processing Technology	269
17.8.1 Large Scale Processing	270
17.8.2 Small Scale Processing	270-271
17.9 Domestic Machineries Developed	271
17.10 Milling methods of pulses	273 to 278
17.10.1 Traditional dry dal milling method	273
17.10.2 Milling techniques of different pulses	274-278
i) Dehulling of Tur (Arhar)	274
ii) Dehulling of Bengal Gram (Chickpea)	276
iii) Dehulling of Black Gram	277
iv) Dehulling of Green Gram	277-278
v) Dehulling of Peas, Lentil and Khesari	278
Chapter-18: Policy Interventions	
18.1 Projects/programme on pulses development	280-284
18.2 Performance over-view: (First to Twelfth Plan)	290 to 292
18.2.1 Area Expansion	290
18.2.2 Production Enhancement	291
18.2.3 Productivity	291
18.2.4 Irrigation	291-292
18.3 Strategies Adopted	292 to
18.3.1 Area expansion	292
18.3.2 Targeted rice fallow area (TRFA	293-296

CONTENT	PAGE NO.
18.3.3 Productivity Enhancement	298
18.3.4 Marketing Strategy	300
18.3.5 Farmer Producer Organization : Empowerment Through Group	302
Chpater-19: Vision 2030	305 to 318
19.1 Road Map-2030	305-318
19.2 Value Addition/Strategy Processing	305-307
19.3Strategy related to research issues	311-312
19.4 Policy Related Strategy	318
Chapter-20 : Way Forward-2030	319-327

LIST OF TABLES

TABLES	PAGE NO.
Executive Summary	
I. Horizontal Expansion	3
II. Production Performance	4
III. Additional Return through Vertical Expansion	4
Chapter -1: Pulses in Indian Context: 2017-18	11 to 22
1.1 Nutritional level of various pulses	11
1.2 Contribution of pulses to food grains basket	13
1.3 Crop contribution to total pulse production	14
1.4 StatesøContribution in Area & Production ó Total Pulses	15
1.5 StatesøContribution in Area & Productionó Kharif Pulses	16
1.6 States@Contribution in Area & Production- Rabi Pulses	17
1.7 States@Contribution in Area & Production- Gram	18
1.8 StatesøContribution in Area & Production ó Pigeonpea (2017-18)	19
1.9 StatesøContribution in Area & Production ó Mungbean (2017-18)	20
1.10 StatesøContribution in Area & Production- Urdbean (2017-18)	21
1.11 StatesøContribution in Area & Production-Lentil (2017-18)	22
Chapter-2 : Plan-wise Pulse Overview	26 to 49
2.1 Contribution of pulses to total foodgrains in India.	26
2.2 Season-wise Pulse Contribution to Total Pulses	27
2.3 Per capita availability of pulses in India	28
2.4 Demand, Production, Growth and Projected Target	29
2.5 (a) India@s imports and exports of pulses	31
2.5 (b) India@s imports and exports of pulses v/s agriculture	31
2.6 Pulse importing and exporting countries of major pulses (2016-17)	33
2.7 Total pulses: Crop/Season-wise contribution	34
2.8 Growth rate of total pulses	35
2.9 Global Ranking: Crop-wise	36
2.10 Global Ranking: Total Pulses	37
2.11 Plan-Wise National Scenario- Total Pulses	38
2.12 Plan-Wise StatesøScenarioóMajor States	40
2.13 Top Potential Districts (2016-17) óTotal Pulses	42
2.14 Plan-Wise National Scenario - Kharif Pulses	43
2.15 Plan-Wise StatesøScenario - Major States	45
2.16 Plan-Wise National Scenario ó Rabi Pulses	47
2.17 Plan-Wise StatesøScenarioóMajor States	49

TABLES	PAGE NO.
Chapter – 3 : Chickpea (Gram)	52 to 78
3.1 Growth rate of chickpea	52
3.2 Per cent share to total pulses	53
3.3 Importing & Exporting Countries	54
3.4 Global Ranking: Major Countries	55
3.5 Plan-Wise National Scenario	56
3.6 Plan-Wise StatesøScenario: Major States	58
3.7 Top Potential Districts: Chickpea (2016-17)	60
3.8 State-wise yield gap and additional return	61
3.9 Recommended varieties & their characteristics	68-78
Chapter -4 : Pigeonpea	80 to 110
4.1 Growth rate of pigeonpea	80
4.2 Per cent share to total pulses	81
4.3 Importing & Exporting Countries	82
4.4 Global Ranking: Major Counties	82
4.5 Plan-Wise National Scenario	84
4.6 Plan-Wise StatesøScenario: Major States	86
4.7 Top Potential Districts: Pigeonpea	88
4.8 State-wise yield gap and additional return	89
4.9 Pigeonpea production technology for different zones.	102
4.10 Agronomic recommendations of ICAR-IIPR for increased production under	103-104
different cropping systems in the country	
4.11 Recommended Varieties/Characteristics	105-110
Chapter-5 : Greengram (Mungbean)	112 to 132
5.1 Growth rate of mungbean	112
5.2 Per cent share to total pulses	112-113
5.3 Plan-wise national scenario	113
5.4 Plan-Wise StatesøScenario ó Major Stateøs	115
5.5 Top potential districts: Mungbean	117
5.6 State-wise yield gap and additional return	118
5.7 Recommended mungbean varieties/characteristics	126-132
Chapter-6 : Black Gram (Urdbean)	113 to 150
6.1 Growth rate of urdbean	133-134
6.2 Per cent share to total pulses	134
6.3 Plan-Wise National Scenario ó Urdbean	135
6.4 Plan-Wise StatesøScenario ó Major Stateøs	137
6.5 Top Potential Districts: Urdbean	139
6.6 State-wise yield gap and additional return: urdbean	140
6.7 Recommended urdbean varieties/characteristics	146-150

TABLES	PAGE NO.
Chapter -7 : Lentil	152 to 169
7.1 Growth rate of lentil	152
7.2 Per cent share to total pulses	153
7.3 Importing & Exporting countries	154
7.4 Global Ranking: Major Countries	155
7.5 Plan-wise national scenario	156
7.6 Plan-wise statesøscenario ó major states	158
7.7 Top Potential Districts: Lentil (2016-17)	160
7.8 State-wise yield gap and additional return	161
7.9 Recommended varieties/characteristics	167-169
Chapter – 8 : Fieldpea	170 to 186
8.1 Growth rate of fieldpea	170-171
8.2 Per Cent share to total pulses	171
8.3 Importing & exporting countries	172
8.4 Global Ranking : Major Countries	173
8.5 Plan-Wise National Scenario	174
8.6 Plan-Wise StatesøScenario - Major Stateøs	176
8.7 Top Potential Districts: Fieldpea (2016-17)	178
8.8 State-wise yield gap and additional return	179
8.9 Recommended varieties of fieldpeas/characteristics	184-186
Chapter-9: Mothbean	189 to 194
9.1 Plan-Wise StatesøScenario óMajor Stateøs	189
9.2 Top Potential Districts (2012-13)	191
9.3 Recommended varieties of mothbean/characteristics	194
Chapter-10 : Horsegram (Kulthi)	196 to 202
10.1 Plan-Wise StatesøScenario - Major Stateøs	196
10.2 Top Potential Districts: Horsegram (2016-17)	197
10.3 Recommended Varieties/Characteristics	201-202
Chapter-11 : Lathyrus (Khesari)	206 to 211
11.1 Plan-Wise StatesøScenario	206
11.2 Top Potential Districts: Lathyrus (2016-17)	208
11.3 Recommended Varieties/Characteristics	211
Chapter-12: Rajmash	216
12.1 Recommended Varieties/Characteristics	216
Chapter-13: Cowpea	222 to 224
13.1 Recommended varieties of cowpea/characteristics	222-224

TABLES	PAGE NO.
Chapter-14: Market Scenario: 2016-17	228 to 234
14.1 National production of pigeonpea, urd, mung & lentil	228-230
I. State-wise area production of pigeonpea	228
II. State-wise area production of <i>urdbean</i>	229
III. State-wise area production of mungbean	229
IV. State-wise area production of gram	230
V. State-wise area production of lentil	230
14.2 Import, Export and Availability	231
14.3 Market Prices/Rates and Arrivals	232-234
I. State-wise market rates and arrivals of pigeonpea	232
II. State-wise market rates and arrivals of urdbean	232
III. State-wise market rates and arrivals of mungbean	233
IV. State-wise market rates and arrivals of lentil	233
V. State-wise market arrivals of gram	234
VI. State-wise market arrivals and prices of gram	234
Chapter -15 : Seed Production	239 to 249
15.1 Crop-wise & Year-wise SRR	239
15.2 Seed Requirement	240
15.3 Breeder seed production and indent	240
15.4 Certified seed production programme under NFSM-Pulses	242
15.5 All India: Crop-wise requirement and availability of certified seed	243
15.6 All India-crop-wise additional breeder seed production targ. & achiev.	244
15.7 Infrastructure: Strengthening of seed production farms	245
15.8 All India-Crop-wise seed production target and achiev. under seed-hub	246
15.9 Seed distribution under Seed Village Programme (2017-18)	247
15.10 State-wise Physical and Financial Progress under Seed Village Programme	248
15.11 Crop-wise Seeds distributed and produced under SVP during 2017-18	249
Chapter-16: Production and Sustainability Constraints Identified	250 to 260
16.1 Technological yield gap exhibiting the production related constraints- FLDs	250
16.2 Yield gap exhibiting the production related constraints among the states	251
16.3 Identified production related constraints and their interventions	252
16.4 All India CFLDs targets and achievement (2015-16 to 2018-19)	253
16.5 All India: CFLDs yield gaps exhibited: Total Pulses (2016-17)	254
16.6 All India: CFLDs yield gaps: Kharif Pulses (2016-17)	254
16.7 All India: CFLDs yield gaps: Rabi Pulses (2016-17)	254
16.8 All India: Varieties Demonstrated under CFLDs	255
16.9 Cluster FLD on Pulses 2017-18	255
16.10 Requirement and availability of certified seeds during 2016-17	256
16.11 Poor Varietal Development (XI th and XII th Plan)	256

TABLES	PAGE NO.
16.12 Limited statesøvarietal choice during last 10 years (2006-2015)	257
16.13 Poor Varietal Diversification (VRR)	257-258
16.14 Crop-Wise Potential Districts With 20-30% Prod. ShareóAll India	260
Chapter-17: Post-Harvest Technology and Management	265 to 272
17.1 Categories of prevalent storage structures.	265
17.2 Processes of marketing of raw produce.	267
17.3 Processing technology developed under R&D in PHT	271-272
Chapter-18: Policy Interventions	284 to 303
18.1 Plan-Wise Intervention (VIII th to XII th Plan)	284
18.2 Recent policy initiatives/interventions taken (2015-16 to 2017-18)	285-287
18.3 Interventions under NFSM-Pulses	288
18.4 Summary of research project funded under of NFSM-Pulses in year 2017-18	289
18.5 Plan-wise trend of growth in pulses	290
18.6 Area coverage under spring/summer pulses	292
18.7 Progress of TRFA	294
18.8 State-wise production of pulses under TRFA during 2017-18	294
18.9 (a) Major technological interventions	297
18.9 (b) Performance based recommendation	297
18.10 Promotion of improved varieties	299
18.11 Crop-wise procurement of pulses enhanced	300
18.12 State-wise procurement of pulses under MSP (PSS)	301
18.13 Crop-wise procurement of pulses under MSP (PSF)	302
18.14 State-wise progress of FPO promotion (As on 30.06.2018)	303
Chpater-19: Vision 2030	305 to 317
19.1 Potential of additional area under pulses	305
19.2 Seed Plan-Total Pulses (2018-19 to 2020-21)	307
19.3 Seed Plan: Kharif Pulses (2018-19 to 2020-21)	308
19.4 Seed Plan: Rabi Pulses (2018-19 to 2020-21)	309
19.5 Crop Specific Strategy/Recommendations (var./ plant protection)	313-317

LIST OF DIAGRAM

DIAGRAM	PAGE NO.
I. Horizontal Expansion	3
II. Production Performance	4
III. Additional Return through Vertical Expansion	5
Chapter -1: Pulses in Indian Context: 2017-18	13 to 22
1.1 Contribution of Pulses to Foodgrains Basket	13
1.2 Crop contribution in Total Pulses	14
1.3 StatesøContribution in Area & Production ó Total Pulses	15
1.4 StatesøContribution in Area & Productionó Kharif Pulses	16
1.5 StatesøContribution in Area & Productionó Rabi Pulses	17
1.6 StatesøContribution in Area & ProductionóGram	18
1.7 StatesøContribution in Area & Production ó Arhar/Tur	19
1.8 State® Contribution in Area & Production-Mungbean	20
1.9 StatesøContribution in Area & Production- Urdbean	21
1.10 StatesøContribution in Area & Production-Lentil	22
Chapter-2 : Plan-wise Pulse Overview	27 to 50
2.1 Contribution of Pulses to Foodgrains Basket	27
2.2 Pulses Availability	29
2.3 Demand vs Production	30
2.4 Import & Export of Pulses	33
2.5 Global Ranking: Crop-wise	36
2.6 Global Ranking: Total Pulses	37
2.7 Plan-Wise National Scenario- Total Pulses	38
2.8 Total Pulses Area: State@s Contribution (XII th Plan)	39
2.9 Total Pulses Production: State® Contribution (XII th Plan)	39
2.10 Total Pulses: Potential District	42
2.11 Kharif Pulses: Plan-Wise Scenario	43
2.12 Kharif Pulses Area: State Contribution (XII th Plan)	46
2.13 Kharif Pulses Production: State@s Contribution (XII th Plan)	46
2.14 Rabi Pulses: Plan-Wise Scenario	47
2.15 Rabi Pulses Area: State@s Contribution (XII th Plan)	50
2.16 Rabi Pulses Production: State Contribution (XII th Plan)	50
Chapter – 3 : Chickpea (Gram)	55 to 60
3.1 Global Scenario : Chickpea	55
3.2 Chickpea: Plan-Wise Scenario	56
3.3 Chickpea Area: State State Contribution (XII th Plan)	59
3.4 Chickpea Production: State & Contribution (XII th Plan)	59
3.5 Chickpea: Potential Districts	60

DIAGRAM	PAGE NO.
Chapter -4 : Pigeonpea	83 to 88
4.1 Global Area : Pigeonpea	83
4.2 Global Production: Pigeonpea	83
4.3 Pigeonpea : Plan-wise Scenario	84
4.4 Pigeonpea Area: State@s Contribution (XII th Plan)	87
4.5 Pigeonpea Production: State® Contribution (XII th Plan)	87
4.6 Pigeonpea: Potential District	88
Chapter-5 : Greengram (Mungbean)	113 to 117
5.1 Mungbean: Plan-wise Scenario	113
5.2 Mungbean Area: State@s Contribution (XII th Plan)	116
5.3 Mungbean Production: State@s Contribution (XII th Plan)	116
5.4 Mungbean: Potential District	117
Chapter-6 : Black Gram (Urdbean)	135 to 139
6.1 Urdbean: Plan-wise Scenario	135
6.2 Urdbean Area: State@s Contribution (XII th Plan)	138
6.3 Urdbean Production: State® Contribution (XII th Plan)	138
6.4 Urdbean: Potential District	139
Chapter -7: Lentil	155 to 160
7.1 Global Area and Production : Lentil	155
7.2 Lentil : Plan-wise Scenario	156
7.3 Lentil Area: State@s Contribution (XII th Plan)	159
7.4 Lentil Production: State@s Contribution (XII th Plan)	159
7.5 Lentil: Potential District	160
Chapter – 8 : Fieldpea	173 to 178
8.1 Global Area and Production : Fieldpea	173
8.2 Fieldpea : Plan-wise Scenario	174
8.3 Pigeonpea Area: State@s Contribution (XII th Plan)	177
8.4 Pigeonpea Production: State@s Contribution (XII th Plan)	177
8.5 Fieldpea: Potential District	178
Chapter-9: Mothbean	
9.1 Mothbean Area: State@s Contribution (XII th Plan)	190
9.2 Mothbean Production: State® Contribution (XII th Plan)	190
9.3 Mothbean: Potential District	191
Chapter-10 : Horsegram (Kulthi)	197 to 198
10.1 Horsegram Area: State@s Contribution (XII th Plan)	197
10.2 Horsegram Production: State® Contribution (XII th Plan)	197
10.3 Horsegram: Potential District	198

DIAGRAM	PAGE NO.
Chapter-11 : Lathyrus (Khesari)	207 to 208
11.1 Lathyrus Area: Stateøs Contribution (XII th Plan)	207
11.2 Lathyrus Production: State® Contribution (XII th Plan)	207
11.3 Lathyrus: Potential District	208
Chapter-15: Seed Production	239 to 249
15.1 Seed Replacement Rate	239
15.2 Breeder Seed Production vs Indent: Total Pulses	241
15.3 Breeder Seed Production: Kharif Pulses	241
15.4 Breeder Seed Production: Rabi Pulses	241
15.5 Availability of Certified Seeds of Pulses Enhanced	242
15.6 Availability of Certified Seed	243
15.7 Production of Breeder seed under ABSP	244
15.8 Production of quality seed under Seed-hub	246
15.9 Seed Village Programme - Phy. & Fin. Progress	248
15.10 Crop-wise Seed Distributed - 2017-18	249
Chapter-16: Production and Sustainability Constraints Identified	253
16.1 All India CFLDs Targets and Achievement (2015-16 to 2018-19)	253
Chapter-17: Post-Harvest Technology and Management	274 to 279
17.1 Flow diagram of dry milling of pulses	274
17.2 Dehulling of tur-large scale	275
17.3 Dehulling of tur- dry method	275
17.4 Dehulling of tur- wet method	276
17.5 Process for dehulling of Bengal gram	276
17.6 Process for milling green gram and blackgram	277
17.7 Green gram/black gram splits-flow chart	278
17.8 Process for dehulling peas	278
17.9 Process for milling of masoor (lentil)	279
17.10 Process for dehulling khesari pulse (lakh)	279
Chapter-18: Policy Interventions	
18.1 Area Coverage under Total Summer Pulse	293
18.2 Major state-wise Area Coverage under Spring/Summer Pulses	293
18.3 Quantity Procured under MSP	300
18.4 Values of Procured Pulses under MSP	301
18.5 State-wise Procurement of Pulses under MSP (PSS)	301
18.6 Crop-wise procurement of pulses under MSP (PSF)	302
Chpater-19: Vision 2030	
19.1 State-wise seed plan total pulses	308
19.2 State-wise seed plan kharif pulses	309
19.3 State-wise seed plan rabi pulses	310
19.4 Season-wise seed plan of pulses	310
19.5 Crop-wise seed plan of pulses	311

LIST OF ANNEXURE

ANNEXURE	PAGE NO.
I. Approved Norms of NFSM óPulses for 2017-18	338
II. District Covered (Identified) under NFSM - Pulses	339
III. Pulses: All India- Crop Calender: Package of Practice	340-355
IV. Specific standards prescribed for certification at field stage for pulses	356
V. Seed Standard	357
VI. Crop-wise and State-wise categories of district having LALP, LAHP, HALP, HAHP	358-379
VII. Crop-wise and State-wise districts of higher and lower National Yield.	380-392
VIII. Crop-wise major Potential Pulse State/Districts for the proposed pulse export	393-395
zone	
IX. Contact Details of DAC&FW, CDD, SDAs etc.	396-406

Executive Summary

1. A giant leap in production

- During 2017-18, pulses were cultivated in approx. 29.99 million ha (Mha) area and recorded the highest ever production of 25.23 million tonnes (Mt) at a productivity level of 841 kg/ha. The exponential growth rate in production of pulses during last year was 9.11 *per cent*.
- Major 10 states to contribute > 90 per cent pulse production have been Madhya Pradesh (> 8 Mt), Rajasthan (>3 Mt), Maharashtra (>3 Mt) Uttar Pradesh (>2 Mt) Karnataka (2 Mt) and Andhra Pradesh (>1 Mt) followed by Gujarat, Jharkhand, Tamil Nadu, and Chhattisgarh producing <1.0 Mt each.
- Under individual crop category, Gram recorded a highest ever production of 11.23 Mt at a record productivity level of 1063 kg/ha in an area of 10.56 Mha. Major 07 states to contribute > 90 per cent in gram production have been Madhya Pradesh (4.60 Mt), Maharashtra (1.78 Mt), Rajasthan (1.67 Mt), Karnataka (0.72 Mt), Andhra Pradesh (0.59 Mt), Uttar Pradesh (0.58 Mt) and Gujarat (0.37 Mt).
- Tur (Arhar) remained at IInd position in total pulse production with 4.25 Mt of production in an area of 4.43 Mha at a productivity level of 960 kg/ha, the ever highest yield. Major states to record the success have been Maharashtra (1.07 Mt), Madhya Pradesh (0. 84 Mt), Karnataka (0.77 Mt), Gujarat (0.34 Mt), Uttar Pradesh (0.33 Mt), Telangana (0.26 Mt) and Jharkhand (0.22 Mt).
- Urd (Blackgram), the IIIrd important crop group, was cultivated in an area of 5.44 Mha (kharif + rabi) and recorded a production of 3.56 Mt at a productivity level of 655 kg/ha. This was the *highest ever area, production and productivity* in this crop. Major contributing states have been MP, Rajasthan, AP, UP, Tamil Nadu, Maharashtra, Jharkhand and Gujarat.
- Similarly, Mung (Greengram) was sown in an area of 4.26 Mha in (kharif + rabi) and recorded a production of 2.01 Mt at yield level of 472 kg/ha. Rajasthan, MP, Maharashtra, Karnataka, Bihar, AP, Odisha, Tamil Nadu, Gujarat and Telangana have been the major states.
- Lentil also recorded an ever highest production of 1.61 Mt in an area of 1.55 Mha at a productivity level of 1034 kg/ha, *the ever highest yield level*. Leading six lentil producing states have been Madhya Pradesh (0.68 Mt), Uttar Pradesh (0.50 Mt), West Bengal (0.15 Mt), Bihar (0.14 Mt), Jharkhand (0.06 Mt) and Rajasthan (0.03 Mt).

2. Equilibrium Production and demand

- As per the Report of the Working Group on Crop Husbandry, Demand & Supply Projection for XIIth Plan (2012-17) of NITI Aayog (erstwhile Planning Commission), the demand of pulses @ 3.09 *per cent* per annum growth during 2016-17 and 2017-18 has been worked out at 22.74 Mt and 23.44 Mt respectively.
- It was for the first time since plan 6 interventions on pulses that the nation inscribed a success by achieving higher pulse production at 23.13 Mt and 25.24 Mt during 2016-17 and 2017-18, meet out the demand.
- Country witnessed the self-sufficiency in pulses.

3. Significant growth with additional return

- During 2017-18, significant growth registered under total pulse production, both over the base year (2014-15) and the normal/XIIth Plan (2012-13 to 2016-17) at the level of 47 *per cent* and 34 *per cent* respectively.
- Major increment recorded in kharif production *i.e.* 62 *per cent* mainly due to lion share contributed by urad (82 *per cent*) followed by tur (52 *per cent*) and mung (34 *per cent*).
- Rabi Pulses recorded a 39 *per cent* hike over the base year (2014-15) and mainly contributed by gram (53 *per cent*) and lentil (55 *per cent*).
- As a consequence of meticulous planning and strategy, the production increase was
 realized both by way of horizontal expansion *i.e.* pulses under new niches as well as the
 vertical expansion by bridging the yield gaps through aggressive technology transfer,
 capacity building, workshops/seminars and consultations with the stake-holders during
 the aforesaid period.
- A total pulse area was increased to about 27 *per cent* over the period. Under seasonal increase, it was 41 *per cent* during kharif followed by rabi 17 *per cent*. Under individual crop category, major increment in area has been recorded in urad (68 *per cent*) followed by Mung (41 *per cent*), gram (28 *per cent*) and tur (15 *per cent*).
- The total pulse productivity increase during the same period has been about 15 *per cent* over the base year (2014-15), comprising rabi and kharif increment at about 19 *per cent* and 15 *per cent* respectively.
- As regards the crop-wise yield performance during 2017-18 over the period, the highest yield increment have been recorded in lentil (47 per cent) followed by tur (32 per cent), gram (20 per cent) and urd (8 per cent) over the base year (2014-15).

- The change over under Area Expansion (horizontal expansion) and production growth over normal and base year are indicated under *Table I & II*.
- The overall pulse change over per ha yield (vertical expansion) and consequently the per unit additional return to farmer to increase their income is about Rs. 6487 *per ha* and Rs. 5308 *per ha* over the base year and normal respectively. The crop-wise details indicated under *Table-III*.

(Table-I): Horizontal Expansion

(Area-Million ha)

Crop	Season		Area	Change over (+/-)		
		Normal*	2014-15	2017-18**	Normal	2014-15
Tur	Kharif	4.19	3.85	4.43	0.24	0.58
Urd	Kharif	2.70	2.48	4.50	1.80	2.02
	Rabi	0.81	0.76	0.94	0.13	0.18
	Total	3.51	3.25	5.44	1.93	2.19
Mung	Kharif	2.49	2.02	3.29	0.80	1.27
	Rabi	0.96	0.99	0.97	0.01	-0.02
	Total	3.46	3.02	4.26	0.80	1.24
Gram	Rabi	8.95	8.25	10.56	1.61	2.31
Lentil	Rabi	1.39	1.47	1.55	0.16	0.08
Other	Kharif	1.81	1.63	1.87	0.06	0.24
Pulses	Rabi	1.97	3.55	1.88	-0.09	-1.67
	Total	3.78	5.19	3.75	-0.03	-1.44
Kharif P	Charif Pulses 11.19 10.00 14.08 2.89		2.89	4.08		
Rabi Pu	lses	14.08	13.56	15.91	1.83	2.35
Total Pu	lses	25.28	23.55	29.99	4.71	6.44

Source: DES, Ministry of Agriculture & FW (DAC&FW), GoI Normal*-Avg. 2012-13 to 2016-17; 2017-18**- IVth Advance Estimate.




Fig.-I: Horizontal Expansion

(Table-II): Production Performance

(Production-Million Tones)

Crop	Production			Change over (+/-)		
	Normal*	2014-15	2017-18**	Normal*	2014-15	
Tur	3.29	2.81	4.25	0.96	1.44	
Urd	2.08	1.96	3.56	1.48	1.60	
Mung	1.61	1.50	2.01	0.40	0.51	
Gram	8.43	7.33	11.23	2.80	3.90	
Lentil	1.08	1.04	1.61	0.53	0.57	
Kharif Pulses	6.55	5.78	9.34	2.79	3.56	
Rabi Pulses	12.29	11.42	15.89	3.60	4.47	
Total Pulses	18.84	17.20	25.23	6.39	8.03	

Source: DES, Ministry of Agriculture &FW (DAC&FW), GoI Normal*-Avg. 2012-13 to 2016-17; 2017-18**- IVth Advance Estimate.




Fig.-II: Production Performance

(Table-III): Additional Return through Vertical Expansion

(Yield-kg/ha, Additional Return-Rs/ha))

Crop	Yield			Yield Gap		Addition Return	
	Normal*	2014-15	2017-18**	Normal*	2014-15	Normal	2014-15
Tur	785	729	960	175	231	8284	11336
Urd	593	604	654	61	50	3240	2646
Mung	466	498	472	6	-26	56	-1728
Gram	942	889	1063	121	174	4972	7304
Lentil	773	705	1039	266	334	9987.5	12878
Kharif Pulses	585	578	663	78	85		
Rabi Pulses	873	843	999	126	156		
Total Pulses	745	730	841	96	111		

Source: DES, Ministry of Agriculture &FW (DAC&FW), GoI Normal*-Avg. 2012-13 to 2016-17; 2017-18**- IVth Advance Estimate.




Fig.-III: Additional Return through Vertical Expansion

4. Pulses import falls by 03 million tonnes in financial year- 2018 (Saving of foreign currency)

- With the increase in population, consumer awareness and affordability of middle /lower middle and other category citizens up to some degree, the demand of pulses has also increased overtime.
- Despite biggest producer, India is also the biggest importer and consumer (23-24 million tonnes) of pulses in the world.
- The year 2014-15 and 2015-16 were the adverse crop year owing to drought and erratic behavior of rainfall across the major pulse growing states.
- The government, however ensured the availability/supply as per demand by way of enhanced imports between 2014-15 to 2016-17 at about 5-6 million tonnes (Mt) per year in their buffer stock on one hand and swung in to action to combat the natural calamities through development programmes, risk management through PMFBY, PSS and PSF procurement etc., on the other.
- The CCEA has empowered the Committee headed by Food Secretary to review the export and import policy on pulses and consider measures such as quantitative restrictions, prior registration and changes in import duties depending on domestic production and demand, local and international prices and global trade volumes.

- Farmer-friendly policy measures have helped to reduce import of pulses. Import of pulses during 2017-18 has declined by about 30 lakh tonnes from previous year, resulting in saving of foreign exchange amounting to Rs 7,698 crore. It is expected that pulses production will be sustained in the country and our import dependence on pulses will come down substantially.
- To ensure that farmers get remunerative prices, the government has imposed import duty and put quantitative restrictions on the various varieties of pulses. Import duty on chickpea has been fixed at 60 per cent, while that for yellow pea is 50 per cent, lentil 30 per cent and tur 10 per cent on 21st Aug., 2017.

5. Export of pulses allowed

- Further, to safeguard farmersø interest, recently the Cabinet Committee on Economic Affairs (CCEA) has given its approval for removal of prohibition on export of all types of pulses to ensure that farmers have greater choice in marketing their produce and in getting better remuneration for their produce.
- The decision comes two months after the government lifted ban dated 15th September 2017 on export of tur, urad and moong dal, although shipments of these varieties were allowed only through permission from agriculture export promotion body APEDA. Export of organic pulses and kabuli chana is permitted in a limited quantity.
- Opening of exports of all types of pulses will help the farmers dispose of their products at remunerative prices and encourage them to expand the area of sowing.

6. Interventions which proved the trigger

- To ensure availability of location specific/recommended high yielding varieties and quality certified seeds at all levels, 150 Pulse Seed Hubs and strengthening of Infrastructure for Enhancing Breeder Seed Production (EBSP-12 Centres) programmes initiated from 2016-17 to 2018-19. The total fund flow for seed hub & EBSP has been Rs. 245.39 crore.
- Incentives for production of certified seeds and seed distribution of pulses were instrumental both in varietal replacement as well as area expansion.
- To ensure availability of quality bio-inputs-rhizobium culture/PSB, micro-nutrients, bio-intensive/bio-pesticides etc.
- To reduce cost of cultivation and timely operations in rainfed areas where > 80% pulses are grown, availability of implements like seed drills, zero-till seed machine/rotavator and ridge-maker etc., ensured through Resource Conservation Technology (RCT)

- components and Custom Hiring Centres (CHCs), especially in Bundelkhand region of U.P./M.P yielded the result.
- In view of favorable response of pulses to 1-2 critical irrigations, priority was given to pulses in tune MIDH/Micro-irrigation scheme under PMKSY.
- For processing and value addition, domestic milling support provided through mini dal mills under *local initiative/flexi fund component* various states including UP., Gujarat, and Maharashtra.
- To ensure effective transfer of technology, Cropping System Based Trainings (CSBTs) were provided to extension workers. Quality cluster demonstrations, both on sole crop and CSBDs were organized which helped in bridging the yield gaps.
- Strong Interface mechanism between State Department of Agriculture and State Agricultural Universities (SAUs), ICAR and KVKs were developed through seminars/workshops/Annual Group Meetings of ICAR etc. These activities were systematically organized during the International Year of Pulses (IYOP-2016).
- Robust monitoring and field visits in all 638 NFSM districts across the country and comprehensive approach in implementation of programme, including all components from seed to post-harvest management, marketing aspects and capacity building etc., yielded wonderful results.
- Dissemination of information through Literature on Pulses (bulletin/leaflet and articles) in both languages by print media as well as digital (dpd.gov.in/Farmers portal/mKisan Portal) including advisories on pulses on monthly basis by the Department including Directorate of Pulses Development, Bhopal.
- In 2015-16, All XI ó ATARIøs/578 KVKs were involved in conducting of Cluster Front Line Demonstrations (CFLDs) of pulses with need based thematic areas on farmersø field. An amount of Rs. 63.38 crore has so far been utilized for ToT through CFLDs and 2.95 crore under regular ICAR-IIPR FLDs.
- In 2016-17 Seed minikits programme of newer varieties of pulses was initiated to popularize improved varieties of pulses. An amount of Rs. 137 crore has so far been utilized and 76.71 crore allocated for 2018-19.
- Applied and Action-Research Projects to ICRISAT, ICARDA, ICAR/SAUs to address biotic and abiotic stress/assessment and providing varieties /recommendations.

Conclusion

- Ever since the government resumed the office in May 2014, it resolved the Union government commitment to achieve/sustain the ambitious twin objectives of food and nutritional security and raising agricultural productivity to making farming remunerative to double the farmer income by 2022.
- Pulses programme were kept to centre stage owing to their nutritional importance to contain the protein malnutrition vis-à-vis their role in the economy of >40 per cent human population and 2/3rd live-stock rearing in the rainfed regions of the country. About 80 % of total pulse area fall in rainfed region.
- To enhance the production and productivity of pulses, in addition to enhanced financial provision under NFSM, the DAC&FW encouraged/propagated adaptation of multi-dimensional seven point strategy suggested by the Honøble Prime Minister, which include: Emphasis on irrigation along with end to end solution on creation of resources for more crops per drop; Provision of quality seeds and nutrients according to the soil quality of each farm; Large investment in warehouse and cold chains to prevent post harvest losses; Promotion of value addition through food processing; Implementation of National Agricultural Markets and e- platform (e-NAM) to eliminate shortcomings of all the 585 centers.; To mitigate the risk by introduction of crop insurance scheme (PMFBY) at a lower cost.; Promotion of allied activities such as *Dairy, Animal Husbandry; Poultry, Bee-keeping, Har med per ped, Horticulture & fisheries*.
- Despite two consecutive droughts/erratic monsoon years between 2014-15 to 2015-16, sustained efforts by both the centre and states, the Indian farmers created a history by a record production of 23.13 Mt in 29.45 Mha at 786 kg/ha yields and 25.24 Mt in 29.99 Mha at the productivity level of 841 kg/ha during 2016-17 and 2017-18, respectively. Record area coverage under pulse crops and productivity is a significant outcome of the recent policies of the government.
- Appropriate policy support, timely availability of quality inputs, enhanced MSP, curbing distress sale through procurement by NAFED/SFAC/FCI, pro-active statesø policies such as marketing under Bhavantar Bhugtan Yojana in MP, Bonus/incentives in Chhattisgarh (@ Rs 1500 per acres) as provision for buffer stocks, imposition of import duties on pulses (Peas-50% from Nov. 8th, 2017; gram-40% w.e.f. Feb 06th, 2018 revised to 60% w.e.f. from March 2018; lentil-30% w.e.f. Dec 21st, 2017; Tur 10% w.e.f. March 28th, 2017), risk management under PMFBY coupled with the implementation of district crop contingency plans (DCCPs) involving pulses etc., were instrumental to provide cushioning to the success.

- Seed Minikits of newer varieties; Creation of Pulses Seed Hubs/ Infrastructures for production of quality seeds; Enhancing Breeder Seed Production (EBSP) at 12 centres in 8 states; Cluster FLDs through ICAR/ATARI/578 KVKs; National Workshops on õPromotion of Pulses in non-traditional Niches; Summer Cultivationö and capacity building activities; Targeted INM-NFSM; Seed production; Strengthening of Bioagent/Bio Control labs (24 Centres SAUs/ICAR); National Demonstration Project & Value Chain Development of Pulses & Millets (111 FPOs), etc., have been the new initiatives taken for the first time.
- Area expansion strategy under RKVY/NFSM as Targeting Rice fallow Areas (TRFA) in 6 eastern states in about 3 Mha; Additional area coverage in spring/Summer intercropping of pulses, tur on rice bunds and transplanting of tur (Dharwad system) Crop Diversification Programmes (CDP) and changing cropping pattern strategies between 2015-16 to 2017-18, paid the dividends. An area of >29 Mha could be stabilized for consecutive two years of 2016-17 & 2017-18.
- Massive programme of pulse seed minikits, production incentives to farmers/seed societies and public sector seed production agencies had the requisite impetus in record production of pulses.
- Decentralized planning of the government also prompted the states to fetch further funding support on Pulses development programmes. AP initiated the procurement and supplied minikits with 75% subsidy and promoted intercropping of redgram in 4.8 million hectare with groundnut and cotton; Jharkhand, Karnataka, Odisha, Telangana, UP and Maharashtra distributed mini dal mills to support processing & value addition.
- Mechanization, line sowing, strengthening of irrigation infrastructure under RKVY and NFSM in convergence with MIS (drip/sprinkler/water carrying pipes) etc., played a critical role in harvest of higher yields. Critical irrigation support was provided through NFSM and PMKSY and 20 per cent of total pulse area brought in the ambit of irrigation.
- Large scale technological demonstrations through cluster frontline demonstrations (CFLDs) by ICAR/ATARI in thematic areas (line sowing, BBF, INM, IPM, Seed treatment FIR etc.,) and also by ICAR-IIPR, AICRP centers coupled with capacity building/training activities under CSBT, could be proved an effective strategy.
- Promising state specific varieties were promoted along-with adoption of good agricultural practices and they played a major role, these include:

- The high yielding varieties GNG 1581, JG 11, JG 63, JG 130, JG 14, JAKI 9218 and Vijay in chickpea; BDN 711, TJT 501, BSMR 736, Maruti, Bahar, Narendra Arhar-1, Asha in pigeonpea; MH 421, SML 668, IMP 2-3, GM-4, HUM 16, IPL 2-14, Pant Mung 5 and PDM 139 in mungbean; PU 30, PU 31, IPU 02-43, KU 96-3, TAU-1, LBG-752, Uttara and KU-300 of urdbean; Pant L-8, Pant L-7, Pant L-6, JL-3, HUL 57, WBL 77 and K 75 and HUDP 15, Prakash, KPMR-400, KPMR 522, Vikas, Aman and Adarsh under Field pea significantly contributed to harvest the highest production.
- The achievements not with-standing, there still remain a large untapped yield potential reservoir for the exploitation of which mission has to step-up its efforts further. A gap between the potential yield that can be achieved at farmerøs field level and what they actually get is very wide. Bridging this yield gap offers an opportunity to produce more even by using the available technologies.
- To meet the domestic demand of pulse requirement and ensure self-sufficiency in pulses, a sustainable production and productivity has to be maintained with the multipronged short-term and long-term strategies. As Agriculture growth is limited, imports may help improve the supply situation in the short term whereas, the long term measures would need to focus on sustainable production system with increased productivity envisaging public capital formation in irrigation, quality seeds of promising varieties and their availability at least 33% SRR, research and efficient use of water, plant nutrition and other necessary inputs including the remunerative prices to the farmers.
- Policy initiatives must lead for efficiency and helping maintaining balance between domestic production and demand. If we strive to achieve these potential yield levels, than the increasing demand requirement of the country can be met in future.
- It is learnt that the success is likely to remain sustainable owing to 100 per cent implementation of Direct Benefit Transfer Mode (DBT) under Crops Development Programme (NFSM-Pulses). The assistance towards critical inputs was ensured in the accounts of the beneficiaries belonging to SCP/TSP/Woman /SMF Categories. A large section of farmers/pulse growers are happy with the DBT and the assistance provided is being utilized in real sense.
- During the last 03 years of programme implementation *i.e.* 2016-17 to 2018-19 there has been substantially growth in production and productivity of pulses with the support and extension work taken-up by both central and state government. The results indicated that the programme is suitable and could be highly viable economic programme for the benefits of farmers as well as country.

Chapter -1

Pulses in Indian Context: 2017-18

1.1 Background

- 1.1.1 The Food Security Act-2013 mandatorily envisages the right to nutritional security as well. To ensure access to adequate quantity of quality food at affordable prices to each individual, as per FSA-2013, is government to priority.
- 1.1.2 Pulses are important commodity group of food crops that can play a vital role to address national food and nutritional security and tackle environmental challenges. Pulses share to total foodgrain basket is around 9-10 *per cent* and critical and inexpensive source of plant-based proteins, vitamins and minerals. Pulses are a Smart Food as these are critical for food basket (dal-roti, dal-chawal), rich source of protein *i.e.* 20-25 *per cent* which is double the protein content of wheat and thrice that of rice and help address obesity, diabetes malnutrition etc. Pulse-wise nutritional status is given in (*Table 1.1*).

(Table–1.1): Nutritional level of various pulses

(Unit-mg/100 g)

Name of foodstuff	Gram	Urd	Mung	Kulthi	Lentil	Pea	Tur	Moth	Khesari	Cowpea
Protein (%)	20	24	25	22	25	22	22	25	31	23
Vit. A (I.U.)	316	64	83	119	450	31	220	16	200	60
Vit. C	3			1				2		
Vit. K	0.29	0.19	-	-	0.25	-	-	-		
Thiamine	0.3	0.41	0.72	0.42	0.45	0.47	0.45	0.45	0.39	0.5
Ribo-flavin	0.51	0.37	0.15	0.2	0.49	0.21	0.51	0.09	0.41	0.48
Nicotinic-acid	2.1	2	2.4	1.5	1.5	3.5	2.6	1.5	2.2	1.3
Biotin (g/100g)	10	7.5	-	-	13.2	-	7.6	-	7.5	202
Choline	194	206	-	-	299	-	183	-	-	-
Folic-acid (g/100g)	125	144	-	-	107	-	83	-	100	-
Inositol	240	90	-	-	130	-	100	-	140	-
Pantothenic-acid	1.3	3.5	-	-	1.6	-	1.5	-	2.6	-
Total No. of	12	11	5	6	11	5	10	6	9	6
Vitamins/Minerals										

Source: Indian Council of Medical Research (ICMR), Hyderabad, 2012.

- 1.1.3 In India, pulses are generally produced in poor soils not suited to other crops, with a minimum use of resources and have a very low water footprint. They are vital constituent of cropping and consumption pattern. Of the total net sown area of 141.40 million hectares, 52 *per cent i.e.* 73.20 million hectares is rainfed. The pulses cultivation occupies major area under this ecology. The rainfed regions of the country supports 40% of human population and 2/3rd of live stock, further 90% of coarse cereals, 80% pulses, 74% oilseeds, 65% cotton and 48% rice is rainfed.
- 1.1.4 Pulses play a greater role in sustaining the economy of the rainfed farming community in a variety of ways. Besides improving soil fertility and physical structure, pulses fit well in mixed/intercropping systems, crop rotation and dry farming, provide green vegetable (pods/beans) and nutritious fodder for cattle as well thereby contributing to a more sustainable food system. Cultivation of pulses builds-up a mechanism to fix

atmospheric nitrogen to N-compounds in their root nodules and tend to fix 72 to 350 kg N per ha per year, thereby meeting their own nitrogen requirements to a great extent. The cultivation of the pulses under irrigation is only about 20% of their cropped area, remaining 80% are grown under *rainfed* conditions. Gram with 35% area under irrigation is the highest pulse crop followed by other pulses crop within < 10% irrigated area.

1.1.5 In addition to their nutritive value, by virtue of broad genetic diversity in food legumes and climate resilience to sustain well in adverse weather situations, the government of India has, undertaken various measures to tackle the issue of domestic supply constraints of pulses on one hand and addressing the issues of calamity prone rainfed areas farmers to double their income by 2022.

1.2 India's status of pulse production

- 1.2.1 The total world acreage under pulses is about 85.40 (Mha) with production of 87.40 (Mt) at 1023 kg/ha yields level. India, with >29 Mha pulses cultivation area, is the largest pulse producing country in the world. It ranks first in area and production with 35 *per cent* and 29 *per cent* respectively. During 2017-18 our productivity at 841kg/ha, has also increased significantly over Eleventh (662 kg/ha) and Twelfth plan (745 kg/ha).
- 1.2.2 Thanks to pro-active pulse programme implementation strategies and robust monitoring mechanism of Govt. of India, significant growth in area, production and productivity of pulses has been recorded in the XIIth Plan (2012-13 to 2016-17), *especially during the last 03 years of this Plan period, despite two consecutive drought years of 2014-15 to 2015-16*. More visible and significant increasing trends during 2016-17 and 2017-18, whereby the pulses production reached at 23.13 Mt and 25.24 Mt respectively, is a grand success story in itself. The productivity of pulses has increased 13 *per cent* at 841 kg/ha during 2017-18 from the level of 743 kg/ha during 2014-15. The production growth has been (47 *per cent*) the ever highest.

1.3 Pulses share to total foodgrain basket

- 1.3.1 In India foodgrains occupy 65% of total gross cropped area comprising cereals in 50% and pulses in about 15%. Within pulses, gram occupies 5% area followed by Urd 3%, Arhar 2% and Mung 2% the other pulses cover about 3% of gross cropped area.
- 1.3.2 Percent share of pulses to total foodgrain production basket remained stagnated between 6-7 *per cent* uptill 2015-16 after the Green Revolution period (1960-70). The area also remained stagnant between 22-24 Mha *i.e.* 19 *per cent* of total food grain area till this period.
- 1.3.3 Deceleration of percent production contribution of pulses to total food grains basket prompted the present dispensation in the Ministry of Agriculture & FW to vigorously pursue the NFSM-Pulses with synergistic approach on Research & Development, procurement, marketing, and import-export policies etc.

1.3.4 The multi-pronged strategy of the government to protect the interest of farmers and the consumers has resulted into enhanced *per cent* contribution of about 9 *per cent* pulses to total food grains during 2017-18 from 6-7 *per cent* till 2015-16 which is the ever highest after 1980-81. 1990-91 the production of food grains and contribution of pulses to total foodgrains basket is depicted *under (Table-1.2)*.

(Table-1.2): Contribution of pulses to food grains basket

{Area- Million ha, Production- Million Tones, Yield-kg/ha}

Year		Pulses		Fo	ood grains		Pulses share to	Pulses share to foodgrains (%)		
	A	P	Y	A	P	Y	A	P		
1950-51	19.09	8.41	441	97.32	50.82	522	19.62	16.55		
1960-61	23.56	12.70	539	115.58	82.02	710	20.38	15.48		
1970-71	22.54	11.82	524	124.32	108.42	872	18.13	10.90		
1980-81	22.46	10.63	473	126.67	129.59	1023	17.73	8.20		
1990-91	37.25	20.36	547	140.83	182.49	1300	26.45	11.16		
2000-01	20.35	11.08	544	121.05	196.81	1626	16.81	5.63		
2010-11	26.40	18.24	691	126.67	244.49	1930	20.84	7.46		
2011-12	24.46	17.09	699	124.76	259.32	2079	19.61	6.59		
2012-13	23.25	18.34	789	120.77	257.12	2129	19.25	7.13		
2013-14	25.21	19.25	764	125.04	265.04	2120	20.16	7.26		
2014-15	23.10	17.16	743	122.07	252.67	2069	18.92	6.79		
2015-16	24.91	16.35	656	123.22	251.57	2042	20.22	6.50		
2016-17	29.44	23.13	786	129.23	275.11	2129	22.78	8.40		
2017-18*	29.99	25.23	841	127.56	284.83	2233	23.51	8.85		




Fig-1.1: Contribution of Pulses to Foodgrains Basket

1.4 Season & crop contribution in total pulse production (2017-18)

Under individual crop category gram with 46 *per cent* production share to total pulses is the highest contributor followed by tur (17 *per cent*), urd (>14%) and mung (8%). The crop-wise APY and *per cent* share to total pulses is given below (*Table-1.3*).

(Table -1.3): Crop contribution to total pulse production

{Area-lakh ha, Production-lakh tons, Yield-kg/ha}

Crop		2017-18*		Contribution (%)		
	Area	Production	Yield	Area	Production	
Gram	105.61	112.29	1063	35.21	44.50	
Tur	44.31	42.54	960	14.77	16.86	
Urd	54.39	35.62	655	18.13	14.12	
Mung	42.57	20.09	472	14.19	7.96	
Other Kharif Pulses	18.71	8.15	436	6.24	3.23	
Other Rabi Pulses	34.34	33.65	980	11.45	13.33	
Total Kharif Pulses	140.83	93.45	664	46.95	37.03	
Total Rabi Pulses	159.10	158.90	999	53.05	62.97	
Total	299.93	252.35	841			

Source: DES, Ministry of Agri. & FW (DAC&FW), Govt. of India; $2017-18*-IV^{th}$ Adv. Est. OKP-other kharif pulses, ORP-other rabi pulses




Fig-1.2: Crop contribution in Total Pulses

1.5 States' Contribution – (2017-18)

1.5.1 Total Pulses

In India, total pulse area and production during 2017-18 has been >299 Lha and 252 Lt respectively. Out of the total area >74 Lha is confined to Madhya Pradesh alone, earning a prime status in pulse production commodity contributing a remarkable 25% of the countryøs pulse area with 32% production, thereby ranking first both in area and production followed by Rajasthan and Maharashtra with 13% each and Uttar Pradesh at 9%.

Compared to normal production, the estimated production during 2017-18 is 30% higher in case of total pulses, 32% gram, 27% arhar, 58% urdbean, 18% mungbean and 40% higher lentil production.

More than 90 *per cent* of total pulse production has been contributed by 10 states of MP, Rajasthan, MS, UP, Karnataka, AP, Gujarat, Jharkhand, Tamilnadu and Chhatisgarh.

(Table-1.4): States@Contribution in Area & Production ó Total Pulses

(Area-Lakh ha, Production-Lakh tons)

States	Area	% Contr.	States	Production	% Contr.
Madhya Pradesh	74.80	24.94	Madhya Pradesh	81.12	32.14
Rajasthan	53.29	17.77	Rajasthan	33.86	13.42
Maharashtra	43.52	14.51	Maharashtra	33.04	13.09
Karnataka	30.21	10.07	Uttar Pradesh	22.08	8.75
Uttar Pradesh	22.68	7.56	Karnataka	18.55	7.35
Andhra Pradesh	14.07	4.69	Andhra Pradesh	12.24	4.85
Gujarat	9.10	3.03	Gujarat	9.26	3.67
Tamilnadu	8.65	2.88	Jharkhand	8.45	3.35
Jharkhand	7.93	2.64	Tamilnadu	5.50	2.18
Chhattisgarh	7.84	2.61	Chhattisgarh	5.43	2.15
Others	27.84	9.28	Others	22.81	9.04
All India	299.93		All India	252.35	




Fig-1.3: States@Contribution in Area & ProductionóTotal Pulses

1.5.2 Kharif Pulses

The total area coverage and production of Kharif Pulses during 2017-18 has been 140 Lha and 93 Lt respectively. Madhya Pradesh outshined with first rank in production with >24 *per cent* production followed by Rajasthan and Maharashtra with 18% and 16%.

About 94 *per cent* of total kharif production was realized from 10 states of MP, Rajasthan, Maharashtra, Karnataka, Uttar Pradesh, Gujarat, Jharkhand, Telangana, Odisha and Andhra Pradesh. *(Table-1.5)*

(Table-1.5): States@Contribution in Area & Productionó Kharif Pulses

(Area-Lakh ha, Production-Lakh tons)

States	Area	% Contr.	States	Prod.	% Contr.
Rajasthan	37.09	26.34	Madhya Pradesh	22.82	24.42
Madhya Pradesh	26.80	19.03	Rajasthan	16.41	17.56
Maharashtra	22.47	15.96	Maharashtra	14.68	15.71
Karnataka	16.01	11.37	Karnataka	10.68	11.43
Uttar Pradesh	8.92	6.33	Uttar Pradesh	6.33	6.77
Gujarat	5.59	3.97	Gujarat	5.23	5.60
Telengana	4.56	3.24	Jharkhand	4.05	4.33
Odisha	4.41	3.13	Telengana	3.51	3.76
Jharkhand	4.11	2.92	Odisha	2.48	2.66
Andhra Pradesh	3.67	2.61	Andhra Pradesh	1.82	1.95
Others	7.19	5.11	Others	5.43	5.82
All India	140.83		All India	93.45	




Fig-1.4: States@Contribution in Area & Productionó Kharif Pulses

1.5.3 Rabi Pulses

All India Rabi pulse acreage and production has been recorded in 159 Lha and >158 Lt. Madhya Pradesh with 30 *per cent* of area and 37 *per cent* of total rabi pulse production in the country outshined at first rank.

More than 90 *per cent* pulse production was recorded from 10 states of MP, MS, Rajasthan, UP, AP, Karnataka, Chhattisgarh, Jharkhand, Gujarat and Bihar (*Table-1.6*).

(Table-1.6): StatesøContribution in Area & Production- Rabi Pulses

(Area-Lakh ha, Production-Lakh tons)

States	Area	% Contr.	States	Production	% Contr.	
Madhya Pradesh	48.00	30.17	Madhya Pradesh	58.30	36.69	
Maharashtra	21.04	13.23	Maharashtra	18.36	11.55	
Rajasthan	16.20	10.18	Rajasthan	17.45	10.98	
Karnataka	14.20	8.93	Uttar Pradesh	15.75	9.91	
Uttar Pradesh	13.76	8.65	Andhra Pradesh	10.42	6.56	
Andhra Pradesh	10.40	6.54	Karnataka	7.87	4.95	
Tamilnadu	6.40	4.03	Chhattisgarh	4.59	2.89	
Chhattisgarh	5.84	3.67	Jharkhand	4.41	2.77	
Bihar	4.39	2.76	Gujarat	4.03	2.54	
Jharkhand	3.82	2.40	Bihar	3.74	2.36	
Others	15.04	9.46	Others	13.99	8.80	
All India	159.10		All India	158.90		




Fig-1.5: States@Contribution in Area & Productionó Rabi Pulses

1.5.4 Gram (Chickpea)

In 2017-18 this crop was cultivated in about 106 Lha. The country harvested a record production of > 112 Lt at a highest productivity level of 1063 kg/ha. As usual, MP has contributed a significant 34% of the total gram area and 41% of total gram production in the country, thereby ranking first both in area and production followed by Maharashtra in area (19% and 16%).

About 98 *per cent* of gram production of the country during the period under report has been realized by 10 states of MP, MS, Rajasthan, Karnataka, AP, UP, Gujarat, CG, Jharkhand, and Telangana (*Table-1.7*).

(Table-1.7): StatesøContribution in Area & Production- Gram

(Area-Lakh ha, Production-Lakh tons)

States	Area	% Contr.	States	Production	% Contr.
Madhya Pradesh	35.90	33.99	Madhya Pradesh	45.95	40.92
Maharashtra	20.00	18.94	Maharashtra	17.84	15.89
Rajasthan	15.72	14.89	Rajasthan	16.70	14.87
Karnataka	12.65	11.98	Karnataka	7.21	6.42
Andhra Pradesh	5.20	4.92	Andhra Pradesh	5.85	5.21
Uttar Pradesh	5.01	4.74	Uttar Pradesh	5.79	5.16
Chhattisgarh	3.18	3.01	Gujarat	3.67	3.27
Gujarat	2.93	2.77	Chhattisgarh	3.16	2.82
Jharkhand	2.33	2.20	Jharkhand	2.93	2.61
Telengana	0.97	0.92	Telengana	1.38	1.23
Others	1.72	1.63	Others	1.81	1.61
All India	105.61		All India	112.29	




Fig-1.6: StatesøContribution in Area & ProductionóGram

1.5.5 Arhar (Pigeonpea)

The countryøs total area coverage and production of tur has been about 44 Lha and 42 Lt respectively. As usual Maharashtra has contributed >28 per cent of area and 25 per cent of total production during this period. With aggressive ToT in various thematic areas, highest ever productivity levels of 960 kg/ha were achieved during 2017-18 (Table-1.8).

About than 97 *per cent* of Arhar production of the country during the period under report has been realized by 10 states of MS, MP, Karnataka, Gujarat, UP, Telangana, Jharkhand, Odisha, AP and TN *(Table-1.8)*.

(Table-1.8): StatesøContribution in Area & Production ó Pigeonpea

(Area-Lakh ha, Production-Lakh tons)

States	Area	% Contr.	States	Production	% Contr.
Maharashtra	12.29	27.73	Maharashtra	10.73	25.22
Karnataka	8.85	19.97	Madhya Pradesh	8.39	19.72
Madhya Pradesh	6.47	14.60	Karnataka	7.68	18.06
Telengana	3.30	7.45	Gujarat	3.37	7.92
Uttar Pradesh	2.82	6.36	Uttar Pradesh	3.32	7.80
Andhra Pradesh	2.79	6.30	Telengana	2.63	6.18
Gujarat	2.71	6.12	Jharkhand	2.22	5.22
Jharkhand	1.94	4.37	Odisha	1.24	2.91
Odisha	1.38	3.11	Andhra Pradesh	1.19	2.80
Chhattisgarh	0.58	1.30	Tamilnadu	0.61	1.44
Others	1.19	2.68	Others	1.17	2.74
All India	44.31		All India	42.54	




Fig.-1.7: StatesøContribution in Area & Production ó Arhar/Tur

1.5.6 Mungbean (Greengram)

During 2017-18, the total coverage under mungbean has been about 42 Lha with a production of 20 Lt. There has been phenomenal increase in area of mungbean in the country from 2015-16 onwards. Rajasthan with >40 per cent area and 37 per cent of production outshined in the total mungbean contribution in the country during year report.

More than 90 *per cent* of mungbean production comes from 10 states of Rajasthan, Madhya Pradesh, Maharashtra, Karnataka, Bihar, Andhra Pradesh, Odisha, TN, Gujarat, and Telangana (*Table-1.9*).

(Table-1.9): States@Contribution in Area & Production \u00f3 Mungbean

(Area-Lakh ha, Production-Lakh tons)

States	Area	% Contr.	States	Production	% Contr.
Rajasthan	17.21	40.43	Rajasthan	7.47	37.18
Maharashtra	4.53	10.64	Madhya Pradesh	2.66	13.23
Madhya Pradesh	4.34	10.19	Maharashtra	1.64	8.17
Karnataka	3.97	9.33	Karnataka	1.28	6.39
Odisha	2.36	5.55	Bihar	1.04	5.19
Tamilnadu	1.85	4.33	Andhra Pradesh	0.99	4.93
Bihar	1.68	3.95	Odisha	0.87	4.31
Gujarat	1.53	3.59	Tamilnadu	0.86	4.29
Andhra Pradesh	1.40	3.29	Gujarat	0.86	4.28
Telengana	0.99	2.33	Telengana	0.65	3.24
Others	2.71	6.37	Others	1.77	8.81
All India	42.57		All India	20.09	




Fig.-1.8: State & Contribution in Area & Production-Mungbean

1.5.7 Urdbean (Blackgram)

Urdbean crop is also gaining momentum since 2015-16 and there has been phenomenal increase in its coverage. During 2017-18 the crop was cultivated in an area of > 54 Lha. The success of this crop was released with a harvest of about 36 Lt at an ever highest yield levels of 655 kg/ha.

About 95 *per cent* of urdbean production comes from 10 states of Madhya Pradesh, Rajasthan, Andhra Pradesh, Uttar Pradesh, Tamil Nadu, Maharashtra, Jharkhand, Gujarat, Karnataka and West Bengal. *(Table-1.10)*

(Table-1.10): States@Contribution in Area & Production- Urdbean

(Area-Lakh ha, Production-Lakh tons)

States	Area	% Contr.	States	Production	% Contr.
Madhya Pradesh	18.24	33.54	Madhya Pradesh	13.48	37.84
Rajasthan	8.39	15.43	Rajasthan	5.24	14.70
Uttar Pradesh	6.14	11.29	Andhra Pradesh	3.75	10.53
Maharashtra	4.84	8.90	Uttar Pradesh	3.15	8.84
Tamilnadu	4.54	8.34	Tamilnadu	2.85	8.01
Andhra Pradesh	4.03	7.41	Maharashtra	1.77	4.98
Jharkhand	1.48	2.72	Jharkhand	1.32	3.70
Gujarat	1.37	2.52	Gujarat	0.98	2.75
Karnataka	1.36	2.50	Karnataka	0.65	1.84
Chhattisgarh	0.98	1.80	West Bengal	0.63	1.77
Others	3.03	5.56	Others	1.80	5.05
All India	54.39		All India	35.62	




Fig.-1.9: States@Contribution in Area & Production- Urdbean

1.5.8 Masoor (Lentil)

The total area covered under lentil has been 16 Lha during 2017-18. The highest ever production of 16 Lt at 1034 kg/ha is a remarkable success.

About 98% has been realized from 07 states of Madhya Pradesh, Uttar Pradesh, West Bengal, Bihar, Jharkhand, Rajasthan and Assam (*Table-1.11*).

(Table-1.11): StatesøContribution in Area & Production-Lentil

(Area-Lakh ha, Production-Lakh tons)

States	Area	% Contr.	States	Production	% Contr.
Madhya Pradesh	5.96	38.35	Madhya Pradesh	6.79	42.25
Uttar Pradesh	4.84	31.15	Uttar Pradesh	4.98	30.99
West Bengal	1.58	10.17	West Bengal	1.55	9.64
Bihar	1.50	9.65	Bihar	1.39	8.64
Jharkhand	0.69	4.47	Jharkhand	0.60	3.75
Rajasthan	0.31	2.02	Rajasthan	0.34	2.11
Assam	0.27	1.74	Assam	0.20	1.24
Chhattisgarh	0.13	0.84	Uttarakhand	0.07	0.44
Odisha	0.11	0.68	Odisha	0.06	0.35
Uttarakhand	0.10	0.64	Chhattisgarh	0.05	0.29
Others	0.05	0.32	Others	0.05	0.28
All India	15.54		All India	16.07	





Fig.-1.10: States@Contribution in Area & Production-Lentil

PULSES OVERVIEW





Chapter- 2

Plan-wise Pulse Overview

2.1 Background

- Demographic dividend refers to the rise in the rate of economic growth due to rising share of working age people in a population. India's demographic dividend- *i.e.* its working-age (15-59 years) population, largely consisting youth between 15-34 years age, provide an inherent edge and potential to its economy to grow much faster than that of many other countries, including neighbouring China.
- India is the second most populous country in the world with >1/6th of the worldøs population. The stock of population changed from 102 crore in 2001 to 121 crore in 2011, the exponential population growth rate being 1.64 during 2001 to 2011. Although, India occupies only 2.2% of the worldøs land area, it supports approx 18% of the worldøs population.
- The census projection report has further revealed that the proportion of the working age population between 15 and 59 years is likely to increase from 58% in 2001 to > 64 per cent by 2021. Such a trend would make the country one of the youngest nations in the world. Thus, one of the Indias@competetive advantages is its demographic dividend.
- The demographic dividend has been regarded as a key factor for economic growth. The existing demographic dividend provides a great opportunity, however, also poses a great challenge. There are many challenges which the country needs to converge these into opportunities. The growth in the working-age ratio is likely to be concentrated in the rainfed and poorest states, mainly cultivating pulses, oilseeds and rearing livestock. Here, the youth have to be strengthened/targeted towards proper nourishment, skill development inagriculture and allied sector, pulses based sustainable cropping system to overcome the issues on climate change and global warming to fully realize the dividend.
- The future agricultural operations are likely to be highly skilled and competitive. The serious challenges to the workforce/youth of these resource poor and rainfed regions viz. lack of skill in scientific crop cultivation, repair and maintenance of farm machineries and implements, production of quality seeds, primary processing, value addition, modern animal husbandry, poor infrastructure (irrigation, go downs/ware houses ,trading centres) and organized pulse markets etc. have been considered by the government while formulating the strategy and roadmap to increase the production of pulses.
- According to the Human Development Report (HDR) published by the United Nations
 Development Programme (UNDP), India is still in the medium human development
 category with countries like China, Sri Lanka, Thailand, Philippines, Egypt, Indonesia,
 South Africa, and even Vietnam has a better rank. Therefore health and education
 parameters need to be improved substantially to make the Indian workforce efficient and

skilled. At the primary level, there are serious problems of health and nutrition that impact the effectiveness of education, learning capacity etc.

• The poor nutritional status of the population is a major challenge where low income, small scales households, has a long term negative implication for economic development. The government has resolved this issue to be addressed through **nutrition-sensitive agriculture interventions**, focusing the pulse crops having multiple nutritional values with essential source of vitamins, micro-nutrient and protein to help attain **nutritional security**. Pulses have better enabling environment to promote dietary and production diversity to address hunger and malnutrition at national level.

Focus on pulses production and consumption can help overcome malnutrition: India should include pulses in the public distribution system

• India, a country with high concentrations of poor and malnourished people, long promoted a cereal-centric diet composed of subsidized staple commodities such as rice and wheat to feed its population of more than a billion. Today, however, dietary patterns are changing. Policy makers, researchers, and health activists are looking for ways to fight hunger and malnutrition in the country. As they shift their focus from calorie intake to nutrition, neglected foods such as pulses (the dried, edible seeds of legumes) are gaining attention. There are three kinds of hunger that needed to be dealt with 6 calorie inadequacy, protein deficiency and micronutrient deficiency.

The Global Nutrition Report 2017

- India is facing a serious burden of under-nutrition, according to a global report which shows that more than half the women of reproductive age in the country suffer from anemia. The significant burdens of three important forms of malnutrition, used as an indicator of broader trends are i.) childhood stunting-38 *per cent* of children under five are affected by stunting-children too short for their age due to lack of nutrients, suffering irreversible damage to brain capacity ii.) anemia in women of reproductive age and iii.) overweight adult women.
- About 21 per cent of children under 5 are defined as øwastedø or ∹severely wastedø meaning they do not weigh enough for their height.
- Over half of women of reproductive age -51 *per cent* suffer from anemia a serious condition that can have long-term health impacts for mother and child.
- More than 22 *per cent* of adult women are overweight, a rising concern as women are disproportionately affected by the global obesity epidemic.

- While the country has shown some progress in addressing under-5 stunting, it has made
 no progress or presents worse outcomes in the percentage of reproductive-age women
 with anemia, and is of course in terms of reaching targets for reducing adult obesity and
 diabetes, the report said.
- The Global Nutrition Report highlights that the double burden of under-nutrition and obesity needs to be tackled as part of Indiags national nutrition strategy.
- The Global Nutrition Report 2017 calls for nutrition to be placed at the heart of efforts to end poverty, fight disease, raise educational standards and tackle climate change. A well-nourished child is one third more likely to escape poverty. They will learn better in school, be healthier and grow into productive contributors to their economies. Good nutrition provides the brainpower, the ;grey matter infrastructureøto build the economies of the future.ö
- With one of the highest rates of child malnutrition in the world, India has won notoriety
 as one of the nutritional basket cases of the world over the past few years. Although India
 has witnessed significant progress in its battle against child malnutrition over the past
 decade, the progress has been quite uneven, and child malnutrition rates still remain high
 in many parts of the country.
- As in the case of adult under nutrition rates, districts with the highest levels of under nutrition seem to be clustered largely in the central parts of the country. The bottom quartile of districts ranked according to child malnutrition rates includes not just districts from the most deprived tribal belts of central and eastern India but also some of the more urbanized districts of the country such as Udaipur in Rajasthan, Aurangabad in Maharashtra, Lucknow in Uttar Pradesh, Patna in Bihar, and Ranchi in Jharkhand. However, overall urban child malnutrition rates are lower than that of rural India.
- The recent Government has undertaken various measures to impart skill to the Indian workforce to reap the benefits of demographic dividends and to make them employable and help secure a decent job to enhance their income. The National Skill Development Corporation India (NSDC) to contribute significantly (about 30 *per cent*) to the overall target of skilling / up-skilling 500 million people in India by 2022, mainly by fostering private sector initiatives in skill development programmes and providing funding.
- Pulses are grown in all three seasons. The three crop seasons for the commodity are:
 - i. *Kharif*: Arhar (Tur), Urd (Blackgram), Moong (Greengram), Lobia (Cowpea), Kulthi (Horsegram) and Moth;
 - ii. *Rabi*: Gram, Lentil, Pea, Lathyrus and Rajmash;
 - iii. Summer: Greengram, Blackgram and Cowpea.

2.2 Pulses share to total foodgrain basket

- In India foodgrains occupy 65% of total gross cropped area comprising cereals in 50% and pulses in about 15%. Within pulses, gram occupies 5% area followed by Urd 3%, Arhar 2% and Mung 2% the other pulses cover about 3% of gross cropped area.
- Percent share of pulses to total food-grain production basket remained stagnated between 6-7 *per cent* uptill 2015-16 after the Green Revolution period (1960-70). The area also remained stagnant between 22-24 Mha *i.e.* 19 *per cent* of total food grain area till this period.
- Deceleration of percent production contribution of pulses to total food grains basket prompted the present dispensation in the Ministry of Agriculture & FW to vigorously pursue the NFSM-Pulses with synergistic approach on Research & Development, procurement, marketing, and import-export policies etc.
- The multi-pronged strategy of the government to protect the interest of farmers and the consumers has resulted into enhanced percent contribution of about 9 *per cent* pulses to total food grains during 2017-18 from 6-7 *per cent* till 2015-16 which is the ever highest after 1980-81, 1990-91 the production of food grains and contribution of pulses to total foodgrains basket is depicted *under (Table-2.1)*.

(Table-2.1): Contribution of pulses to total foodgrains in India.

{Area- Million ha, Production- Million Tonnes, Yield-kg/ha}

Year		Pulses			Foodgrains				oodgrains
10	A	P	Y	A	P	Y	A	P	YI
1950-51	19.09	8.41	441	97.32	50.82	522	19.62	16.55	84.48
1960-61	23.56	12.70	539	115.58	82.02	710	20.38	15.48	75.92
1970-71	22.54	11.82	524	124.32	108.42	872	18.13	10.90	60.09
1980-81	22.46	10.63	473	126.67	129.59	1023	17.73	8.20	46.24
1990-91	24.66	14.26	578	127.84	176.39	1380	19.29	8.08	41.88
1995-96	22.28	12.31	552	121.01	180.42	1491	18.41	6.82	37.02
2000-01	20.35	11.08	544	121.05	196.81	1626	16.81	5.63	33.46
2001-02	22.01	13.37	607	122.78	212.85	1734	17.93	6.28	35.01
2002-03	20.50	11.13	543	113.86	174.77	1535	18.00	6.37	35.37
2003-04	23.46	14.91	635	123.45	213.19	1727	19.00	6.99	36.77
2004-05	22.76	13.13	577	120.00	198.36	1652	18.97	6.62	34.93
2005-06	23.39	13.39	598	121.60	208.60	1715	18.41	6.42	34.87
2006-07	23.76	14.11	594	124.07	211.78	1707	19.15	6.66	34.80
2007-08	23.63	14.76	625	124.07	230.78	1860	19.05	6.40	33.58
2008-09	22.09	14.57	660	122.83	234.47	1909	17.98	6.21	34.55
2009-10	23.28	14.66	630	121.33	218.11	1798	19.19	6.72	35.03
2010-11	26.40	18.24	691	126.67	244.49	1930	20.84	7.46	35.80
2011-12	24.46	17.09	699	124.76	259.32	2079	19.61	6.59	33.61
2012-13	23.25	18.34	789	120.77	257.12	2129	19.25	7.13	37.06
2013-14	25.21	19.25	764	125.04	265.04	2120	20.16	7.26	36.03
2014-15	23.10	17.16	743	122.07	252.67	2069	18.92	6.79	35.91
2015-16	24.91	16.35	656	123.22	251.57	2042	20.22	6.50	32.13
2016-17	29.45	23.13	785	129.23	275.11	2129	22.78	8.40	36.88
2017-18*	29.99	25.24	841	127.56	284.83	2233	23.51	8.86	37.66




Fig-2.1: Contribution of Pulses to Foodgrains Basket

(Table - 2.2): Season-wise Pulse Contribution to Total Pulses

{Area-lakh ha, Production-Lakh Tones, Yield-kg/ha}

Year	Kharif Pulses			ont. to T	Total	Ra				tal Pulses					
					Pulses						Pulses				
	A	P	Y	A	P	ΥI	A	P	Y	A	P	YI	A	P	Y
2000-01	106.58	44.48	417	52.4	40.2	77	96.90	66.27	684	47.6	59.8	126	203.48	110.75	544
2001-02	107.22	48.38	451	48.7	36.2	74	112.86	85.30	756	51.3	63.8	124	220.08	133.68	607
2002-03	99.50	41.51	417	48.5	37.3	77	105.46	69.74	661	51.5	62.7	122	204.96	111.25	543
2003-04	116.83	61.65	528	49.8	41.4	83	117.75	87.41	742	50.2	58.6	117	234.58	149.05	635
2004-05	113.17	47.17	417	49.7	35.9	72	114.46	84.12	735	50.3	64.1	127	227.63	131.30	577
2005-06	106.80	48.65	456	47.7	36.3	76	117.12	85.20	727	52.3	63.7	122	223.91	133.84	598
2006-07	106.76	47.95	449	46.0	33.8	73	125.16	94.02	751	54.0	66.2	123	231.92	141.98	612
2007-08	114.90	64.03	557	48.6	43.4	89	121.44	83.58	688	51.4	56.6	110	236.33	147.62	625
2008-09	98.09	46.86	478	44.4	32.2	72	122.85	98.80	804	55.6	67.8	122	220.94	145.66	659
2009-10	105.82	42.04	397	45.5	28.7	63	127.00	104.58	823	54.5	71.3	131	232.82	146.62	630
2010-11	123.20	71.20	578	46.7	39.0	84	140.82	111.21	790	53.3	61.0	114	264.02	182.41	691
2011-12	111.90	60.58	541	45.7	35.4	77	132.72	110.31	831	54.3	64.6	119	244.62	170.89	699
2012-13	99.54	59.16	594	42.8	32.3	75	133.03	124.27	934	57.2	67.7	118	232.57	183.42	789
2013-14	103.33	59.95	580	41.0	31.1	76	148.85	132.60	891	59.0	68.9	117	252.18	192.55	764
2014-15	99.98	57.31	573	42.4	33.4	79	135.55	114.22	843	57.6	66.6	116	235.53	171.52	728
2015-16	113.14	55.30	489	45.4	33.8	74	135.98	108.18	796	54.6	66.2	121	249.11	163.48	656
2016-17	143.63	95.85	667	48.7	41.4	84	150.83	135.47	898	51.2	58.5	114	294.47	231.31	785
2017-18*	140.83	93.45	664	47.0	37.0	79	159.10	158.90	999	53.0	63.0	119	299.93	252.35	841

2.3 Per capita availability of pulses in India

As a result of stagnant pulse production and continuous increase in population, the per capita availability of pulses has decreased considerably. The *per capita* per day availability of pulses in 1951 was 60 g that dwindled down to a provisional level of 47.2 g in the year 2014. The *per capita* per year availability shows the same decreasing trend from 22.1 kg in 1951 to 17.2 kg in 2014.

Per capita availability enhanced during 2017-18. In conformity to FSA- 2013 to ensure nutritional security to vegetarian population, the per capita per day availability of pulses which dwindled down to a provisional level of 41-42 g (15-16 kg/annum) between 1991 to 2013, is now attend at the level of 53 g per head/day i.e > 19 kg/annum/person (*Table-2.3*).

(Table- 2.3): Per capita availability of pulses in India

Year	Pulses .	Availability
	(g <i>per capita per</i> day)	(kg <i>per capita</i> per year)
1951	60.7	22.1
1961	69.0	25.2
1971	51.2	18.7
1981	37.5	13.7
1991	41.6	15.2
1992	34.3	12.5
1993	36.2	13.2
1994	37.2	13.6
1995	37.8	13.8
1996	32.7	12.0
1997	37.1	13.5
1998	32.8	12.0
1999	36.5	13.3
2000	31.8	11.6
2001	30.0	10.9
2002	35.4	12.9
2003	29.1	10.6
2004	35.8	13.1
2005	31.5	11.5
2006	32.5	11.8
2007	35.5	12.9
2008	41.8	15.3
2009	37.0	13.5
2010	35.4	12.9
2011	43.0	15.7
2012	41.6	15.2
2013	43.3	15.8
2014	46.4	16.9
2015	43.8	16.0
2016	43.0	15.7
2017 (P*)	52.9	19.3

P - Provisional figures are based on IIIrd Advance Estimates of production for 2017-18, Source: Press Information Bureau, Ministry of Agriculture & Farmers Welfare.*




Fig.- 2.2: Pulses Availability

2.4 Projected Demand: (XIth & XIIth Plan)

The XIIth Plan Working Group of Planning Commission (NITI Aayog) had worked-out a demand of pulses from 2012-13 to 2017-18 presented in *Table-2.4*. The working group projected a production gap of 1.79 to 5.70 Mt up till 2015-16.

As a result of phenomenal increase in production of pulses during 2016-17 & 2017-18, consequent upon the implementation of short term, mid-term and long term strategy to promoter the pulse sector, the government meet the pulse demand and moved closer to achieve its target to attend nutritional security by attending self-sufficiency production.

(Table- 2.4): Demand, Production, Growth and Projected Target

(Qty: Million Tonnes)

Year	Demand *	Production	Growth (%)	Gap	Target
2007-08	17.29	13.61		-3.68	17.00
2008-09	17.82	13.65	0.29	-4.17	18.00
2009-10	18.37	13.68	0.22	-4.69	18.50
2010-11	18.94	13.72	0.29	-5.22	19.00
2011-12	19.53	13.75	0.22	-5.78	20.00
2012-13	20.13	18.34	33.38	-1.79	18.24
2013-14	20.75	19.25	4.96	-1.5	19.00
2014-15	21.39	17.15	-10.91	-4.24	19.50
2015-16	22.05	16.35	-4.66	-5.70	20.05
2016-17	22.74	23.13	41.47	0.39	20.75
2017-18*	23.44	25.24	9.12	1.80	22.90

Note: *Demand includes seed, feed and wastage and based on behaviouristic approach.

2017-18*- IVth Adv. Est.

Source: XIIth Plan Working Group (Planning Commission) @3.09% growth rate.




Fig.-2.3: Demand vs Production

2.5 Import/Export and Availability

Import: The import of pulses in India during April, 2014 to March, 2015 was 45.85 lakh tonnes worth Rs.17063 crores against the value of Rs.17196.87 crore for total foodgrains, Rs.121319 crore for total agricultural imports and against Rs.2737087crore for total National Import. The import during April, 2015 to March, 2016 was 57.98 lakh tonnes worth Rs.21176 crore against the import value of Rs. 26841.87 crore for total foodgrains, Rs.140289 before for total agricultural import and Rs.2490298 crore for total National import respectively during this period. The share of Agricultural import to National import was 4.43% and 5.63% respectively during April, 2014 to March, 2015 and April, 2015 to March, 2016.

{Dry Peas contributes the single largest share in India's import basket of pulses registering in the total pulses import}.

Export:The pulses export of the country during April, 2014 to March, 2015 was 2.22 lakh 31 ones worth Rs.1218 crore against the value of Rs. 59500.54 crore for total foodgrains, Rs.239681crore for total agricultural exports and against Rs.1896445 crore for total National export. The export during April, 2015 to March, 2016 was 2.56 lakh tonnes worth Rs.1553 crore against the export value of Rs. 42622.29 crore for total foodgrains, Rs.215396 crore for total agricultural export and Rs.1716378crore for total National export respectively during this period. The share of agricultural export to National export was 12.64% and 12.55% respectively during April, 2014 to March, 2015 and April, 2015 to March, 2016.

{Chickpeas contributes the single largest share in India's export basket of pulses registering 85.64% and 84.87% share in the total pulses export during 2014-15 and 2015-16 respectively}.

(Table- 2.5-a): India imports and exports of pulses

(Quantity - Lakh tonnes, Values -Crore)

Year	Import		Expor	ts
	Import Quantity	Import Value	Export Quantity	Export Value
2007-08	28.35	5375	1.64	526
2008-09	24.74	6246	1.36	540
2009-10	35.10	9813	1.00	408
2010-11	27.78	7512	2.09	870
2011-12	34.96	9448	1.74	1068
2012-13	40.13	13345	2.03	1285
2013-14	31.78	11037	3.46	1749
2014-15	45.85	17063	2.22	1218
2015-16	57.98	21176	2.56	1553
2016-17	112.82	22160	1.54	1228
2017-18	82.96	14462	1.35	1120

Source: DGCI&S, Ministry of Commerce, Kolkata;

(Table- 2.5-b): India@s imports and exports of pulses v/s agriculture

(Rs.in Crore)

Year			Import			Exports				
	Total I	Pulses	Total Agriculture		Total National	Total 1	Pulses	Tot:		Total National
					Imports			Agriculture		Exports
	Actual	% to	Actual	% to		Actual	% to	Actual	% to	
	Value	Agri.	Value	Natio nal		Value	Agri.	Value	Natio nal	
2007-08	5375	23.84	22550	2.23	1012312	526	0.70	74673	11.39	655864
2008-09	6246	21.75	28719	2.09	1374436	540	0.67	81065	9.64	840755
2009-10	9813	18.05	54365	3.99	1363736	408	0.48	84444	9.99	845534
2010-11	7512	14.71	51074	3.03	1683467	870	0.77	113047	9.94	1136964
2011-12	9448	13.47	70165	2.99	2345463	1068	0.58	182801	12.47	1465959
2012-13	13345	13.94	95719	3.59	2669162	1285	0.57	227193	13.90	1634318
2013-14	11037	12.87	85727	3.16	2715434	1749	0.67	262779	13.79	1905011
2014-15	17063	14.06	121319	4.43	2737087	1218	0.51	239681	12.64	1896445
2015-16	21176	18.26	140289	5.63	2490298	1553	0.77	215396	12.55	1716378
2016-17	22160				2577666	1228				1849429
2017-18	14462				3001016	1120				1955541

Source: DGCI&S, Ministry of Commerce, Kolkata.

2.6 Exim policies in favour of pulses have paid

2.6.1 Import

The Cabinet Committee on Economic Affairs (CCEA) as a major step empowered the Committee headed by Food Secretary to review the export and import policy on pulses and consider measures such as quantitative restrictions, prior registration and changes in import duties of pulses depending on domestic production and demand, local and international prices and global trade volumes.

Farmer-friendly policy measures have helped reduce import of pulses. Import of pulses declined by 30 lakh tonnes from previous year, resulting in saving of foreign exchange amounting to Rs 7,698 crore.

The government ensured the availability supply as per demand by way of enhanced imports between 2014-15 to 2016-17 at about 5-6 million tons per year in their buffer stock on one hand and swung in to action to combat the natural calamities through development programmes risk management through PMFBY, PSS and PSF procurement on the other.

Because of back to back record production, Import duty on chickpeas has been fixed at 60%, while that for yellow peas is 50%, 30% for lentils and 10% for tur. Peas which accounted for major share in Indiags pulses import declined.

The government has also imposed a quantitative cap of 2 lakh tonnes per year on tur dal (Mozambique) and 3 lakh tonnes on urad and moong (Mynmar). Recently, the government had imposed quantitative restrictions on some of the pulses to check cheaper imports.

2.6.2 Export of Pulses

The duties on import were imposed and export was also encouraged to support the farmers. "The Cabinet Committee on Economic Affairs (CCEA) has given its approval for removal of prohibition on export of all types of pulses to ensure that farmers have greater choice in marketing their produce and in getting better remuneration for their produce.

The government lifted ban on export of tur, urad and moong dal, although shipments of these varieties were allowed only through permission from agriculture export promotion body APEDA. All varieties of pulses, including organic pulses, have been made \pm freeø for export and kabulichana has also been permitted in a limited quantity. Gram which accounted for major share in Indiaøs pulses export increased.

Opening of exports of all types of pulses will help the farmers dispose of their products at remunerative prices and encouragethem to expand the area of sowing.

2.6.3 Buffer Stock

As a major policy to support the consumers as well, the government has taken a number of steps to sustain high pulses production and procured 20 lakh tonnes of pulses directly from the farmers by ensuring minimum support price or market rates, whichever is higher.

India Pulses and Grains Association (IPGA) has appreciated the buffer stocking policy as a measure to correct price distortions, offer support to pulses selling below MSP (minimum support price) and revitalize the milling industry.




Fig.- 2.4: Import & Export of Pulses

(Table-2.6): Pulse importing and exporting countries of major pulses (2016-17)

Pulses	Top 5 Export Destinations	Top 5 Import Sources
Peas	Shri Lanka DSR (96.3%), Myanmar	Canada (54.5%), Russia (10.3%),
(PisumSativum)	(1.6%), Bhutan (1.4%), Nepal	Luthuania (9.0%), France (6.8%),
	(0.5%), U Arab EMTS (0.09%).	USA (6.4%)
Chickpeas	Pakistan (21.6%), U Arab EMTS	Australia (85.1%), Russia (4.7%),
(Garbanzos)	(10.6%), Algeria (11.6%), Saudi	Tanzania (3.8%), USA (1.4%),
	Arab (9.5%), Sri Lanka (7.3%)	Canada (0.91%),
Moong/Urad	USA (39.96%), Sri Lanka (13.05%),	Myanmar (70.37%), Kenya
	UK (9.86%), Australia (7.77%),	(7.43%), Australia (6.32%),
	Malaysia (7.63%)	Tanzania (3.15%), Uzbekistan
		(2.60%).
Lentils (Masur)	Sri Lanka DSR (43.39%),	Canada (89.58%), USA (7.47%),
	Bangladesh (18.11%), U Arab	Australia (2.88%), Turkey (0.03%),
	EMTS (8.35%), Egypt ARP	Mozambique (0.03%).
	(3.98%), USA (3.67%)	
Pigeon Peas(Tur)	USA (40.79%), U Arab EMTS	Myanmar (46.35%), Tanzania
	(18.28%), Canada (11.28%), UK	(18.71%), Mozambique (15.36%),
	(10.75%), Singapore (5.11%),	Malawi (12.56%), Sudan (3.36%)

(%) figures in parenthesis indicates percentage share of global import/export

(Table -2.7): Total pulses: Crop/Season-wise contribution

{Area-lakh ha, Production-lakh tonnes, Yield-kg/ha}

Crop	Season	Area	% Contr. to	Production Production	% Contr. to	Yield
			Tot. Pulses		Tot. Pulses	
Arhar	Kharif	41.90	16.58	32.88	17.45	785
Urd	Kharif	27.00	10.68	14.72	7.81	545
	Rabi/Summer	8.14	3.22	6.10	3.24	749
	Total	35.13		20.82		593
Moong	Kharif	24.93	9.86	10.51	5.58	421
	Rabi/Summer	9.62	3.81	5.60	2.97	582
	Total	34.55		16.11		466
Horse	Kharif	2.27	0.90	1.06	0.56	466
gram	Rabi/Summer	2.09	0.83	0.98	0.52	468
	Total	4.36		2.04		467
Moth	Kharif	10.53	4.16	3.32	1.76	315
Chickpea	Rabi	89.45	35.39	84.25	44.72	942
Lentil	Rabi	13.94	5.52	10.77	5.72	773
Peas &	Rabi	9.33	3.69	8.81	4.68	945
Beans						
Lathyrus	Rabi	4.58	1.81	3.84	2.04	837
Other	Kharif	5.30	2.10	3.03	1.61	572
Pulses	Rabi/Summer	3.69	1.46	2.54	1.35	689
	Total	8.99		5.58		620
Kharif Pul	ses	111.92	44.28	65.51	34.77	585
Rabi/Sumi	mer Pulses	140.84	55.72	122.90	65.23	873
Total Pulse	es	252.77		188.41		745

Source: DES, DAC&FW, GoI, ND (Normal-Avg. 2012-13 to 2016-17). (figures in parenthesis indicates % share of crop).

2.7 Growth Rate

From 1950-51 to 2006-07, the total acreage under pulses has almost been stagnated but for 1990-91 (24.66 million ha), however, the maximum growth rate in area was recorded between the period from 2002-03 to 2003-04 at 14%. Maximum production growth rate of 34% and maximum yield growth rate of 17% were also observed during the same period. The highest production (23 million tonnes) & yield (785 kg/ha) was recorded during 2016-17 followed by 19 & 18 million tones during 2013-14 & 2012-13 (Table-2.8).

(Table-2.8): Growth rate of total pulses

(Area-Million ha, P-Million tones, Y-kg/ha, Growth Rate (GR)-%)

Year	Area	GR	Prod.	GR	Yield	GR	% coverage
		- GIL		011		511	under irrigation
1950-51	19.09		8.41		441		9.4
1955-56	23.22	4.3	11.04	6.3	476	1.6	8.4
1960-61-	23.56	0.3 -	12.7	3.0	539	2.6 -	8
1965-66	22.72	-0.7	9.94	-4.3	438	-3.7	9.4
1967-68*	22.65	-0.1	12.1	4.3	534	4.4	8.7
1970-71	22.54	-0.1	11.82	-0.5	524	-0.4	8.8
1975-76	24.45	1.7	13.04	2.1	533	0.3	7.9
1980-81	22.46	-1.6	10.63	-3.7	473	-2.3	9.0
1985-86	24.42	1.7	13.36	5.1	547	3.1	8.5
1990-91	24.66	0.2	14.26	1.3	578	1.1	10.5
1995-96	22.28	-1.9	12.31	-2.7	552	-0.9	12.9
1996-97	22.45	0.2	14.24	3.1	635	3.0	12.7
1997-98	22.87	0.4	12.98	-1.8	567	-2.1	11.3
1998-99	23.5	0.6	14.91	3.0	634	2.4	12.1
1999-00	21.12	-2.0	13.42	-2.0	635	0.0	16.1
2000-01	20.35	-0.7	11.08	-3.5	544	-2.9	12.5
2001-02	22.01	1.6	13.37	4.1	607	2.3	13.3
2002-03	20.50	-1.4	11.13	-3.4	543	-2.1	14.4
2003-04	23.46	2.9	14.91	6.8	635	3.4	13.6
2004-05	22.76	-0.6	13.13	-2.4	577	-1.8	13.9
2005-06	22.39	-0.3	13.39	0.4	598	0.7	15
2006-07**	23.76	1.2	14.11	1.1	594	-0.1	15.4
2007-08	23.63	-0.1	14.76	0.9	625	1.0	16.2
2008-09	22.09	-1.3	14.57	-0.3	660	1.1	16
2009-10	23.28	1.1	14.66	0.1	630	-0.9	16.2
2010-11	26.4	2.7	18.24	4.9	691	1.9	14.8
2011-12	24.46	-1.5	17.09	-1.3	699	0.2	16.1
2012-13	23.26	-1	18.34	1.5	788	2.5	18.6
2013-14	25.23	8.5	19.27	5.1	764	-3.1	19.70
2014-15	23.55	-6.7	17.15	-11.0	728	-4.7	NA
2015-16	24.91	5.8	16.35	-4.7	656	-9.9	NA
2016-17	29.45	18.2	23.13	41.4	785	19.6	NA
2017-18*	29.99	1.6	25.24	9.1	841	7.1	NA

Note: The yield rates given above have been worked out on the basis of production & area figure taken in ±000 units. ** Green Revolution period, N.A.- Not available.; 2017-18*- IVth Advance Estimate.

Source: Agricultural Statistics at a Glance, 2016. Directorate of Economics and Statistics, Ministry of Agriculture& FW, Govt. of India (Website http://www.dacnet.nic.in/ean).

2.8 Production Trends

2.8.1 Global Scenario: Crop-Wise

The total world acreage under pulses as recorded during 2014 is about 851.91 lakh ha with production at 774.73 lakh tones and productivity 909 kg/ha (Table -2.9).

In the world, pulses are grown by 198 countries. Beansdry was cultivated by 152 countries, which contributed about 35.93 % area to total world area, Chickpea by 58 contributed about 16.41%, Peasdry by 98 contributed 8.14%, Pigeonpea by 23 contributed 8.26 %, Lentil by 56 contributed by 5.31% and others 25.95%. The share to World production of Beans dry was 31.64% followed by Chickpea 17.72%, Peasdry 14.44%, Pigeonpea 6.31%, Lentil 6.23% and others 23.66%.

(Table-2.9): Global Ranking: Crop-wise

{Area-lakh ha, Production-lakh tonnes, Yield-kg/ha}

Crop	Area	% to Total	Production	% to Total	Productivity
Chickpea	139.81	16.41	137.31	17.72	982
Lentil	45.24	5.31	48.27	6.23	1067
Pigeon pea	70.33	8.26	48.90	6.31	695
Pea	69.32	8.14	111.86	14.44	1614
Beans	306.13	35.93	245.16	31.64	801
Total Pulses	851.91		774.73		909

Source: FAO Statistics 2014.




Fig.-2.5: Global Ranking: Crop-wise

2.8.2 Global Scenario: Total Pulses

The total world acreage under pulses as recorded during 2014 is about 851.91 lakh ha with production at 774.73 lakh tones and productivity 909 kg/ha. It reveals that the India ranked first in area and production with 35% and 25% respectively of world area and production. However, in case of productively Bahrain stood first with 18485 kg/ha. Thus it is also evident that the country¢s productivity at 660 kg/ha is far below the world average productivity of 909 kg/ha (Table-2.10).

(Table-2.10): Global Ranking: Total Pulses

{Area- lakh ha, Production-Lakh Tones, Yield-kg/ha}

Country	Area	% Cont.	Country	Prod.	% Cont.	Country	Yield
India	303.09	35.58	India	199.8	25.79	Bahrain	18485
Niger	54.7	6.42	Myanmar	59.77	7.72	Ireland	5886
Myanmar	42.03	4.93	Canada	58.28	7.52	Israel	5576
Nigeria	38.49	4.52	China	41.13	5.31	Belgium	4445
Brazil	32.09	3.77	Brazil	33.06	4.27	Tajikistan	3985
Canada	28.7	3.37	Ethiopia	26.13	3.37	Denmark	3952
China	23.85	2.80	USA	23.95	3.09	Trinidad &Tobago	3919
Tanzania	20.68	2.43	Russian	22.94	2.96	United Kingdom	3755
			Fed.				
Mexico	18.35	2.15	Australia	22.47	2.9	Netherlands	3639
Kenya	17.19	2.02	Nigeria	22.05	2.85	Switzerland	3638
Others	272.74	32.02	Others	265.15	34.22	Others	1068
World	851.91		World	774.73	World		909
					India		660

Source: FAO Statistics 2014.




Fig.-2.6: Global Ranking: Total Pulses

2.8.3 National Scenario: Plan Periods (Total Pulses)

A visit to different plan periods records a slight growth in total production and productivity from VIIIth Plan 1992-97 with 7% & 9% respectively. The area remained almost stagnant, stabilized up-till Xth plan. However, the XIIth plan analysis shows that the increasing per cent change trend under area (+ 5%) and production (+18%) of total pulses over previous plan periods (COPP) is given at (Table-2.11).

(Table – 2.11): Plan-Wise National Scenario- Total Pulses

(Area-Mha, Production-MTonnes, Yield-kg/ha)

Plan	Area	% COPP	Prod.	% COPP	Productivity	% COPP
First Plan (1951-56)	21.09		10.04		476	
Second Plan (1956-61)	23.71	12.42	11.75	17.03	496	4.10
Third Plan (1961-66)	23.86	0.63	11.14	-5.19	467	-5.79
Fourth Plan (1969-74)	22.21	-6.92	10.90	-2.15	491	5.11
Fifth Plan (1974-79)	23.32	5.00	11.71	7.43	502	2.32
Sixth Plan (1980-85)	23.08	-1.03	11.77	0.51	510	1.56
Seventh Plan (1985-90)	23.08	0.00	12.55	6.63	544	6.63
Eighth Plan (1992-97)	22.47	-2.64	13.34	6.29	594	9.18
Ninth Plan (1997-02)	21.97	-2.23	13.15	-1.42	599	0.82
Tenth Plan (2002-07)	22.44	2.14	13.35	1.52	595	-0.61
Eleventh Plan (2007-2012)	23.97	6.80	15.85	18.73	662	11.19
Twelfth Plan (2012-2017)	25.28	5.46	18.84	18.86	745	12.53

Source: DES, MoA&FW (DAC&FW), Govt. of India; % COPP is percentage change over previous plan.




Fig.-2.7: Total Pulses: Plan-Wise National Scenario

2.8.4 States' Scenario: Plan Analysis (Xth –XIIth)

- *Tenth Plan (2002-2007)*: The total pulse area in the country during the Tenth plan was 224.60 lakh hectares with a total production of 133.48 lakh tonnes. The same trend of pulses scenario was observed during the Tenth plan Madhya Pradesh ranked first in area (43.27 lakh hectares or 19.27 %) with a total production of 31.46 lakh tonnes or (23.57 % of the total production). While, Maharashtra was placed second with respect of area and its production 35.32 lakh hectares (15.73 %) and 19.98 lakh tonnes (14.97 %) followed by Rajasthan 31.77 lakh hectares (14.15 %) & 12.96 lakh tonnes (9.71 %).
- *Eleventh Plan (2007-2012)*: During Eleventh plan period the total pulses area and production were 239.75 lakh ha and 139.82 lakh tonnes respectively. Out of 239.75 lakh hectares about 53 % area under rabi and 46 % area under kharif were covered. However, approx. 59 % share of rabi production and 41 % share of kharif production exhibited in total pulse production in the country which explained the productivity of Rabi pulses is much higher than the kharif pulses. The state-wise analysis exhibited first rank to Madhya Pradesh, both in area and production with 47.75 lakh hectares and 35.98 lakh tonnes which was 19.92 % and 25.73 % respectively. Rajasthan ranked second in coverage with 16.90 % (40.51 lakh hectares) while at production front, state of Maharashtra ranked at second with 17.76 % (24.84 lakh tonnes) followed by Uttar Pradesh with 14.19 % (19.83 lakh tonnes). Maharashtra ranked third in area with 14.87% and Rajasthan ranked fourth in production with 14.00% (19.57 lakh hectares) of country@ production while in area, Karnataka stood at IVth rank with 10.05 % of country coverage during that XIth plan.
- Twelfth plan (2012-2017): In India, total pulse area and production irrespective of Twelfth Plan was 252.78 lakh hectares and 188.42 lakh tonnes respectively. Out of the total area, 57.52 lakh hectares is confined to Madhya Pradesh alone, earning a good pulse status and position contributing a remarkable 22.75% of the country total area and a production of 52.46 lakh tonnes, thereby ranking first both in area and production followed by Rajasthan in area (39.89 lakh hectares, 15.78% of the total area). While Rajasthan ranked third in production with 12.28% of the total pulse production and Maharashtra which ranked second (25.68 lakh tonnes or 13.63% of the total production); Uttar Pradesh was hardly placed at the forth rank in production (17.63 lakh tonnes or 9.36% of the total production). While Karnataka is on the forth rank in respect of area (25.74 lakh ha or 10.18%).

The overall area, production and productivity increasing trend during the last three plan period.

(Table-2.12): Plan-Wise StatesøScenarioóMajor States

{Area-lakh ha, Production-lakh tonnes, Yield-kg/ha}

States		Tenth Plan	% to	Eleventh	% to	*Twelfth	nes, 11eta-kg/na/ % to
States		1 (11111 1 11111	Country	Plan	Country	Plan	Country
A.P	A	19.71	8.78	19.76	8.24	12.70	5.03
	P	12.09	9.05	14.49	10.36	10.61	5.63
	Y	613		733		835	
Bihar	Α	6.5	2.9	5.79	2.42	5.16	2.04
	P	4.95	3.71	4.98	3.56	4.88	2.59
	Y	761		859		945	
Chhattisgarh	Α	9.06	4.03	8.52	3.55	8.80	3.48
	P	4.52	3.39	5.12	3.66	6.29	3.34
	Y	499		601		714	
Gujarat	Α	8.04	3.58	8.49	3.54	7.18	2.84
	P	5.14	3.85	6.74	4.82	6.47	3.44
	Y	639		794		902	
Haryana	Α	1.74	0.77	1.68	0.7	1.16	0.46
	P	1.26	0.94	1.33	0.95	0.91	0.48
	Y	725		792		781	
Jharkhand	Α	2.74	1.22	4.01	1.67	6.30	2.49
	P	1.72	1.29	3.1	2.21	6.24	3.31
	Y	626		772		991	
Karnataka	Α	20.78	9.25	24.09	10.05	25.74	10.18
	P	7.82	5.86	12.11	8.66	14.25	7.56
	Y	376		503		554	
Madhya	Α	43.27	19.27	47.75	19.92	57.52	22.75
Pradesh	P	31.46	23.57	35.98	25.73	52.46	27.84
	Y	727		754		912	
Maharashtra	Α	35.32	15.73	35.65	14.87	37.08	14.67
	P	19.98	14.97	24.84	17.76	25.68	13.63
	Y	566		697		693	
Orissa	Α	7.01	3.12	8.28	3.45	8.10	3.21
	P	2.81	2.11	3.88	2.78	4.27	2.27
	Y	401		469		528	
Punjab	Α	0.39	0.17	0.23	0.09	0.48	0.19
	P	0.32	0.24	0.19	0.14	0.42	0.22
	Y	811		858		872	
Rajasthan	Α	31.77	14.15	40.51	16.9	39.89	15.78
	P	12.96	9.71	19.57	14	23.14	12.28
	Y	408		483		580	
Tamil Nadu	A	5.45	2.43	5.97	2.49	7.74	3.06
	P	2.19	1.64	2.34	1.67	5.12	2.72
	Y	402	40	391		661	
U.P.	A	27.31	12.16	23.58	9.84	22.83	9.03
	P	22.37	16.76	19.83	14.19	17.63	9.36
W . 5	Y	819	1.02	841	0.70	773	4.0=
West Bengal	A	2.32	1.03	1.86	0.78	2.71	1.07
	P	1.75	1.31	1.47	1.05	2.53	1.34
	Y	754		787		934	
All India	A	224.6		239.75		252.78	
	P	133.48		139.82		188.42	
	Y	594		583		745	

Source: DES, DAC&FW,ND (*Twelfth plan - Avg. 2012-13 to 2016-17)




Fig.-2.8: Total Pulses Area: State@s Contribution (XIIth Plan)




Fig.-2.9: Total Pulses Production: State@s Contribution (XIIth Plan)

2.8.5 District Scenario (2016-17) – Potential Districts

The micro analysis at district level was also carried out and presented in (Table 2.13). The intra-state analysis revealed that Gulbarga district of Karnataka had the highest production with 3.45 per cent share to India followed by Latur of MH (2.86%) and Narsinghpur of MP (1.90%). In respect of area coverage, District-wise area, production and yield of top ten districts of India in respect of production are presented below which contributed 15.29% and 20.01% of area and production of the country.

The yield levels of potential district are also above the national average yield level except Vijayapur of Karnataka and Nagur & Bikaner district of Rajasthan Out of ten potential districts four districts was belongs to Madhya Pradesh State during 2016-17 & rest of the other were two from Karnataka, Maharshtra and Rajsthan.

(Table-2.13): Top Potential Districts óTotal Pulses

{Area- lakh ha, Production-Lakh Tones, Yield-kg/ha}

Name of	State	Aı	·ea	Produ	iction	Yie	eld
districts		Area	% to	Prod.	% to	Yield	ΥI
			India		India		
Gulbarga	Karnataka	6.49	2.20	7.98	3.45	1230	157
Latur	MH	3.55	1.20	6.63	2.86	1867	238
Narsinghpur	MP	2.71	0.92	4.4	1.90	1622	206
Vijayapur	Karnataka	5.66	1.92	4.25	1.83	751	96
Nagaur	Raj.	7.04	2.39	4.21	1.82	598	76
Bikaner	Raj.	6.51	2.21	3.97	1.71	609	78
Yavatmal	MH	3.41	1.15	3.86	1.66	1133	144
Damoh	MP	3.2	1.08	3.8	1.64	1185	151
Sagar	MP	3.58	1.21	3.79	1.63	1060	135
Satna	MP	2.88	0.97	3.4	1.47	1183	151
Total above		45.03	15.29	46.29	20.01	1028	131
All India		294.47		231.31		786	

Source: State Department of Agriculture.




Fig.-2.10: Total Pulses: Potential District

2.8.6 National Scenario: Plan Periods (Kharif Pulses)

A visit to different plan periods records a slight growth in total Area and production from X^{th} Plan 2002-07 with 4.3% and 4.8% respectively. Highly variability seen like sudden growh with area (+54%) and production (+47%) during VIIth plan and decreasing trend for area (-35%) and production (-33%) during IXth plan is given at (Table-2.14).

(Table-2.14): Plan-Wise National Scenario - Kharif Pulses

(Area-Million ha, Production-Million Tonnes, Yield-kg/ha)

Plan	Area	%	Prod.	%	Productivity	%
		COPP		COPP		COPP
Sixth Plan (1980-85)	10.53		4.48		425	
Seventh Plan (1985-90)	16.32	54.99	6.63	47.99	406	-4.51
Eighth Plan (1992-97)	16.10	-1.35	7.10	7.09	441	8.55
Ninth Plan (1997-02)	10.41	-35.34	4.71	-33.66	452	2.60
Tenth Plan (2002-07)	10.86	4.32	4.94	4.88	455	0.54
Eleventh Plan (2007-2012)	11.08	2.03	5.69	15.18	514	12.90
Twelfth Plan (2012-2017)	11.19	0.99	6.55	15.11	585	13.81

Source: DES, MoA&FW (DAC&FW), Govt. of India; % COPP is percentage change over previous plan.




Fig.-2.11: Kharif Pulses: Plan-Wise Scenario

2.8.7 States' Scenario: Plan Analysis (Xth -XIIth)

- *Tenth Plan (2002-2007)*: With a total coverage of 108.61 lakh hectares and a total production of 49.38 lakh tonnes, In area and production, Maharashtra ranked first with 24.58 lakh hectares (22.63%) and 13.54 lakh tonnes (27.42%) of the total area and production under kharif pulses in the country. Rajasthan trailed to second in area (21.97 lakh hectares) with 20.23% of the total kharif area. Rajasthan was the second largest producer with 6.06 lakh tonnes (12.27%) while Karnataka third in acreage of 13.59 lakh hectares (12.51%) during the plan period with a mere 4.81 lakh tonnes of production (9.74%), placed at third rank. The highest yield was recorded by the state of Bihar (910 kg/ha) followed by Punjab (778 kg/ha) and Haryana (645 kg/ha). Lowest yield observed in Rajasthan only (276 kg/ha).
- *Eleventh Plan (2007-2012):* The area and production under kharif pulses during twelfth plan were 110.78 lakh hectares and 56.94 lakh tonnes respectively. The state-wise contribution to total kharif pulses exhibited that the state stands first in acrerage were Rajasthan with 26.91 lakh ha (24.29%) followed by Maharashtra (19.85%), Karnataka (12.83%) and Madhya Pradesh (9.28%). While at production front, Maharashtra ranked first with 14.08 lakh tonnes which are 24.73% of country total kharif production. Karnataka stands third position with respect to production 6.43 lakh tones with (11.29%) followed by Madhya Pradesh (8.31%). The highest yield was recorded by the state of Bihar (1014 kg/ha) followed by Punjab (813 kg/ha) and Haryana (780 kg/ha) with the over all National yield average of (514 kg/ha). Lowest yield was observed in C.G. *i.e.*, 342 kg/ha only.
- Twelfth plan (2012-2017): The total area coverage and production of Kharif Pulses in India during the Twelfth plan was 111.93 lakh hectares and 65.52 lakh tonnes respectively, out of which Rajasthan ranked first (25.34 lakh hectares) and contributed 22.64% of total area while in production Maharashtra ranked first with 19.94% (13.06 lakh tonnes) and ranked second in area accounting for 18.29% (20.48 lakh hectares) of the total area. Madhya Pradesh ranked second with 16.29% of the country production (10.67 lakh tonnes) and stands third in area with 13.99% (15.66 lakh hectares) and Rajasthan stood third in production which accounted for 15.51% (10.16 lakh tonnes) of the total Kharif pulses during the period. In case of productivity, state of Bihar ranked first with (1180 kg/ha) followed by Jharkhand (915 kg/ha) and Gujarat (834 kg/ha). Lowest yield was observed in the state of Chhattisgarh (394 kg/ha).

Overall trend of area, production and yield of last three plan periods have shown increasing trend in production and productivity.

(Table-2.15): Plan-Wise States

ø Scenario - Major States

{Area-lakh ha, Production-lakh tonnes, Yield-kg/ha}

States		X th Plan	% to AI	XI th Plan	% to AI	*XII th Plan	nes, Yield-kg/ha) % to AI
A.P	Λ	9.08		7.21	6.51	2.99	2.67
A.P	A P		8.36				
	Y	3.89 428	7.87	3.23 448	5.67	1.67	2.54
D'1	1		0.02		0.62	556	0.51
Bihar	A	0.89	0.82	0.69	0.62	0.57	0.51
	P	0.81	1.63	0.7	1.23	0.67	1.02
	Y	910		1014		1180	
Chhattisgarh	A	2.35	2.17	2.22	2	2.08	1.85
	P	0.79	1.6	0.76	1.33	0.82	1.25
	Y	336		342		394	
Gujarat	Α	6.31	5.81	6.13	5.53	4.89	4.37
	P	3.79	7.67	4.48	7.87	4.07	6.22
	Y	601		731		834	
Haryana	Α	0.62	0.57	0.5	0.45	0.26	0.23
	P	0.4	0.81	0.39	0.68	0.18	0.27
	Y	645		780		695	
Jharkhand	Α	2.08	1.92	2.48	2.24	3.69	3.29
	P	1.22	2.46	1.67	2.93	3.37	5.15
	Y	587		673		915	
Karnataka	Α	13.59	12.51	14.21	12.83	13.72	12.26
	P	4.81	9.74	6.43	11.29	7.11	10.86
	Y	354	2111	452		518	2000
Madhya	A	9.5	8.75	10.28	9.28	15.66	13.99
Pradesh	P	4.45	9	4.73	8.31	10.67	16.29
Tracesir	Y	468		460	0.31	681	10.27
Maharashtra	A	24.58	22.63	21.99	19.85	20.48	18.29
Manarashra	P	13.54	27.42	14.08	24.73	13.06	19.94
	Y	551	27.42	640	24.73	638	17.74
Odisha	A	4.9	4.51	5.06	4.57	4.66	4.16
Odisila	P	1.92	3.89	2.38	4.18	2.51	3.83
	Y	392	3.69	470	4.10	538	3.63
Punjab	A	0.27	0.25	0.16	0.14	0.10	0.09
r unjao	P	0.27	0.23	0.10	0.14	0.10	0.09
	Y	778	0.43	813	0.23	779	0.12
Rajasthan	A	21.97	20.23	26.91	24.29	25.34	22.64
Kajasulali	P	6.06	12.27	9.66	16.97	10.16	15.51
	Y	276	12.27		10.97	401	13.31
Tamilnadu	A	1.88	1.73	359 1.61	1.45	2.34	2.09
Tammadu	P			0.79	1.43	1.59	2.42
	Y	0.77	1.56		1.39		2.42
Talangana		410		491	4.33	679 4.22	2 77
Telangana	A P			4.79			3.77
				1.93	3.39	2.36	3.60
IID	Y	0.0	0.1	402	7.65	558	7.00
U.P.	A	8.8	8.1	8.47	7.65	8.82	7.88
	P	5.59	11.31	5.74	10.08	5.55	8.46
Wast D. 1	Y	635	0.5	678	0.44	628	0.65
West Bengal	A	0.54	0.5	0.49	0.44	0.73	0.65
	P	0.35	0.71	0.33	0.58	0.48	0.74
	Y	648		673		663	
All India	A	108.61		110.78		111.93	
	P	49.39		56.94		65.52	
	Y	455		514		585	

Source: DES, DAC&FW,ND (*Twelfth plan - Avg. 2012-13 to 2016-17)




Fig.-2.12: Kharif Pulses Area: State@s Contribution (XIIth Plan)




Fig.-2.13: Kharif Pulses Production: State@s Contribution (XIIth Plan)

2.8.8 National Scenario: Plan Periods (Rabi Pulses)

A visit to different plan periods records a growth in VIIth, XIth, & XIIth Plan for area (32%, 11% & 9% and for production (36%, 21% & 21%). The area remained almost stagnant, stabilized up-till XIIth plan. However, the XIIth plan analysis shows that the increasing per cent change trend under area (+ 9%) and production (+21%) of total pulses over previous plan periods (COPP) is given at (Table-2.16). Decreasing trend seen only during VIIIth to IXth Plan.

(Table-2.16): Plan-Wise National Scenario ó Rabi Pulses

(Area-Million ha, Production-Million Tonnes, Yield-kg/ha)

Plan	Area	%	Prod.	%	Productivity	%
		COPP		COPP		COPP
Sixth Plan (1980-85)	12.62		7.33		581	,
Seventh Plan (1985-90)	16.72	32.49	10.03	36.83	600	3.28
Eighth Plan (1992-97)	15.25	-8.79	10.32	2.89	677	12.81
Ninth Plan (1997-02)	11.55	-24.26	8.43	-18.31	730	7.85
Tenth Plan (2002-07)	11.58	0.26	8.39	-0.47	725	-0.73
Eleventh Plan (2007-2012)	12.88	11.23	10.15	20.98	788	8.77
Twelfth Plan (2012-2017)	14.08	9.31	12.29	21.08	872	10.66

Source: DES, MoA&FW (DAC&FW), Govt. of India; % COPP is percentage change over previous plan.




Fig.-2.14: Rabi Pulses: Plan-Wise Scenario

2.8.9 States' Scenario: Plan Analysis (Xth -XIIth)

- Tenth Plan (2002-2007): Total area of (115.99 lakh hectares) and production of 84.10 lakh tonnes of Rabi pulses were observed during the Tenth plan in India. Out of these, 29.11% of area (33.77 lakh hectares) and 32.12% of production (27.01 lakh tonnes) were contributed by Madhya Pradesh alone which ranked first followed by Uttar Pradesh ranked second, could cover 15.96% of the total area (18.52 lakh hectares) and produce 19.95% of production (16.78 lakh tonnes) while with respect to area Maharashtra at third place could hardly cover 9.26% (10.74 lakh hectares) and in production Andhra Pradesh is in third position with 9.75% (8.20 lakh tones) of the country total Rabi pulse during the plan period. The highest state average yield exhibited in Punjab (917 kg/ha) followed by Uttar Pradesh (906 kg/ha) and Madhya Pradesh (800 kg/ha) has been above the National average yield of 725 kg/ha. The lowest yield and the plan period was recorded in Tamilnadu (398 kg/ha) followed by Karnataka (418 kg/ha) and Odisha (420 kg/ha).
- Eleventh plan (2007-2012): The total area and production under Rabi pulses during the Eleventh plan were 128.84 lakh hectares and 101.55 lakh tonnes respectively. Madhya Pradesh ranked first both in area and production with 37.47 lakh hectares and 31.24 lakh tonnes which are 29.08% and 37.77% of the countryøs total rabi pulse acreage and production respectively followed by Uttar Pradesh with 11.73 % and 13.88% (15.11 lakh hectares and 14.10 lakh tonnes) and Maharashtra was placed third in area & Production both as 10.60% (13.66 lakh hectares) and 10.59% (10.75 lakh tonnes) of the total Rabi pulses during the period. The highest state average yield exhibited in Punjab (1063 kg/ha) followed by Gujarat (960 kg/ha) and Andhra Pradesh (937 kg/ha) has been above the National average yield of 788 kg/ha. The lowest yield was recorded on the state of Tamilnadu 356 kg/ha followed by Odisha (466 kg/ha) and Karnataka (574 kg/ha).
- Twelfth plan (2012-17): All India Rabi pulse acreage and production has been recorded at 140.84 lakh hectares and 122.90 lakh tonnes. Madhya Pradesh stood at first in area and production, covering 41.86 lakh hectares (29.72%) with a production of 41.80 lakh tonnes (33.01%). Maharashtra ranked second with 16.60 lakh hectares of area (11.79%), Rajasthan with third position which covered 12.98 lakh hectares of area (10.56%) and UP shared the fourth rank having an area of 14.00 lakh hectare (9.94%). However, Rajasthan stood second in terms of production contributing 12.98 lakh tonnes (10.56%), Maharshtra in third position with a production of 12.62 lakh tonnes (10.27%) and UP in fourth position 12.09 lakh tonnes (9.84%), (Table 1.16). The highest productivity was recorded in plan period in Telangana (1315 kg/ha) followed by Jharkhand (1098 kg/ha) and Gujarat (1047 kg/ha). The lowest yield was noticed in the state of Odisha (513 kg/ha).

Overall trend of area, production and yield of last three plan periods have shown increasing trend in area however, production and productivity declined during XI^{th} plan period from X^{th} plan period and significantly increased during XII^{th} plan period.

(Table-2.17): Plan-Wise StatesøScenarioóMajor States

{Area-lakh ha, Production-lakh tonnes, Yield-kg/ha}

		th				ction-lakn tonnes	
States		X th Plan	% to AI	XI th Plan	% to AI	*XII th Plan	% to AI
A.P	A	10.63	9.16	10.99	8.53	9.71	6.89
	P	8.2	9.75	10.31	10.15	8.94	7.28
	Y	771		937		921	
Bihar	A	5.61	4.84	5.1	3.96	4.60	3.26
	P	4.15	4.93	4.28	4.21	4.21	3.43
	Y	740		839		917	
Chhattisgarh	A	6.7	5.78	6.3	4.89	6.73	4.78
	P	3.73	4.44	4.36	4.29	5.47	4.45
	Y	557		692		813	
Gujarat	A	1.73	1.49	2.36	1.83	2.29	1.63
	P	1.35	1.61	2.27	2.23	2.40	1.95
	Y	780		960		1047	
Haryana	A	1.12	0.97	1.18	0.91	0.90	0.64
-	P	0.86	1.02	0.94	0.93	0.73	0.59
	Y	768		798		806	
Jharkhand	A	0.66	0.57	1.53	1.19	2.61	1.85
	P	0.5	0.6	1.42	1.4	2.87	2.33
	Y	758		931		1098	
Karnataka	A	7.2	6.21	9.88	7.67	12.01	8.53
	P	3.01	3.58	5.67	5.59	7.14	5.81
	Y	418		574		594	
Madhya Prd.	A	33.77	29.11	37.47	29.08	41.86	29.72
J	P	27.01	32.12	31.24	30.77	41.80	34.01
	Y	800		834		998	
Maharashtra	A	10.74	9.26	13.66	10.6	16.60	11.79
	P	6.44	7.66	10.75	10.59	12.62	10.27
	Y	600		787		760	
Odisha	A	2.12	1.83	3.22	2.5	3.44	2.44
	P	0.89	1.06	1.5	1.48	1.77	1.44
	Y	420		466		513	
Punjab	A	0.12	0.1	0.06	0.05	0.38	0.27
3	P	0.11	0.13	0.07	0.07	0.34	0.28
	Y	917		1063		902	
Rajasthan	A	9.81	8.46	13.6	10.55	14.54	10.33
5	P	6.9	8.21	9.91	9.76	12.98	10.56
	Y	703		729		892	
Tamilnadu	A	3.57	3.08	4.36	3.39	5.41	3.84
	P	1.42	1.69	1.55	1.53	3.53	2.87
	Y	398		356		653	
Telangana	A			1.49	1.15	1.28	0.91
	P			1.42	1.4	1.68	1.37
	Y			957		1315	
U.P.	A	18.52	15.96	15.11	11.73	14.00	9.94
	P	16.78	19.95	14.1	13.88	12.09	9.84
	Y	906		933		863	
West Bengal	A	1.78	1.54	1.38	1.07	1.98	1.40
-	P	1.4	1.66	1.13	1.12	2.05	1.66
	Y	787		822		1034	
All India	A	115.99		128.84		140.84	
	P	84.1		101.55		122.90	
	Y	725		788		873	

Source: DES, DAC&FW,ND (*Twelfth plan - Avg. 2012-13 to 2016-17)




Fig.-2.15: Rabi Pulses Area: State@ Contribution (XIIth Plan)




Fig.-2.16: Rabi Pulses Production: State@s Contribution (XIIth Plan)

CHICKPEA (GRAM)





Chapter-3 Chickpea (Gram)

Botanical Name : Cicer arietinum L.

Synonym : Chickpea, Bengalgram, Chana and Gram

Origin : South West Asia ó probably Afganisthan and/or Persia

Chromosome nos. : 2n = 16

3.1 Economic Importance

In India chickpea is the most important pulse crop and mostly grown under rainfed conditions (65%), remaining 35% with critical irrigation support. During XIIth Plan (2012-2017), it was grown in > 35 per cent of total pulse area (252.43 Lha) contributing to about 45% of total pulse production. It is mainly consumed as \oplus Dalø(split cotyledons) and *chole*. Many attractive dishes viz.,ó sweets, snacks and namkeen are also prepared from its flour called besan. Also eaten as whole fried or boiled and salted. Fresh green leaves (sag) are used as vegetables and green grains as *hare chole or cholia*. Straw of gram is an excellent fodder while both husk and bits of \oplus Dalø are valuable cattle feed. Leaves consist of mallic and citric acid and are very useful for stomach ailments and blood purifier.

3.2 Nutritive Value

Protein	18-22%	Calcium	280 mg/100 g
Carbohydrate	61-62%	Iron	12.3 mg/100 g
Fat	4.5 %	Phosphorus	301 mg/100
Fibre	22-23%	Calorific value	368-396 Kcal/100 g

Agronomic Significance: Leaving about 30-50 kg N/ha for successive crops, especially cereals. Intercrop cereals also get benefited through \pm Nø supplied by way of fixation in gram.

3.3 Growth Rate

From 1980-81 to 2009-10, the total acreage under pulses has almost slightly (\pm) being showed, however, the maximum growth rate in area was recorded during 2013-14 and 2016-17 with growth rate of 16% & 14%. Maximum production growth rate of 32% and maximum yield growth rate of 16% were also observed during 2016-17. The highest area (10.57 million ha) and production (11.16 million tonnes) was recorded during 2016-17 followed by 9.93 million ha & 9.53 million tonnes during 2013-14 (Table 3.1).

(Table-3.1): Growth rate of chickpea

(A- Million ha, P- Million tones, Y-kg/ha, Growth Rate (GR) - %)

Year			Chick		ng/na, Growin i	
	Area	GR	Production	GR	Yield	GR
1980-81	4.33		6.58		1522	
1985-86	5.79	33.7	7.80	18.5	1348	-11.4
1990-91	5.36	-7.5	7.52	-3.6	1404	4.1
1995-96	4.98	-7.0	7.12	-5.4	1429	1.8
1996-97	6.85	37.5	5.57	-21.7	814	-43.1
1997-98	7.56	10.5	6.13	10.1	811	-0.3
1998-99	8.47	12.0	6.80	10.9	803	-1.0
1999-00	6.15	-27.4	5.12	-24.7	833	3.7
2000-01	5.19	-15.6	3.86	-24.7	744	-10.7
2001-02	6.42	23.7	5.47	42.0	853	14.7
2002-03	5.91	-7.9	4.24	-22.6	717	-15.9
2003-04	7.05	19.3	5.72	34.9	811	13.1
2004-05	6.71	-4.7	5.47	-4.3	815	0.4
2005-06	6.93	3.2	5.60	2.4	808	-0.7
2006-07*	7.49	8.2	6.33	13.1	845	4.5
2007-08	7.54	0.7	5.75	-9.2	762	-9.8
2008-09	7.89	4.6	7.06	22.8	895	17.4
2009-10	8.17	3.5	7.48	5.9	915	2.3
2010-11	9.19	12.4	8.22	10.0	895	-2.2
2011-12	8.30	-9.7	7.70	-6.3	928	3.7
2012-13	8.52	2.7	8.83	14.7	1036	11.7
2013-14	9.93	16.5	9.53	7.9	960	-7.4
2014-15	8.25	-16.9	7.33	-23.0	889	-7.4
2015-16	8.40	1.8	7.06	-3.7	840	-5.4
2016-17	9.63	14.6	9.38	32.8	974	16.0
2017-18*	10.56	9.7	11.23	19.7	1063	9.1

Source: DES, Ministry of Agri. &FW (DAC&FW), Govt. of India; 2017-18*- IVth Adv. Est.

(Table-3.2): Per cent share to total pulses

(Area- lakh ha, Production-Lakh Tones, Yield-kg/ha)

Year		Gram		To	tal Pulses	i 114, 1 1	% Share to Total Pulses		
	A	P	Y	A	P	Y	A	P	YI
1980-81	43.28	65.85	1522	224.45	106.24	473	19.3	62.0	321
1990-91	53.56	75.21	1404	372.55	203.68	547	14.4	36.9	257
1995-96	49.79	71.16	1429	331.77	170.10	513	15.0	41.8	279
2000-01	51.85	38.55	744	203.48	110.75	544	25.5	34.8	137
2001-02	64.16	54.73	853	220.08	133.68	607	29.2	40.9	140
2002-03	59.06	42.37	717	204.96	111.25	543	28.8	38.1	132
2003-04	70.48	57.18	811	234.58	149.05	635	30.0	38.4	128
2004-05	67.15	54.69	815	227.63	131.30	577	29.5	41.7	141
2005-06	69.26	56.00	808	223.91	133.84	598	30.9	41.8	135
2006-07	74.94	63.34	845	231.92	141.98	612	32.3	44.6	138
2007-08	75.44	57.49	762	236.33	147.62	625	31.9	38.9	122
2008-09	78.93	70.6	895	220.94	145.66	659	35.7	48.5	136
2009-10	81.69	74.76	915	232.82	146.62	630	35.1	51.0	145
2010-11	91.86	82.21	895	264.02	182.41	691	34.8	45.1	130
2011-12	82.99	77.02	928	244.62	170.89	699	33.9	45.1	133
2012-13	85.22	88.32	1036	232.57	183.42	789	36.6	48.2	131
2013-14	99.27	95.26	960	252.18	192.55	764	39.4	49.5	126
2014-15	82.51	73.32	889	235.53	171.52	728	35.0	42.7	122
2015-16	83.99	70.58	840	249.11	163.48	656	33.7	43.2	128
2016-17	96.26	93.78	974	294.47	231.31	785	32.6	40.5	124
2017-18*	105.61	112.29	1063	299.93	252.35	841	35.2	44.5	126

Source: DES, Ministry of Agri. &FW (DAC&FW), Govt. of India; 2017-18*- IVth Adv. Est.

3.4 Import/Export

• Import: The import of pulses in India during April, 2014 to March, 2015 was 4.19 lakh tonnes worth Rs.1334.96 crores against the value of Rs.17196.87 crore for total foodgrains, Rs.121319.02 crore for total agricultural imports and against Rs.2737086.58 crore for total National Import. The import during April, 2015 to March, 2016 was 10.31 lakh tonnes worth Rs.4453.72 crore against the import value of Rs. 26841.87 crore for total foodgrains, Rs.140288.69 crore for total agricultural import and Rs.2490298.08 crore for total National import respectively during this period. The share of Chickpea import to Agricultural import was 1.10% and 3.17% respectively during April, 2014 to March, 2015 and April, 2015 to March, 2016.

{Chickpeas contribute the single largest share in India's export basket of pulses registering 85.64% and 84.87% share in the total pulses export during 2014-15 and 2015-16 respectively}.

• Export: The pulses export of the country during April, 2014 to March, 2015 was 1.90 lakh tonnes worth Rs.1021.57 crore against the value of Rs. 59500.54 crore for total foodgrains, Rs.239681.04 crore for total agricultural exports and against Rs.1896445.47 crore for total National export. The export during April, 2015 to March, 2016 was 2.17 lakh tonnes worth Rs.1337.64 crore against the export value of Rs. 42622.29 crore for total foodgrains, Rs.215395.68 crore for total agricultural export and Rs.1716378.05 crore for total National export respectively during this period. The share of Chickpea export to Agricultural export was 0.43% and 0.62% respectively during April, 2014 to March, 2015 and April, 2015 to March, 2016.

(Table-3.3): Importing & Exporting Countries

(Quantity-Thousand Tonnes)

S.No.	I	mport		\	Export	
S.NO.	Country	Aveg.*	% Share	Country	Aveg.*	% Share
1	Australia	376.61	53.73	Pakistan IR	63.45	31.00
2	Austria	131.79	18.80	Algeria	32.91	16.08
3	Russia	76.87	10.97	Turkey	24.70	12.07
4	Sri Lanka DSR	32.49	4.64	Sri Lanka DSR	14.77	7.21
5	Tanzania Rep	18.11	2.58	U Arab EMTS	11.26	5.50
6	Thailand	12.63	1.80	Saudi Arab	9.00	4.40
7	Mexico	12.23	1.74	Spain	5.40	2.64
8	USA	6.89	0.98	Tunisia	5.31	2.59
9	Myanmar	6.31	0.90	Egypt ARP	4.48	2.19
10	Ethiopia	5.72	0.82	Libya	3.94	1.92
11	Canada	4.96	0.71	Iraq	3.43	1.67
12	Argentina	2.30	0.33	Kuwait	2.99	1.46
13	Mozambique	2.15	0.31	Malaysia	2.17	1.06
14	Sudan	2.14	0.31	Jordan	1.99	0.97
15	Uzbekistan	1.95	0.28	France	1.88	0.92
16	El Salvador	1.93	0.27	Iran	1.79	0.88
	Ukraine	1.91	0.27	Vietnam Soc.	1.75	0.85
				Rep		
17	Others	10.78	1.54	Others	5.27	2.57
	Total	700.95		Total	204.70	

Source: Ministry of Commerce and Industry; Aveg. *- 2012-13 to 2016-17

3.5 Production Trends

3.5.1 Global Scenario

India ranks Ist in area and production in the world, followed by Pakistan, Iran and Australia. The highest productivity of 3759 kg/ha is observed in China followed by Israel, Repbl. of Moldova and Bosnia & Herzegovina. Indiaøs productivity was 995 kg/ha.

(Table-3.4): Global Ranking: Major Countries

{Area- Lakh ha, Production-Lakh Tonnes, Yield-kg/ha}

Country	Area	%	Country	Prod.	%	Country	Yield
·		Cont.	•		Cont.	•	
India	99.27	71.00	India	98.80	71.95	China	3759
Pakistan	9.50	6.79	Australia	6.29	4.58	Israel	3559
Iran	5.94	4.25	Myanmar	5.62	4.09	Repbl of	
						Moldova	3556
Australia	5.08	3.63	Ethiopia	4.59	3.34	Bosnia	
			-			&Herzegovina	3204
Turkey	3.88	2.78	Turkey	4.50	3.28	Yemen	3093
Myanmar	3.84	2.75	Pakistan	3.99	2.91	Jordan	2632
Ethiopia	2.40	1.71	Iran	2.62	1.91	Hungary	2250
Uni. Rep.	1.16	0.83	Mexico	1.72	1.25	Uzbekistan	2192
Tanzania							
Malawi	1.15	0.82	USA	1.27	0.93	Egypt	2124
World	139.81		World	137.31		World	982
						India	995

Source: FAO Statistics 2014.




Fig.-3.1: Global Scenario: Chickpea

3.5.2 National Scenario: Plan Periods

Plan-wise performance revelaed, the highest area and production increased during XIth plan over the previous plan period. The details plan-wise performance is given below:

(Table-3.5): Plan-Wise National Scenario

(Area-Million ha, Production-Million Tonnes, Yield-kg/ha)

Plan	Area	%	Prod.	%	Yield	%
		COPP		COPP		COPP
Fourth Plan (1969-74)	7.65		4.89		639	
Fifth Plan (1974-79)	6.71	-12.29	5.52	12.88	823	28.70
Sixth Plan (1980-85)	7.18	7.00	4.71	-14.67	656	-20.26
Seventh Plan (1985-90)	6.73	-6.27	4.66	-1.06	692	5.55
Eighth Plan (1992-97)	6.86	1.93	5.28	13.30	770	11.16
Ninth Plan (1997-02)	6.76	-1.46	5.48	3.79	811	5.32
Tenth Plan (2002-07)	6.82	0.89	5.47	-0.18	802	-1.06
Eleventh Plan (2007-2012)	8.22	20.53	7.24	32.36	881	9.82
Twelfth Plan (2012-2017)	8.95	8.88	8.43	16.43	941	6.81

Source: DES, Twelfth Plan (2012-2017); % COPP is percentage Change over previous plan




Fig.-3.2: Chickpea: Plan-Wise Scenario

3.5.3 State's Scenario: Plan Analysis (Xth -XIIth)

- *Tenth Plan (2002-2007)*: A total of 68.18 lakh ha of area and 54.72 lakh tonnes of gram production were observed in the country during the plan. Madhya Pradesh ranked first in terms of area and production (38.23% and 42.52%) followed by Maharashtra with 13.93% and 10.84% and Rajasthan with 13.77% and 11.60%, respectively. Andhra Pradesh has recorded and yield of 1147 kg/ha followed by Bihar with 913 kg/ha. The lowest yield observed in Karnataka (479 kg/ha) followed by Maharashtra (624 kg/ha) and Odisha (645 kg/ha).
- *Eleventh plan (2007-12)*: The total area and production of gram in the country were 82.18 lakh hectares and 72.42 lakh tonnes respectively. Madhya Pradesh outstanding position in area coverage and production of gram (35.34% and 38.12%) of the total area and production of the country), followed by Rajasthan (16.04% and 13.12%) and Maharashtra (15.27% and 14.14%) respectively. The highest productivity was recorded in plan period in Andhra Pradesh (1270 kg/ha) followed by Bihar (1067 kg/ha) and Gujarat (1048 kg/ha). The lowest yield was noticed in the state of Karnataka (600 kg/ha).
- *Twelfth Plan (2012-2017)*: The total area and production of gram during twelfth Plan was 89.45 lakh hectares and 84.25 lakh tonnes respectively. Madhya Pradesh ranked first contributing an area of (34.39% and 40.32% of total area and production of country). Maharashtra is on the second rank for area 15.48 lakh ha (17.30%) and third for production 12.12 lakh tones (14.39%). Whereas, Rajasthan stood second in production (14.43%) and third in area (15.48%). The highest yield was recorded in the state of Telangana (1474 kg/ha) followed by Gujarat (1178 kg/ha) and West Bengal (1148 kg/ha). The lowest yield was recorded in Karnataka (619 kg/ha).

Chickpea is a major pulse in India which contributed about 35% of area & 45% of total pulse production. Overall trend of area, production and yield for the last three plan periods has shown a significant increase.

(Table-3.6): Plan-Wise StatesøScenario: Major States

{Area- Lakh ha, Production-Lakh Tonnes, Yield-kg/ha}

State		X th Plan	% to AI	XI th Plan	% to AI	*XII th Plan	% to AI
Andhra Pradesh	A	4.30	6.30	6.07	7.38	4.50	5.03
	P	4.93	9.01	7.71	10.65	4.93	5.86
	Y	1147		1270		1096	
Bihar	A	0.69	1.01	0.60	0.72	0.60	0.67
	P	0.63	1.15	0.64	0.88	0.68	0.80
	Y	913		1067	0.00	1124	0.00
Chhattisgarh	A	2.07	3.04	2.44	2.97	2.87	3.21
eminus gum	P	1.54	2.81	2.22	3.06	2.74	3.25
	Y	744	2.01	910	0.00	953	0.20
Gujarat	A	1.49	2.18	1.88	2.28	1.72	1.92
Gujarat	P	1.23	2.25	1.97	2.72	2.02	2.40
	Y	826	2.23	1048	2.72	1178	2.10
Haryana	A	1.05	1.53	1.01	1.23	0.55	0.61
Tiai yana	P	0.79	1.44	0.85	1.18	0.48	0.57
	Y	752	1.77	842	1.10	876	0.57
Karnataka	A	4.95	7.26	8.13	9.89	10.55	11.80
Karnataka	P	2.37	4.33	4.88	6.74	6.54	7.76
	Y	479	4.55	600	0.74	619	7.70
Madhya Pradesh		26.06	38.23	29.04	35.34	30.76	34.39
Madnya Pradesh	A P	23.26					
	Y		42.52	27.6 950	38.12	33.97	40.32
M-1 1. (893	12.02		15.07	1104	17.20
Maharashtra	A	9.50	13.93	12.55	15.27	15.48	17.30
	P	5.93	10.84	10.24	14.14	12.12	14.39
0.11.1	Y	624	0.45	816	0.40	783	0.40
Odisha	A	0.31	0.45	0.40	0.49	0.43	0.48
	P	0.20	0.36	0.29	0.41	0.33	0.39
- · · ·	Y	645	10.55	725	1.504	770	15.40
Rajasthan	A	9.39	13.77	13.18	16.04	13.85	15.48
	P	6.35	11.60	9.50	13.12	12.16	14.43
	Y	676		721		878	
Tamilnadu	Α	0.06	0.09	0.07	0.09	0.07	0.07
	P	0.04	0.08	0.05	0.07	0.04	0.05
	Y	667		714		652	
Telangana	Α			0.97		0.91	1.02
	P			1.10		1.35	1.60
	Y			1135		1474	
Uttar Pradesh	Α	7.57	11.11	5.65	6.87	5.14	5.74
	P	6.80	12.43	5.32	7.35	4.62	5.48
	Y	898		942		899	
West Bengal	A	0.41	0.60	0.23	0.28	0.27	0.30
	P	0.37	0.67	0.24	0.33	0.31	0.37
	Y	902		1043		1148	
All India	A	68.18		82.18		89.45	
	P	54.72		72.42		84.25	
	Y	803		881		942	

Source: DES, DAC&FW, ND (*Twelfth plan - Avg. 2012-13 to 2016-17)




Fig.-3.3: Chickpea Area: State@ Contribution (XIIth Plan)




Fig.-3.4: Chickpea Production: State® Contribution (XIIth Plan)

3.5.4 District Scenario (2016-17) – Potential Districts

The intra-regional analysis at the district level as depicted in (Table -3.7) revealed the highest contribution in production of Latur, MH (2.66%) followed by Sagar (2.63%) and Damoh (2.59%) of M.P. State. District-wise area, production and yield of top ten district of India in respect of production are presented below which contributed 16.59 per cent and 22.40 per cent of total area and production of chickpea in the country.

The yield was revealed that the potential districts yield is higher than the National average yield (974 kg/ha). Out of Ten potential districts six belongs to Madhya Pradesh whereas, two of Maharashtra, one each belongs from Karnataka and Rajasthan.

(Table-3.7): Top Potential Districts: Chickpea

{Area-Lakh ha, Production-Lakh Tonnes, Yield-kg/ha}

Name of	State		Area	Produ	uction	Yi	eld
districts		Area	% to India	Prod.	% to India	Yield	YI
Latur	MH	2.10	2.18	2.50	2.66	1190	122
Sagar	MP	1.78	1.85	2.47	2.63	1386	142
Damoh	MP	1.74	1.80	2.43	2.59	1400	144
Gulbarga	Karnataka	1.85	1.92	2.17	2.32	1174	121
Satna	MP	1.20	1.25	2.10	2.24	1750	180
Vidisha	MP	1.72	1.79	1.99	2.13	1157	119
Bikaner	Raj.	2.27	2.35	1.94	2.06	854	88
Narsinghpur	MP	0.98	1.02	1.88	2.00	1911	196
Raisen	MP	1.20	1.25	1.79	1.91	1489	153
Nanded	MH	1.14	1.19	1.75	1.87	1533	157
Total above		15.98	16.60	21.01	22.41	1315	135
All India		96.26		93.78		974	

Source: State Department of Agriculture.




Fig.-3.5: Chickpea: Potential Districts

(Table-3.8): State-wise yield gap and additional return

(Yield: kg/ha; Return: Rs./ha.)

State			Gap o		Gap over SAY		Yield 2016- 17	Additional return by bridging yield		
									gap (l	Rs/ha)
	IP	FP	SAY	Actual	%	Actual	%		FP	SAY
AP	1651	1553	1085	98	6	566	52	960	3920	22640
Assam	1082	749	692	333	44	390	56	651	13320	15600
Bihar	2063	1798	1159	265	15	904	78	1120	10600	36160
Haryana	2135	1803	848	332	18	1287	152	1189	13280	51480
UP	1518	1522	916	-4	0	602	66	1114	-160	24080
Maharashtra	1441	1125	752	316	28	689	92	891	12640	27560
Gujarat	1620	1379	1179	241	17	441	37	1116	9640	17640
Uttarakhand	2000	1552	868	448	29	1132	130	1000	17920	45280
Tamil Nadu	887	741	648	146	20	239	37	648	5840	9560
Rajasthan	1690	1446	842	244	17	848	101	911	9760	33920
Chhattisgarh	981	911	912	70	8	69	8	1171	2800	2760
Manipur	1142	814	895	328	40	247	28	0.0	13120	9880
MP	1432	1052	1100	380	36	332	30	1100	15200	13280
Karnataka	1459	1231	619	228	19	840	136	590	9120	33600
Punjab	1598	1453	1191	145	10	407	34	1375	5800	16280
J & K	794	619	558	175	28	236	42	0	7000	9440
WB	2033	1402	1157	631	45	876	76	1013	25240	35040
Average	1502	1244	907	257	21	594	66	990	10296	23776
Rice fallow c	hickpea									<u>.</u>
Assam	1257	901	692	356	40	565	82	651	14240	22600
Bihar	1346	1037	1159	309	30	187	16	1120	12669	7667
Manipur	1298	950	895	348	37	403	45	0	14616	16926
West	1198	950	1157	248	26	41	4	1013	10664	1763
Bengal	1170	930	1137	240	20	41	4	1013	10004	1/03
Average	1275	960	976	315	33	299	31	928	13871	13145

Source-Annual Report- 2016-17, GoI, DPD, Bhopal (Ave. 2013-14 to 2015-16).

State Average Yield - E&S (Ave. 2011-12 to 2015-16)

IP: Improved Practise FP: Farmers Practise SAY: State Avergae Yield.

3.6 Chickpea Agronomy

3.6.1 Economic Classification

- **Desi or brown gram** (*C arietinum L*.): Colour of seed coat ranges yellow to dark brown, usually small in seed size but some large size varieties are also found. Plants are small with good branching ability, especially suited for late sown and rice fallow areas (Ch.No. 2n = 14, 16). Most widely cultivated group in India (90% of total world).
- **Kabuli or white gram** (*C. Kabulianum*): Colour of seed coat white with bold and attractive seed size. Plants are taller than desi and stand more or less erect. Yield potential is poor on account of poor branching as compare to :desiø (Ch.No. 2n = 16). Some small seeded white coloured grain also comes under this category.

3.6.2 Botanical Description

It is a small herbaceous branched plant with maximum height of 45-60 cm. Roots include a strong central tap root with extensive lateral branches spread out in all directions in upper soil layers. There are numerous bacterial nodules found on primary and secondary roots, used as a site of atmospheric $\pm N\phi$ fixation. Stem is branched with numerous granular hairs on it. Flowers are typical Papilionaceous consisting of five sepals, five petals (consists of one *standard*, two *wings* and two *keels*), ten stamens (nine fused to form one staminal column and one free) and a *carpel* with the *style* borne laterally on the *ovary*. They are singly, axillary, racemes usually solitary having pink or white colour with pink or blue shades. Anthesis takes place between 9 am to 3 pm. Self- fertilization before opening is the rule but cross pollination upto 5-10% extent also takes place by insect. Pods are rectangular, swollen structure, about 2 cm long and usually contain 2 seed in them. A single plant produces 50-150 pods. Seeds are spherical in shape, wrinkled or smooth with a pointed beak. *Its head is similar to chicken's head with a characteristic 'beak' hence called as chickpea*.

3.6.3 Production Technology

- Climate: Being a winter season legume, it requires fairly cold and dry climate. However, severe cold and frost, especially at flowering/pod initiation stage, are injurious for developing flowers into seed. The crop is well suited to the areas having 600-900 mm rainfall. Excessive rains, soon after sowing or at flowering and fruiting or hailstorms at ripening, cause heavy losses.
- Soil & Field preparation: Grown in a wide range of soils viz light sandy loam to moderately heavy loam in north to black cotton soils of Central Plateau. However, sandy loams to clay loam soil, free from excessive salt and neutral in reaction with drainage facility, are best for gram. Soils having >8.5 P^H. It requires clodded and rough seed bed for good aeration in root zone, obtained by one deep ploughing and a cross harrowing.
- Sowing time: Rainfed –Ist fortnight of October in central and south India and IInd fortnight of October in North India. Irrigated–Ist fortnight of November in North India and IInd fortnight of October in central and southern India; Late sowing– Ist week of December in rice fallows of NEPZ or in irrigated conditions, where field are vacated very late by kharif crops.
- Seed Rate & Spacing: Small seededó50-60 kg/ha; Bold seeded and late sowingó80-90 kg/ha (Small seeded varieties are recommended for late sown conditions). Rainfedó 30 cm x 10 cm; Irrigatedó 45 cm x 10 cm; Late sowing ó 25 cm x 10 cm.
- Seed treatment: Treat the seed with thiram or carbendazim @ 2 g/kg of seed before 3 days of sowing followed by seed inoculation with a solution/jaggery having dual culture of Rhizobium and PSB. One packet of both the culture is enough for 10kg seed. Seed treatment with rhizobium @ 5 g per kg seed and soil inoculation of phosphate solubilizing bacteria @ 500 g per ha by mixing with 50 kg well decompose FYM just at

the time of sowing improves the FUE. For correcting Zn deficiency, foliar spray of 0.5 kg ZnSO₄ with 0.25 kg lime or soil application of ZnSO₄ @ 25 kg per ha to one crop on Zn deficient soils is helpful to both the crop of pulse based cropping system. Mo deficiency can be corrected by applying 1 kg sodium molybdate per ha and for boron deficient soils foliar spray of B @ 1.0 ó 1.5 kg B per ha or soil application of 4 kg borax. Spray 1.0 per cent FeSO₄ to recoup the crop from Fe deficiency.

• Varieties: Selection of the varieties should be as per the adaptability to the region, availability of resources, time of sowing. etc. Short duration varieties like JG 16, JG 11, JAKI 9218, JG 14, Vijay, Vikas, Vishal, JGK-1, KAK 2 etc. are popular/helped in expanding chickpea area in southern and central part of the country. The yield potential of these early varieties remained almost similar to long duration varieties. The ofther recent released varieties of short durations namely Rajas, Pusa 547, RVG 202, RVG 203, JAKI 9218, JGK 1, KAK 2, Shubhra may be grown for better yield.

A number of Fusarium wilt resistant varieties which helps to reduce losses due to diseases and stabilise chickpea yield like GNG 1581, CSJ 515, RVG 202, JGK 5, JG 6, JG 16, Digvijay, Gujarat Gram 3, BG 391, Ujjawal, GLK 26155, HK 05-169, BDNG 797, RSG 991 etc may be adopted by the farmers.

The varieties for specific situations such as **Kabuli** ó KAK-2 (>40 g/100seeds); Pusa Chamatkar (BG-1053), ICCV-2, Pusa Kabuli 1003 (BG-1003), JGK-1, Haryana Kabuli Chana-1; **Rice-Chickpea cropping system** (late sown up to end of December) ó Udai (KPG-59), Pusa-372, RSG-963, PBG-1, Pant G-186 and JG-74; **mild saline soil** ó Karnal Chana 1 (CSG-8962);**drought prone areas**óRSG-888, Annegiri;**high fertile and high rainfall/irrigated areas** ó DCP-92-3 are also recommended.

- Cropping system: Gram in rotation with cereal crops helps in controlling soil-borne diseases. The most common cropping system based on chickpea are as follows-Rotation: Kharif fallow ó Gram, Paddy ó Gram, Maize ó Gram, Bajra ó Gram and Jowar ó Gram; Inter cropping: Chickpea + Mustard (2:1 to 4:1); Chickpea + Linseed (2:2); chickpea + wheat/ Barley (2:2), Chickpea + Safflower (2:2) and Chickpea + Coriander (2:2).
- Water management: -Gramø grown as rainfed crop in general in India, invariably suffers from moisture stress as -terminal drought, at most critical pod development stage due to high atmospheric and soil temperature coupled with high wind velocity. So, to minimize transpiration loss and conserving residual soil moisture for longer time, a foliar spray of 2% KCl is giving promising results.

However, under assured irrigation, one irrigation each at maximum branching and pod development resulted in 25-70% increase in yield in absence of winter rain. In no case, irrigation should be given earlier than four weeks after sowing and during active flowering because earlier situation is harmful for maximum $\pm N\phi$ fixation as the Rhizobial bacteria work only in aerobic conditions and later, excess irrigation may reverse the crop

again to vegetative phase with severe depression in yield due to ultimately shorter reproductive phase.

- Plant Nutrient Management: About 5 tonnes FYM or compost or biogas spent slurry with 50% recommended dose of fertilizers (RDF) plus rhizobium inoculation for better yields and FUE. Recommended fertilizer dose is 15-20 kg N and 40 kg P₂O₅ per ha as basal dressing in separate furrow bands before sowing chickpea. Application of fertilizer is based on soil testing. In late sown chickpea after rice, apply 40 kg N per ha as basal dose. On S deficient soils, use 20 kg S as gypsum, iron pyrites or single super phosphate to meet the S demands of chickpea. Application of 25 kg zinc sulphate and 10 kg borax per ha has positive effect on root growth, BNF and yield.
- Weed management: Major weeds infestation in gram are *Chenopodium spp*. (Bathua), Fumaria parviflora (gajri), Lathyrus aphaca (Chatri matri), Vicia sativa (ankari), Crisium arvense (Kateli), Melilotus alba (senji), Asphodelus enuifolius (jungli piaji), Convolvulus arvensis (Krishan neel), Phalaris minor and Avena ludoviciana.

Being a dwarf stature crop, gram suffers adversely by heavy weed infestation up to 30-45 days after sowing (DAS), the critical period. One hand weeding/inter culture with hand hoe or wheel hoe at 30 DAS and another at 55-60 DAS, if second flush of weeds appear heavily other-wise crop will suppress the weed by it self. A mechanical operation is always better than the herbicide based as later also provides aeration to the roots for maximum efficacy of $\pm N\emptyset$ fixing bacteria as well as soil moisture conservation for its longer availability by breaking soil capillaries and creating dust mulch.

However, an alternate Integrated weed management practice is application of either of Fluchoralin (Basalin) as pre plant incorporation or Pendimethalin (Stomp) as Pre emergence @ 0.75-1.0 kg *a.i.*/ha and one hand weeding in between 30-45 DAS. Also application of Oxyflourfen 100-125 g a.i. /ha or 400 to 500 g or ml /ha at 0-3 DAS controls wide spectrum of weeds in the crop.

- Harvesting and threshing: Crop become ready for harvest when leaves begin to fall, stem and pod turn brown or straw in colour and seeds are hard and rattle (most important) with 15% moisture inside them. Over ripening may lead to fall of pods as well as shattering and seed cracking if seed moisture falls below 10% due to delay in harvesting. The crop is allowed to dry for 2-4 days on threshing floor (depending on situation) and threshed by manually or bullock/power drawn thresher followed by winnowing. The clean seed should be sun dried for 3-4 days to bring their moisture content at 9-10%. Now they should be safely stored in appropriate bins and fumigated to protect them from bruchids.
- Yield: By adopting good management practices, an average yield of 15-20 Q/ha can easily be obtained.

3.6.4 Plant Protection Measures

A. Disease


The important disease of Chickpea are Collar rot, Sclerotinia stemrot, Botrytis Grey Mold, wilt, Dry root rot. Symptoms of these disease and their suitable control measures are given below:

i) Collor Rot

Symptoms: The collar region of plant is constricted and begins to rot. White mycelia stands with minut mustard seed-sized sclerotial bodies are seen over the affected tissue. The affected seedlings turn yellow and wilt. It may be seen in seedling & vegetative growth stage.

Control Measures

i) Application of calcium fertilizer; ii) Seed treatment with fungicide carboxin @ 3 g /kg of seed; iii) Crop rotations with cereals such as wheat, sorghum and millets, and remove undecomposed debris from the field before sowing.



ii) Dry Root Rot

Symptoms: The whole plant dries up and turns straw-colored. Roots become black and brittle and have only a few lateral roots at all. It may be seen in flowering & podding stage.

Control Measures

i) Seed treatment with *Tricoderma viride* @ 4g /kg seed or Thiram (2g) + Carbandizm (1g) @ 3 g per kg of seed or Carbendazim @g/kg of seed; ii) Follow crop rotation; iii) Timely sowing to avoid post-flowering drought and heat stresses, which aggravate the disease.



iii) Wilt

Symptoms: The main cause of this disease is a fungus (*Fusarium* oxysporum). Plant become yellowish and finally dry out. Roots turn black and ultimately decompose. It may be seen in seedling stage & advance stage of plant growth.



i) Seed treatment with Tricoderma viride @ 4g /kg seed or Thiram (2g + Carbandizm (1g) @ 3 g per kg of seed or Carbendazim @ 2 g/kg of seed; ii) Sowing should be during third week of October; iii) Deep Planting (8-10 cm) in light soil; iv) In case of heavy incidence avoid cultivation for 03 to




04 years; v) Grow resistant varieties: *Desi-* JG 315, JG 322, JG 16, JG 11, JG 12, JSC 37, JSC 55, JAKI 9218; *Kabuli-* JGK 1, JGK 2, JGK 3 (Gulabi)- JGG 1.

iv) Sclerotinia Stem Rot

Symptoms: Ito initial stage is visible on the stem near the ground. Brown colour spots may be seen on affected stem which later girdle it White cottony growth of the fungus with hard, black colored sclerotia may be seen on these spots on the stem.

Control Measures

i) Use of disease free seed; ii) Grow resistant varieties like *G-543*, *Gaurav*, *Pusa-261*; iii) Deep summer ploughing & avoid flood irrigation iv) Spraying chlorothalonil @ 2 g/liter of water; v) After harvest, the diseased plants should not be allowed to stand in the field but should be destroyed by burning.



v) Botrytis Grey Mold

Symptoms: Brown necrotic spots appear on twigs, petioles, leaves and flowers. The affected stem finally breaks and the plant dies. In humid weather, the fungus grows rapidly.

Control Measures

i) Use disease free seed; ii) Seed treatment; iii) Grow resistant varieties like Pusa-1003, K-551, BG 276, GL 92162, IPCK 2004-29; iv) Adopt late sowing (first fortnight of November) and wider row spacing; v) Spray the crop with carbendazim (Bavistin) @ 1.5 g/liter or mancozeb @ 3g/liter of water.



B.INSECT-PEST

i) Cutworm

Nature of Damage: Serious pest in low lying areas where fields are cloddy. The larvae hidden under these clods during the day time & cause damage during night. It may be seen in Seedling, vegetative growth stage & reproductive stage. The caterpillar cut the plants at ground level. Larvae feed on leaves, stems and roots.

Control Measures

i) Summer deep ploughing; ii) Crop rotation; iii) Intercropping with wheat or linseed or mustard; iv) Grow marigold on bunds; v) Apply phorate 10 G @ 10 kg/ ha before sowing; vi) Spray insecticides like quinalphos 25 EC @ 2 ml /liter or Profenophos 50 EC @ 2 ml /liter.



ii) Gram Pod Borer

Nature of Damage: It damages almost all the pods in and causes nearly 20-30% annual yield losses in India. The eggs (1 mm diameter) are laid singly on the leaflets, flowers, immature pods and stem. Larvae can be green, brown, yellow, or pink, but are usually striped, irrespective of their colour. Larvae feed on leaves during the vegetative phase and on flowers and pods during the reproductive phase; ii) Large larvae cut round holes in the pod walland devour the seed inside.



Control Measures

i) Early sowing, grow short duration varieties; ii) Intercropping with coriander, linseed, marigold, mustard, sunflower or wheat; iii) Use moderate resistant cultivars like ICCV10, Vijay, ICCV-7 and ICCL-86103, PBG-3; iv) Install bird perches@ 40-50 /ha; v) Sprayneem seed extract (5%);vi) Apply HaNPV@250 LE/ha or Sprayindoxacarb@1 ml/lit or Emamectin benzoate 5 SG @ 0.2 g /lit of water at10-15 days interval if needed.

3.7 Recommendation to achieve higher production

To bridge the yield gap of 1014 kg/ha between the potential and realized yield following technologies are recommended:

- Deep summer ploughing once in 3 years to eliminate dormant pupae.
- Application of fertilizer based on soil test value.
- Seed treatment with Trichoderma (6 g/kg) and Carboxin (Vitavax) (1g/kg).
- Grow wilt resistant/ tolerant varieties of the region: JG 315, JG 12, JG 11, JAKI 9218, JGK-1, JGK-2, JGK-3, KAK2 etc.
- Install bird perches @ 50/ha at flowering stage and remove the perches at grain ripening stage.
- Nipping should be done when crop is at 15-20 cm height.
- Two irrigations first at branching and second at pod initiation stage.
- Weed control should be done at right time.
- Seed treatment with Ammonium Molybdate @ 1g/kg of seed in the areas of chickpeasoybean cropping system.
- Spray of crude NSKE 5% or Azadirachtin 0.03% (300 ppm), Neem oil based WSP 2500-5000 ml/ha at pre-flowering stage at 15 days interval.

.

(Table-3.9): Recommended Gram varieties & their characteristics

Variety	Source	Release/	Area of adoption	Ave. yield	Days to	Special characteristics
		Notification Year	(Zone/State)	(q/ha)	maturity	
RSG-44	RAU, Durgapura	1991	Rajasthan	20-23	135-150	Tol. to drought and frost, double podded
KPG-59 (Uday)	CASUAT	1992	NWPZ (Punjab, Haryana, Delhi, North Rajasthan & West U.P., Bihar & West Bengal).	20.0	135-140	Tolerant to root rot & wilt stunt. Tolerant to pod borer. Bold seeded.(late sown)
Bharati (ICCV-10)	ICRISAT	1992	SZ (A.P., Karnataka, Odisha & Tamilnadu) CZ (MP, Maharashtra, Gujarat).	18-20	95/-100	Resistant to <i>Fusarium</i> wilt & dry root rot.
Sadabahar	CSAUAT	1992	Uttar Pradesh	21-23	145-150	Tolerant to wilt.
Pusa-372 (BG-372)	IARI	1993	NEPZ (East UP, Bihar, WB). NWPZ (Punjab, Haryana, Delhi, Rajasthan CZ (MS & Gujarat). (late sown)	21-23 14.0 14-15	135-150	Moderately resistant to wilt, blight & root rot., Small seed, light brown
Sweta (ICCV-2)	ICRISAT	1993	Maharashtra, A.P	12-13	80-90	Kabuli gram variety . Resistant to wilt & Botrytis grey mould.
Pusa 329	IARI	1993	NWPZ (Punjab, Haryana, Delhi, North Rajasthan & West U.P).	21-23	145-155	Moderately resistant to Wilt, bold seeded
Vijay (Phule G-81-1-1)	MPKV	1994	CZ (MP, Maharashtra, Gujarat).	19-21	105-110	Resistant to wilt, Tolerant to terminal moisture stress.
Pragati (K-3256)	CSAUAT	1994	Uttar Pradesh.	17-20	140-150	Tolerant to wilt.
Vardan (GNG- 663)	RAU, Sriganga- nagar	1995	NWPZ (Punjab, Haryana, Delhi, North Rajasthan & West U.P).	22-25	150-155	Resistant to wilt.
GPF-2 (GF-89-36)	PAU	1995	NWPZ (Punjab, Haryana, Delhi, North Rajasthan & West U.P).	21-23	152	Resistant to wilt & tolerant to Ascochyta blight. Seed yellowish brown
Pusa-362	IARI	1995	NWPZ (Punjab, Haryana, Delhi,	23-24	145-150	Tolerant to wilt, Bold seeded.

Variety	Source	Release/	Area of adoption	Ave. yield	Days to	Special characteristics
		Notification Year	(Zone/State)	(q/ha)	maturity	
(BG-362)			North Rajasthan & West U.P).			
KWR-108	CSAUAT	1996	NEPZ (East Uttar Pradesh,	20-23	130-135	Resistant to wilt, Seeds are dark
			Bihar, West Bengal).			brown and small.
JG-218	JNKVV	1996	Madhya Pradesh.	18-19	115-120	Early maturing, Tolerant to wilt.
Vishal	MPKV	1996	CZ (MP, Maharashtra, Gujarat).	20.00	110-115	Resistant to wilt, Tolerant to pod
(Phule G-87207)						borer, Early maturing.
Alok (KGD-	CSAUAT	1996	NWPZ (Punjab, Haryana, Delhi,	19-21	140-150	Med. Bold, Res. to Wilt & Root rot
1168)			North Rajasthan & West U.P).			
Pant G-186	GBPUAT	1996	Uttar Pradesh	18-20	135-140	Tolerant to wilt & late sown. Small seeded
Hirwa Chaffa (AKGS-1)	PKV	1996	Maharashtra	15-17	105-110	Green seeded
Samrat (GNG-	RAU, Sriganga-	1997	NWPZ (Punjab, Haryana, Delhi,	20-22	145-150	Res.to Ascochyta blight. Tol. to wilt
469)	nagar		North Rajasthan & West U.P).			and root rot. Suitable for rainfed and irrigated areas.
Pusa-391 (BG-391)	IARI	1997	CZ (MP, Maharashtra, Gujarat).	17-18	110-120	Moderately resistant to wilt & root rot. Bold seeded. Light brown
PDG-3 (GF 89- 133)	PAU, Ludhiyana	1997	Punjab	15-17	160-165	Tolerant to pod borer.
Karnal Chana-1	CSSRI, Karnal	1997	NWPZ (Punjab, Haryana, Delhi,	22-25	140-147	Recommended for salt affected
(CSG 8962)			North Rajasthan & West U.P).			areas; Wilt resistant.
DCP-92-3	IIPR	1997	NWPZ (Punjab, Haryana, Delhi,	19-20	145-150	Lodging and wilt resistant.
			North Rajasthan & West U.P).			Yellowish brown and medium bold seeds. Suitable for high fertility and excessive moisture conditions.
JGG-1	JNKVV	1997	Madhya Pradesh	13-15	120-125	Seed pink
(BG-1003) (Pusa Kabuli)	IARI	1999	NEPZ (East Uttar Pradesh, Bihar, West Bengal).	17-19	140-150	White bold seeded, tolerant to wilt.

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
JG-11	ICRISAT/ PKV/JNKVV	1999	SZ (Odisha, Karnataka, A.P. & Tamilnadu)	15-17	95-100	Resistant to wilt, moderately resistant to root rot. Bold seeded
Guj. Gram-1	GAU	1999	CZ (MP,Maharashtra,Gujarat)	17-22	115-120	Wilt resistant, Dark brown, medium bold.
Dharwad Pragati (BGD-72)	IARI	1999	CZ (MP,Maharashtra,Gujarat)	25-30	115-120	Resistant to wilt & root rot, bold seeded
CO-3	TNAU	1999	Tamilnadu	9-11	80-85	Bold seeded, Resistant to wilt & Collar rot
CO-4	TNAU	1999	Tamilnadu	9-11	80-85	Bold seeded
JG-322	JNKVV	1999	Madhya Pradesh	18-20	110-115	Suitable for wilt prone areas.
WCG-2 (Surya)	Meerut Uni.	1999	Uttar Pradesh	20-25	135-150	Res.to rot, tol. to stunt & dry root rot
L-551(Kabuli)	PAU	1999	Punjab	18-20	135-140	Wilt tolerant.
Gujarat Gram 2 (GCP-107)	GAU	1999	Gujarat	22-24	95-100	Tolerant to wilt and bold seeded
Pusa Chamatkar (G 1053) kabuli	IARI	1999	NWPZ (Punjab, Haryana, Delhi, North Rajasthan & West U.P)	17-19	140-150	Tolerant to wilt
Gujarat Gram-4 (GCP-105)	GAU	2000	NEPZ (East U.P., Bihar, West Bengal).	18-20	135-130	Resistant to wilt. Seeds are dark brown.
PKV Kabuli-2 (KAK 2)	PKV	2000	CZ (MP, CG, MS, Gujarat)	17-18	125-130	Bold seeded
SAKI-9516 (Jawahar G-16)	JNKVV	2001	CZ (MP, Maharashtra, Gujarat)	18-20	110-120	Resistant to wilt.
Vaibhav (RG 2918)	IGKV	2001	Chhattisgarh	14-15	110-115	Seeds wrinkled and bold
Kranti (ICCC-37)	ICRISAT	2001	Andhra Pradesh	16-20	90-100	Resistance to Wilt & Dry root rot
WCG-10 (Pant G-10)	GBPUAT	2001	Maharashtra, Haryana, U.P.	21-25	147	Resistant to root rot, Mod. Resis. To stunt virus, wilt and dry root rot

Variety	Source	Release/	Area of adoption	Ave. yield		Special characteristics
		Notification Year	(Zone/State)	(q/ha)	maturity	
Haryana Kabuli 1 (HK- 89-131)	CSSHAU	2002	Haryana	20	142	Resistant to wilt
Virat (Kabuli)	MPKV	2002	Maharashtra	20	108-118	Resistant to wilt
JG-130	JNKVV	2002	Madhya Pradesh	15-16	110-115	Bold, Res.to wilt.
Jawahar Gram- 1(JGK 1)	JNKVV	2002	CZ - M.P., Maharashtra, Gujarat, Bundel khand region of U.P.	15-18	110-115	Mod. Resistant to wilt
Vihar(Phule G- 95311)	MPKV	2002	Karnataka, A.P., TN, Odisha	16-18	90-100	Seed Bold, Resistant to wilt
Anubhav (RSG 888)	RAU	2003	Punjab, Haryana, Delhi, North Rajasthan & WestU.P).	20-22	130-135	For rainfed, Moderately resistant to wilt & root rot
Pusa 1088	IARI	2003	Delhi	25-30	Med. early	Res. to wilt and root rots diseases.
Pusa 1103	IARI	2004	Delhi	19-23	Early	Resistant to root diseases.
Pusa 1105	IARI	2004	Delhi	25-30	Med.early	Mod. Resistant to root diseases.
Anuradha	Research station, Berhanpur	2004	West Bengal	22-25	120-130	Mod. Resistant to wilt.
Haryana Kabuli Chana 2 (HK 94 134)	CCS HAU	2004	U.P and Bihar	14	138	Resistant to wilt, Collor rot, dry root rot.
Asha (RSG 945)	ARS, Duragapura	2005	Rajasthan	17	75-80	Mod. Res. to dry root rot and wilt.
Digvijay	MPKV	2006	Maharashtra	19	105-110	Resistant to fusarium wilt
JG-63	JNKVV	2006	MP	20-25	110-120	Resistance to Wilt, Dry root rot & Mod. Resis. To Collor rot & <i>Helicoverpa</i> Species.
Akash (BDNG- 797)	MPKV	2007	Maharashtra	15-16	102	Resistant to wilt, tolerant to pod borer
Rajas (Phule-G-9425-9)	MPKV	2007	Punjab, Haryana, Uttarakhand, Delhi, Rajasthan and Jammu	18	136	Resistant to fusarium wilt
JGK-2	JNKVV	2007	M.P.	15	95-110	Resistant to collar rot, root rot, Mod. Res.to wilt and dry root rot

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
Lam shanaya(LBeG 7)	ANGRAU	2007	M.P.	20-25	90	Tolerant to Wilt and rot condition
JGK-3 (JGK 19)	JNKVV	2007	M.P.	14-15	92-121	Resistant to wilt
Jawahar Gram 226 (JG 226)	JNKVV	2007	M.P.		112-115	Resistant to wilt and root rot complex
GNG 421 (Gauri)	ARS, Sri Ganga Nagar	2007	Rajasthan	18	127-160	Tolerant to dry root rot, stunt and wilt
GNG 1488 (Sangam)	ARS, Sri Ganga Nagar	2007	Rajasthan	18	99-157	Tol .to dry root rot and stunt
RSG- 991(Aparna)	ARS, Duragapura	2007	Rajasthan	12-15	130-135	Mod. Res. to dry root rot, wilt, collar rot
RSG-896 (Arpan)	ARS, Duragapura	2007	Rajasthan	12-15	130-135	Mod. Res. to dry root rot, wilt, pod borer
RSG-902 (Aruna)	ARS, Duragapura	2007	Rajasthan	15-20	130-135	Mod. Res. to dry root rot, wilt, pod borer
JAKI 9218	PDKV	2008	Maharashtra	18-20	93-125	Resistant to <i>fusarium</i> wilt, root rot and collar rot
GNG 1581 (Ganguar)	ARS, Srigan- ganagar	2008	NWPZ	24.00	127-177	Resistant to water logging condition
WCG 3 (vallabh colour chana)	SVBPUAT, Meerut	2008	Uttar Pradesh	19-21	175	Bold seeded, protein content 22.8 %.
JG 6	JNKVV	2008	M.P.	20.00	103-132	Resistant to fusarium wilt and moderate resistant to dry root, Tolerantto pod borer
Pusa 2024	IARI	2008	Delhi	25-28	145	Moderately resistant against soil borne diseases and drought
BGD 103	UAS	2009	Karnataka	11-13	95-100	Resistant to fusarium wilt
JG 14	JNKVV	2009	M.P.	20-25	113	Moderate resistant to wilt, dry root and pod borer

74 Pulses in India: Retrospect and Prospects -2018

Variety	Source	Release/	Area of adoption	Ave. yield		Special characteristics
		Notification Year	(Zone/State)	(q/ha)	maturity	
Shubra (IPCK 2004-29)	IIPR	2009	CZ	21.00	104-108	Moderate resistant to wilt, escape terminal moisture stress and heat
Ujjawal (IPCK 2004-29)	IIPR	2010	CZ	20.00	103-111	Moderate resistant to wilt and tolerant to BGM, escape terminal moisture stress and heat
Phule G 0517	MPKV	2010	M.S., M.P., Karnataka	18.00	105-110	Tolerant to fusarium wilt, 59.4g/100 seed weight
Pant Kabuli chana 1	GPBUAT, Pantnagar	2010	Uttarakhand	30.00	120-122	Resistant to Botrytis grey mould
PKV Kabuli 4	PDKV	2010	Maharashtra, Madhya Pradesh	15-16	100-110	Moderately resistant to fusarium wilt dry rot and Botrytis grey mould
Gujarat Junagarh Gram 3 (GJG 0207)	JAU, Junagarh	2010	Gujarat	15.00	98-100	Moderately resistant to wilt and tolerant to pod borer
GPF 2	PAU	2010	NWPZ	22.00	134-163	Plants grow erect with thick stem resistant in lodging
MNK 1	UAS, Raichur	2011	Karnataka, A.P., Odisha, and Tamil Nadu	13.00	95-110	Moderately resistant to wilt
RSG 974 (Abhilasha)	ARS,Duragapura	2010	Rajasthan		130-135	Moderately resistant to wilt, dry root rot BGM and sterility mosaic
Raj Vijay Kabuli gram 101 (JSC 42)	RVSKVV	2012	Madhya Pradesh	15-20	90-110	Resistant to fusarium wilt and moderate tolerant to pod borer
CSJK 6	Durgapura	2012	NHZ	11	188	Suitable for normal sown irrigated condition. It posses dull white seed colour with 32.8g/100seeds) It matures in 188 days.
Raj Vijay gram 201 (JSC 40)	RVSKVV	2012	Madhya Pradesh	20-25	95-113	Resistant to fusarium wilt
HK 4 (HK 05-	CCSHAU	2012	NEPZ	15.00	136	Resistant to wilt, bold seeded

Pulses in India: Retrospect and Prospects -2018

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
169)					-	
Raj Vijay Kabuli gram 202	RVSKVV	2012	CZ	18-20	105	Suitable for late sown condition in paddy/cotton/soyabean-chickpea cropping system. Mod, resistant against wilt and dry root rot and collar rot
PBG -5	PAU, Ludhiana	2012	Punjab	17.00	160-165	Resistant to ascochyta blight disease
Raj Vijay Kabuli gram 203 (RVG- 203)	RVSKVV	2012	CZ	19-20	100	Suitable for planting under irrigated and late sown conditions. It is mod, resistant against wilt and dry root rot.
PKV harita (AKG 9303-12)	PDKV	2012	Vidarbha region of Maharashtra	12-18	106-110	Bold seeded, tolerant to wilt and drought, useful for culinary purpose
GLK 26155 (L 555)	Ludhiana	2012	NWPZ	22.8	146	Suitable for timely planting under irrigated condition. It has light yellow or creamy seed colour with 27.8 g/ 100 seeds weight. It matures in 146 days.
Phule G 0027	Rahuri	2012	NHZ	14	196	Suitable for normal sown irrigated condition. It has attractive white seed colour.
GJG 0809	Junagarh	2013	NHZ	16.0	157	Irrigated, medium brown colour attractive seed (21.5 g/100 seed), mod. Resistant to wilt & stunt, root rot & tolerant to ascochyta blight.
GNG 1969	Sriganga nagar	2013	NWPZ	22.0	146	Irrigated, suitable for normal sown irrigated condition. It posses creamy beige seed colour with 26.2 g/100 seeds.

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
GNG 1958	Sriganga nagar	2013	NWPZ (Punjab, Haryana, Delhi, North Rajasthan, West U.P).	26.8	145	Irrigated, suitable for normal sown irrigated condition. It matures in 145 days. It has brown seed colour with 25.4 g average 100-seed wt.
GLK 28127	Ludhiana	2013	NWPZ	21.0	149	Irrigated, large seeds (36.0 g/100 seeds), light yellow or creamy colour with irregular owl head.
NBeG 3	Nandyal (SVRC)	2013	Andhra Pradesh	23	110	Large seeded desi variety, (24.0 g/100 seed), tolerant to drought with good rooting quality, tolerant to wilt.
Vallabh Kabuli Chana-1 (WCGK 2000-16)	Modipuram (CVRC)	2015	NWPZ	23	147	Large seeds (27.5 g/100 seeds), white colour, 77 Moderately resistant to Fusarium wilt.
Birsa Chana 3	BAO, Jharkhand (SVRC)	2015	Jharkhand	18-20	115-118	Desi type, resistant to lodging, shattering, wilt disease and tolerant to gram pod borer.
Bidisha (BG 1084) WBG 29	Berhampur, WB (SVRC)	2015	West Bengal	25	131	Medium maturing desi variety with small seeds (12-14 g/100 seeds)
GNG 2144	Sriganganagar (CVRC)	2016	NWPZ	22.8	133	Desi type chickpea with medium bold seeds (15.9g/100seeds). It has tolerance to fusarium wilt disease
NBeG 119	Nandyal (SVRC)	2016	SZ	18.8	90-95	Large seeded (38.8g/100seeds) kabuli variety.
JGK 5	Jabalpur	2016	MP	15-17	110-115	Extra large seeded (55-60 g/100 seeds) kabuli variety having

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
	(CVRC)					resistant to <i>Fusarium Wilt</i> and moderate resistance to root rot.
CSJ 515	Durgapura (CVRC)	2016	NWPZ	24	135	Small brown colour seeds (15.9g/100seeds), mod. Resistant to dry root rot, wilt & collar rot and tolerant to Ascochyta blight and BGM
BDNGK 798	ARS, Jalna	2016 (SVRC)	Maharashtra	16-18	120-135	Kabuli type with white colour grain (29 g/100 seed), Moderately resistant to wilt and stunt.
Gujarat Junagarh Gram 6 (GJG 6)	Junagarh	2016 (SVRC)	Gujarat	16-18	120-130	Tolerant to wilt and stunt, suitable for normal sowing
JG 36 (Jawahar Gram 36)	JNKVV, Jabalpur	2016 (SVRC)	MP	18-20	110-120	Semi spreading plant with dark brown seed. Tolerant to wilt.
GBM 2	UAS, Raichur	2016 (SVRC)	Karnataka	18-20	100-110	Tall and erect plant type, suitable for mechanical harvesting
GJG 0809	Junagadh	2017 (CVRC)	NHZ	16	157	Medium brown colour attractive seed (21.5g/100seeds), mod. Resistant to wilt & stunt, root rot & tolerant to Ascochyta blight.
GNG 2171	Sriganganagar	2017 (CVRC)	NWPZ	20.14	163	Yellow coloured seed of size 15.9 gm/100seeds; tolerant to fusarium wilt disease
Indira Chana 1	IGKVV, Raipur	2017 (SVRC)	Chhattisgarh	16	101-116	Erect plant type with more primary branches, resistant to wilt
Nandyal Gram 49 (NBeG 49)	ARS, Nandyal	2017 (SVRC)	Andhra Pradesh	14-20	90-105	Semi spreading plant type with medium height, tolerant to wilt

Variety	Source	Release/	Area of adoption	Ave. yield	Days to	Special characteristics
		Notification Year	(Zone/State)	(q/ha)	maturity	
Pant Gram-4	GBPAUT	2017	Uttarakhand	18-20	126-162	Semi erect plant type, tolerant to
(PG 065)	(Patnagar, UT)	(SVRC)				wilt, BGM and dry root rot
Pant Gram -3	GBPAUT	2017	Uttarakhand	18-24	139-152	Semi erect plant type, medium seed
(PG 043)	(Patnagar, UT)	(SVRC)				(24.4 g/100 seed) tolerant to wilt
						and BGM
Pant Kabuli	GBPAUT	2017	Uttarakhand	16-17	135-163	Semi erect plant type, kabuli variety
gram-2 (PG 071)	(Patnagar, UT)	(SVRC)				(31.4 g/100 seed) tolerant to wilt
						and BGM
Dheera	ARS, Nanadyal	2017	Andhra Pradesh	20-25	90-105	Semi erect plant type, kabuli variety
(NBeG 47)		(SVRC)				, Suitable for mechanical harvesting

NHZ-North Hilly Zone (H.P.,J.K & U.P.hills), CZ- Central Zone (MP.,Maharashtra, Chhattisgarh, Gujarat), SZ- South Zone (A.P., Karnataka, Tamil nadu, Odisha) NEPZ-North East plane Zone (East Uttar Pradesh, Bihar, Jharkhand, West Bengal).NWPZ- North West Plane Zone (Punjab, Haryana, Delhi, West UP & North Rajasthan), Res. = Resistant, Tol. = Tolerant, Mod. = Moderately, BGM- Botrytis grey mould.

SVRC- State Variety Release Committee, CVRC-Central Variety Release Committee CZ- Central Zone, NWPZ- North Western Plain Zone, NEPZ-North Eastern Plain Zone, SZ-South Zone, NHZ-Northern Hills Zone.

Source: - Project Coordinator's Report, Annual Group Meet on Chickpea - Aug. 2017, AICRP, ICAR, IIPR, Kanpur.

PIGEONPEA (ARHAR)



Chapter - 4

Pigeonpea

Botanical Name : Cajanus cajan (L.) Millsp.

Synonym : Red gram, Tur

Origin : Africa

Chromosomes : 2n = 22

4.1 Economic Importance

In India pigeonpea (Arhar), the IIndimportant pulse crop after chickpea, is largely cultivated under rainfed conditions (95%), remaining 5% is grown with critical irrigation support. India ranks Ist in area and production in the world with 80% and 67% of worldøs acreage and production respectively (*FAO Stats.*,2014). During XIIth Plan this crop occupied16.6% of the total pulse area contributing to 17% of the total pulse production.

It is mainly consumed in split form as 'dal' and is a preferential pulse for Indians. Seeds are also rich in iron, iodine, essential amino acids like lycine, tyrocene, cystine and arginine. The outer seed layer along with the kernel part provides a valuable feed/concentrate to milch cattle. The husk of pods and leaves constitute a valuable cattle fodder. The dry sticks of the plant are used for fuel, thatches, storage bins (baskets) making etc.

4.2 Nutritive Value

Protein	22.3 %	Calcium	73 mg/100 g
Fat	1.7 %	Phosphorus	304 mg/100 g
Minerals	3.5 %	Iron	5.8 mg/100 g
Dietary Fiber	1.5 %	Calorific value	335 Kcal/100 g
Carbohydrate	57.6 %	Moisture	13.4%

4.3 Agronomic Significance

It is favoured legume as component crop in mixed/intercropping, hedge crop on the rice bunds/ vegetable crops. *Pigeonpea is a choice crop for improving soil characteristics due to deep tap root system with valuable properties and bed of fallen leaves besides restorer of nitrogen to the tune of 6-69 kg N/ha in a long duration pigeonpea over a period of 40 weeks (Kumar Rao and Dart., 1987, ICRISAT).*

4.4 Growth Rate

From 1980-81 to 2009-10, the total acreage under pulses has almost slightly (\pm) being showed, however, the maximum growth rate in area and producton was recorded with 35% & 90% during 2016-17. The highest area (5.34 million ha) and production (4.87 million tonnes) was also recorded during the same period (Table 4.1).

(Table-4.1): Growth rate of pigeonpea

(A- Million ha, P- Million tones, Y-kg/ha, Growth Rate (GR)-%)

Year				onpea	ng/na, Growin	
	Area	GR	Production	GR	Yield	GR
1980-81	2.84		1.96		689	
1985-86	3.18	12.0	2.44	24.7	767	11.3
1990-91	3.59	12.8	2.42	-1.0	673	-12.3
1995-96	3.45	-4.1	2.31	-4.4	670	-0.4
1996-97	3.51	1.9	2.66	15.2	757	13.0
1997-98	3.36	-4.4	1.85	-30.5	551	-27.3
1998-99	3.44	2.4	2.71	46.4	787	43.0
1999-00	3.43	-0.4	2.69	-0.5	786	-0.2
2000-01	3.63	6.0	2.25	-16.6	618	-21.3
2001-02	3.33	-8.4	2.26	0.6	679	9.8
2002-03	3.36	0.9	2.19	-3.3	651	-4.2
2003-04	3.52	4.7	2.36	7.8	670	3.0
2004-05	3.52	0.1	2.35	-0.4	667	-0.5
2005-06	3.58	1.8	2.74	16.7	765	14.6
2006-07**	3.56	-0.5	2.31	-15.5	650	-15.0
2007-08	3.73	4.6	3.08	32.9	826	27.1
2008-09	3.38	-9.3	2.27	-26.3	671	-18.8
2009-10	3.47	2.6	2.46	8.8	711	6.0
2010-11	4.37	26.0	2.86	16.1	655	-7.9
2011-12	4.01	-8.2	2.65	-7.2	662	1.1
2012-13	3.89	-2.9	3.02	13.9	776	17.2
2013-14	3.90	0.3	3.17	5.0	813	4.7
2014-15	3.85	-1.3	2.81	-11.6	729	-10.4
2015-16	3.96	2.8	2.56	-8.8	646	-11.3
2016-17	5.34	34.8	4.87	90.2	912	41.1
2017-18*	4.43	-17.0	4.25	-12.7	960	5.2

Source: DES, Ministry of Agri. &FW (DAC&FW), Govt. of India; 2017-18*- IVth Adv. Est.

(Table-4.2): Per cent share to total pulses

(Area- lakh ha, Production-Lakh Tones, Yield-kg/ha)

Year		Tur		To	tal Pulses		% Sha	% Share to Total Pulses		
	A	P	Y	A	P	Y	A	P	ΥI	
1980-81	28.42	19.57	689	224.45	106.24	473	12.7	18.4	145	
1990-91	35.93	24.17	673	372.55	203.68	547	9.6	11.9	123	
1995-96	34.47	23.09	670	331.77	170.10	513	10.4	13.6	131	
2000-01	36.32	22.46	618	203.48	110.75	544	17.8	20.3	114	
2001-02	33.28	22.6	679	220.08	133.68	607	15.1	16.9	112	
2002-03	33.59	21.86	651	204.96	111.25	543	16.4	19.6	120	
2003-04	35.16	23.56	670	234.58	149.05	635	15.0	15.8	105	
2004-05	35.19	23.47	667	227.63	131.30	577	15.5	17.9	116	
2005-06	35.81	27.38	765	223.91	133.84	598	16.0	20.5	128	
2006-07	35.62	23.14	650	231.92	141.98	612	15.4	16.3	106	
2007-08	37.26	30.76	826	236.33	147.62	625	15.8	20.8	132	
2008-09	33.78	22.66	671	220.94	145.66	659	15.3	15.6	102	
2009-10	34.66	24.65	711	232.82	146.62	630	14.9	16.8	113	
2010-11	43.67	28.61	655	264.02	182.41	691	16.5	15.7	95	
2011-12	40.07	26.54	662	244.62	170.89	699	16.4	15.5	95	
2012-13	38.93	30.23	776	232.57	183.42	789	16.7	16.5	98	
2013-14	39.04	31.74	813	252.18	192.55	764	15.5	16.5	106	
2014-15	38.54	28.07	729	235.53	171.52	728	16.4	16.4	100	
2015-16	39.63	25.61	646	249.11	163.48	656	15.9	15.7	98	
2016-17	53.38	48.73	912	294.47	231.31	785	18.2	21.0	116	
2017-18*	44.31	42.54	960	299.93	252.35	841	14.8	16.9	114	

Source: DES, Ministry of Agri. &FW (DAC&FW), Govt. of India; 2017-18*- IVth Adv. Est.

4.5 Import/Export

- Import: The import of pulses in India during April, 2014 to March, 2015 was 5.75 lakh tonnes worth Rs.2635.85 crore against the value of Rs.17196.87 crore for total foodgrains, Rs.121319.02 crore fortotal agricultural imports and against Rs.2737086.58 crore for total National Import. The import during April, 2015 to March, 2016 was 4.63 lakh tonnes worth Rs.3318.23 crore against the import value of Rs. 26841.87 crore for total foodgrains, Rs.140288.69 crore for total agricultural import and Rs.2490298.08 crore for total National import respectively. The share of Pigeonpea import to Agricultural import was 2.17% and 2.37% respectively during April, 2014 to March, 2015 and April, 2015 to March, 2016.
- Export: The pulses export of the country during April, 2014 to March, 2015 was 0.012 lakh tonnes worth Rs.8.82 crore against the value of Rs. 59500.54 crore for total foodgrains, Rs.239681.04 crore for total agricultural exports and against Rs.1896445.47 crore for total National export. The export during April, 2015 to March, 2016 was 0.040 lakh tonnes worth Rs.52.55 crore against the export value of Rs. 42622.29 crore for total foodgrains, Rs.215395.68 crore for total agricultural export and Rs.1716378.05 crore for total National export respectively during this period. The share of Pigeonpea export to Agricultural export was 0.004% and 0.024% respectively during April, 2014 to March, 2015 and April, 2015 to March, 2016.

(Table-4.3): Importing & Exporting Countries

(Quantity-Thousand Tonnes)

S.No.	IN	IPORT			EXPORT	
	Country	Avg.*	% Share	Country	Avg.*	% Share
1	Myanmar	235.51	44.05	USA	0.90	25.50
2	Tanzania Rep	131.21	24.54	UK	0.62	17.54
3	Mozambique	94.70	17.71	U Arab EMTS	0.50	14.17
4	Malawi	38.39	7.18	Canada	0.41	11.64
5	Sudan	17.00	3.18	Singapore	0.21	5.98
6	Kenya	11.15	2.09	Mozambique	0.16	4.60
7	Uganda	4.38	0.82	Thailand	0.16	4.58
8	Nigeria	1.26	0.23	Brunei	0.11	3.20
9	Benin	0.22	0.04	Malaysia	0.09	2.67
10	Afghanistan Tis	0.21	0.04	Saudi Arab	0.07	1.86
11	Canada	0.18	0.03	Australia	0.06	1.76
12	Australia	0.12	0.02	Angola	0.04	1.26
13	Malaysia	0.10	0.02	Sri Lanka DSR	0.04	1.16
14	Sri Lanka Dsr	0.09	0.02	Kuwait	0.03	0.75
15	Ethiopia	0.06	0.01	Tanzania Rep	0.02	0.57
16	U Arab Emts	0.03	0.01	Korea RP	0.02	0.43
17	USA	0.03	0.005	New Zealand	0.01	0.33
	Total	534.62		Total	3.53	

Source: Ministry of Commerce and Industry; Aveg.*- 2012-13 to 2016-17.

4.6 Production Trends

4.6.1 Global Scenario

India rank first in area (79%) and production (67%) at Global level. Mynmar stands second position in area (8.70%) followed by Kenya (3.93%) respectively. Tanzania occupy forth position in area (3.56%) and 5th position in production (2.48%). In productivity, Saint Vincent & Grenadines ranked first with 7926 kg/ha followed by Trinidad & Tobago (4103 kg/ha) and Malawi (4100 kg/ha). While, India& productivity is only 587 kg/ha.

(Table-4.4): Global Ranking: Major Counties

(Area-Lakh ha, Production-Lakh tonnes, Yield-kg/ha)

Country	Area	%Cont.	Country	Prod.	%Cont.	Country	Yield
India	56.02	79.65	India	32.90	67.28	Saint Vincent &	7926
						Grenadines	
Myanmar	6.12	8.70	Myanmar	5.75	11.76	Trinidad &	4103
						Tobago	
Kenya	2.76	3.93	Malawi	3.35	6.85	Malawi	4100
Tanzania	2.51	3.56	Kenya	2.75	5.61	Bangladesh	2500
Haiti	1.11	1.57	Tanzania	2.48	5.07	Philippines	1664
Malawi	0.82	1.16	Haiti	0.90	1.84	Grenada	1451
World	70.33		World	48.90		World	695
						India	587

Source: FAO, Statistics 2014




Fig.-4.1: Global Area: Pigeonpea




Fig.-4.2: Global Production: Pigeonpea

4.6.2 National Scenario: Plan Periods

Plan-wise performance revelaed, the highest area and production increased during XIIth plan over the previous plan period. The details plan-wise performance is given below:

(Table-4.5): Plan-Wise National Scenario

(Area-Million ha, Production-Million Tonnes, Yield-kg/ha)

Plan	Area	%	Prod.	%	Productivity	%
		COPP		COPP		COPP
Sixth Plan (1980-85)	3.03		2.27		749	
Seventh Plan (1985-90)	3.35	10.61	2.49	9.83	744	-0.71
Eighth Plan (1992-97)	3.48	3.75	2.43	-2.59	698	-6.12
Ninth Plan (1997-02)	3.44	-1.13	2.35	-3.12	684	-2.01
Tenth Plan (2002-07)	3.51	2.04	2.39	1.56	681	-0.47
Eleventh Plan (2007-2012)	3.79	8.03	2.66	11.56	703	3.26
Twelfth Plan (2012-2017)	4.19	10.5	3.29	23.6	785	11.6

Source: DES, Twelfth Plan (2012-2017); % COPP is percentage Change over previous plan




Fig.-4.3: Pigeonpea: Plan-wise Scenario

4.6.3 States' Scenario: Plan Analysis (Xth -XIIth)

- *Tenth Plan* (2002-2007): The country¢s total area coverage and production of tur were 35.07 lakh hectares and 23.88 lakh tonnes respectively. The state-wise trend shows that Maharashtra ranked first both in respect of area and production (30.87% and 31.30%) followed by Karnataka (16.00% and 12.12%). The third place occupied by Andhra Pradesh in area (13.28%) and U.P. (15.15%) in production. The highest yield recorded by Bihar (1194kg/ha) followed by Haryana (1032 kg/ha) and U.P. (953 kg/ha). The lowest yield recorded in the state of A.P. (451 kg/ha) followed by Chhattisgarh (482 kg/ha) and Karnataka (515 kg/ha).
- Eleventh Plan (2007-2012): The country total area coverage and production of tur were 37.89 lakh hectares and 26.64 lakh tonnes respectively. The state-wise trend shows that Maharashtra ranked first both in respect of area and production (30.68% and 33.44%) followed by Karnataka (18.69% and 14.75%). The third place occupied by Andhra Pradesh in area (13.14%) and U.P. 10.97%) in production. The highest yield recorded by Bihar (1286 kg/ha) followed by Haryana (1080 kg/ha) and Gujrat (1008 kg/ha). The lowest yield recorded in the state of A.P. (450 kg/ha) followed by C.G. (491 kg/ha) and Karnataka (555 kg/ha).
- Twelfth Plan (2012-2017): The country total area coverage and production of tur were 41.90 lakh hectares and 32.88 lakh tonnes respectively. The state-wise trend shows that Maharashtra ranked first both in respect of area and production (29.61% and 29.07%). Madhya Pradesh stood second position in production (15.82%) followed by Karnataka (15.68%). The third place occupied by Madhya Pradesh (13.29%) in area. The highest yield recorded by Bihar (1682 kg/ha) followed by Gujrat (1118 kg/ha) and Haryana (1040 kg/ha) and the lowest yield observed in the state of A.P. (489 kg/ha) followed by C.G. (581 kg/ha) and Karnataka (632 kg/ha).

The overall trend of area, production and yield shown increasing trend during the last three Plan Period.

(Table-4.6): Plan-Wise StatesøScenario: Major States

(Area-Lakh ha, Production-Lakh tonnes, Yield- kg/ha)

State		X th Plan	% to AI	XI th Plan	% to	N-Lakn tonnes, 11 XII th Plan*	% to
					AI		AI
A.P	A	4.66	13.28	4.98	13.14	2.22	5.29
	P	2.1	8.78	2.24	8.39	1.09	3.30
	Y	451		450	639	489	
Bihar	A	0.36	1.04	0.28	0.73	0.22	0.52
	P	0.43	1.79	0.36	1.35	0.37	1.11
	Y	1194		1286		1682	
Chattisgarh	A	0.56	1.59	0.55	1.46	0.57	1.37
	P	0.27	1.13	0.27	1.03	0.33	1.01
	Y	482		491		581	
Gujarat	A	2.81	8.02	2.64	6.96	2.46	5.86
	P	2.38	9.95	2.66	9.97	2.75	8.35
	Y	847		1008		1118	
Haryana	A	0.31	0.87	0.25	0.67	0.10	0.24
	P	0.32	1.32	0.27	1.02	0.10	0.32
	Y	1032		1080		1040	
Jharkhand	A	0.78	2.21	1.00	2.64	2.04	4.87
	P	0.55	2.3	0.77	2.89	2.04	6.21
	Y	705		770		1000	
Karnataka	A	5.61	16	7.08	18.69	8.17	19.49
	P	2.89	12.12	3.93	14.75	5.16	15.68
	Y	515		555		632	
Madhya Prd.	A	3.18	9.07	4.06	10.72	5.57	13.29
	P	2.32	9.71	2.57	9.63	5.20	15.82
	Y	730		633		934	
Maharashtra	A	10.83	30.87	11.62	30.68	12.41	29.61
	P	7.47	31.3	8.91	33.44	9.56	29.07
	Y	690		767		770	
Odisha	A	1.29	3.69	1.37	3.62	1.38	3.30
	P	0.93	3.88	1.16	4.37	1.24	3.77
	Y	721		847		895	
Punjab	A	0.08	0.24	0.05	0.13	0.03	0.08
	P	0.07	0.31	0.04	0.17	0.03	0.09
	Y	875		800		912	
Rajasthan	A	0.18	0.52	0.19	0.51	0.15	0.36
	P	0.11	0.46	0.13	0.5	0.12	0.36
	Y	611		684		805	
Tamilnadu	A	0.39	1.12	0.32	0.83	0.57	1.37
	P	0.24	0.99	0.23	0.86	0.51	1.56
	Y	615		719		894	
Telangana	A			3.2		2.79	6.65
	P			1.21		1.53	4.66
	Y			378		550	
Uttar Pradesh	A	3.8	10.85	3.3	8.72	3.00	7.17
	P	3.62	15.15	2.92	10.97	2.63	8.00
	Y	953		885		876	
All India	A	35.07		37.89		41.90	
	P	23.88		26.64		32.88	
	Y	681		703		785	




Fig.-4.4: Pigeonpea Area: State® Contribution (XIIth Plan)




Fig.-4.5: Pigeonpea Production: State & Contribution (XIIth Plan)

4.6.4 District Scenario (2016-17) – Potential Districts

The intra-state analysis at the district level is presented in Table 6 4.7. Inter-district analysis across the country revealed that the highest area and production of pigeonpea is in Gulberga district of Karnataka which are 7.29 *per cent* and 11.12 *per cent* respectively of countryøs total tur area and production. District-wise area, production and yield of top ten districts of India in respect of production are presented below which together contribute to 30.17 *per cent* and 48.73 *per cent* of area and production of the country.

The yield of Potential districts recorded higher than the National average yield except Vijayapur district of Karnataka.

(Table-4.7): Top Potential Districts: Pigeonpea

{Area-Lakh ha, Production-Lakh Tonnes, Yield-kg/ha}

Name of		A	rea	Prod	luction	Yiel	d
districts	State	Area	% to India	Prod.	% to India	Yield	YI
Gulbarga	Karnataka	3.89	7.29	5.42	11.11	1466	161
Latur	MH	1.18	2.20	3.98	8.18	3384	371
Vijayapur	Karnataka	3.89	7.29	3.30	6.78	894	98
Yavatmal	MH	1.74	3.25	2.34	4.80	1348	148
Amravati	MH	1.34	2.52	1.55	3.19	1156	127
Nagpur	MH	0.58	1.09	1.46	3.00	2509	275
Akola	MH	0.60	1.12	1.33	2.72	2219	243
Narsinghpur	MP	0.66	1.24	1.29	2.64	1952	214
Osmanabad	MH	1.17	2.20	1.16	2.37	985	108
Bharuch	Gujarat	1.06	1.98	1.14	2.33	1077	118
Total above		16.10	30.17	22.97	47.13	1426	156
All India		53.38		48.73		913	

Source: State Department of Agriculture




Fig.-4.6: Pigeonpea: Potential District

(Table-4.8): State-wise yield gap and additional return

(Yield: kg/ha: Return: Rs./ha.)

State	Vi	eld (kg/	(ha)	Gap ove	or FD	Cana	vor	Yield	Addit	
State	1 10	eiu (Kg/	на)	Gap ove		Gap o		2016-		
						SA			retur	•
								17	bridgin	0.0
									gap (F	
	IP	FP	SAY	Actual	%	Actual	%		FP	SAY
AP	1953	1498	493	455	30	1460	296	380	25025	80300
Bihar	2082	1765	1669	317	18	413	25	1581	17435	22715
Gujarat	1442	1098	1092	344	31	350	32	1156	18902	19250
Haryana	1897	1530	1104	368	24	793	72	915	20222	43633
Jharkhand	1513	1192	987	322	27	526	53	1002	17692	28948
Karnataka	1017	812	557	205	25	460	83	749	11293	25318
MP	1336	1043	819	293	28	517	63	1133	16097	28417
Maharashtra	1189	930	692	259	28	497	72	1042	14263	27335
Nagaland	1191	702	891	489	70	300	34	0	26895	16482
Punjab	1542	1282	944	260	20	598	63	860	14300	32890
Odisha	1236	888	880	348	39	356	40	884	19140	19553
Rajasthan	750	632	703	118	19	47	7	1088	6490	2585
Telangana	1518	1112	457	407	37	1061	232	676	22367	58373
TN	965	808	940	157	19	25	3	655	8653	1393
Tripura	895	616	716	279	45	179	25	0	15345	9827
UP	1773	1339	867	434	32	906	104	1074	23852	49812
Average	1394	1078	863	316	29	530	61	943	17373	29177

Source-Annual Report- 2016-17, GoI, DPD, Bhopal (Ave. 2013-14 to 2015-16)

State Average Yield – E&S (Ave. 2011-12 to 2015-16)

IP: Improved Practise FP: Farmers Practise SAY: State Avergae Yield

4.7 Pigeonpea Agronomy

4.7.1 Economic Classification : Based on plant and pod character and maturity duration, Arhar belongs to two groups:

- *i)* Cajanus cajan var. bicolor-group includes late maturing types with tall and bushy character. Flowers are yellow with purple streaked at the end of the branch. The standard which is largest of five petals in the flower possesses red veins on the dorsal side. Pods are relatively longer dark in colour with 4-5 seeds inside.
- *ii)* Cajanus cajan var. flavus- group includes early maturing varieties with shorter bushy plant types having flowers at several points along the branches. Flowers are yellow and pods are plain, shorter with 2-3 seeds inside.
- *iii*) On the basis of crop duration, there are three distinct groups: (i) early duration 6 100 to 150 days, (ii) medium duration -150-180 days, and (iii) late duration 6 180-300 days.

4.7.2 Botanical Decription

The pigeonpea has its origin in India. The plant is an erect shrub with considerable variation in height from 1-4 meter, depending upon the variety, growing season and management practices. Branching mostly begins from 6th to 10th node *i.e.* from 15-25 cm above ground. Leaves are trifoliately compound with central leaflets, longer than laterals.

Inflorescence is axillary raceme often forming a terminal panicle, open in the evening and remain open for whole night and up to noon of the next day. Self-pollination is a general rulebefore opening the flowers. However, cross-pollination may also occur to some extent; Pod Length varies from 5-10 cm and width from 0.6 to 0.9 cm and colour variation from green to dark brown; Seeds are round or lens shaped with colour variationfrom dirty white to silver white, light brown to chestnut brown and dark mottled brown to pinkish black with yellow cotyledons; Root System consists of a well developed central tap root with numerous secondary and lateral branches bearing nodules. Usually tall and upright variety produce longer and more deeply penetrating roots whereas spreading type produce shallower, more spreading and denser roots.

4.7.3 Production Technology

- Soil & Field preparation: It is successfully grown in black cotton soils, well drained with a pH ranging from 7.0-8.5. Pigeonpea responds well to properly tilled and well drained seedbed. A deep ploughing with soil turning plough in fallow/waste lands, zero tillage sowing under intensive cropping system and Broad Bed Furrow/Ridge-furrow planting in low lying as well as intercropping areas is recommended. Raised Bed method of planting by dibbling at 2 inches depth with Row to Row distance 4 to 5 feet also 15 feet gap (2 pairs of Tur on bed) under intercropping of soybean under transplanting (Dharwad method/SPI), 5 X 3 and 3 X 1.5 feet spacing is recommended.
- Climate: The crop prefers warm tropical and subtropical climate with fairly moist and warm weather during vegetative period. During the flowering and ripening stages, it requires bright sunny weather for setting of fruits. Temperature requirement varies from 30-35°C for germination, 20-25°C for active growth, 15-18°C during flowering and pod setting and 35-40°C at maturity. The crop is highly susceptible to frost at the time of flowering. Cloudy weather and excessive rainfall at flowering time damage the crop to a great extent.
- Sowing time: Early arhar should be sown in first fortnight of June with pre-sowing irrigation, so that the succeeding crop can be sown with the least delay. Late sowing crop is more likely to be damaged by frost in northern parts of India. However, under rainfed conditions sowing may be done immediately after soil saturating rains (10-15 cm) have started. In *diara lands* which are prone to flood, sowing must be delayed by mid September.
- **Sowing Method**: Three systems of sowings are practiced for pigeon pea. The common is flat sowing, the other methods are broadbed-furrow (BBF) for extra-early group and ridge-and-furrow for the late maturity group.
 - **Bund Plantation of Pigeonpea:** To meet the pigeonpea *dal* requirement, instead of growing a sole/ mixed crop farmers opt for bund-cropping in rice growing areas in Bihar, Orissa, Chhatttisgadh etc. Such pigeonpea growing areas and mixed cropping areas are not

accounted in total cropped area under pigeonpea. Delayed cropping is required under aberrant rainfall condition. So, under compulsion farmers may select suitable varieties.

- Transplanting of Pigeonpea: Apprehension of crop failure due to delayed monsoon can be managed or avoided by transplanting of pigeonpea. Before the onset of monsoon seedling can be made ready, thus saving 30 to 35 days of seedling stage and crop duration. It gives higher yields with minimized plant population, which is also effective for agronomic management. However, watering need to be ensured at transplanting stage. This technology has a limitation that cannot be adopted at large scale due to lack of irrigation facilities.
- Seed Rate & Spacing: The seeding rate of pigeonpea depends on the desired plant density for a genotype (early, medium or late), cropping system (pure crop, mixed crop, or inter crop), germination rate of seed and mass of seed.

 Early Maturing Var.ó20-25 kg/ha (Row to Row-45-60 cm & Plant to Plant-10-15 cm).

 Medium/Late Maturing Var.- 15-20 kg/ha (Row to Row- 60-75 & Plant to Plant-15-20 cm).
- **Seed Treatment:** Fungicide: Thiram (2gm) + Carbendazim (1gm) or Thiram @ 3 gm or Tricoderma virdie 5-7g /kg of seed; Culture: Rhizobium and PSB culture 7-10 g /kg seed.
- Varieties: The variety may be selected as per the adaptability to the region, agro-eco sitiation, sowing time and the cropping system (Table 63.11).

Pigeonpea Varieties identified for delayed planting

Date of sowing	Promising, Genotypes	Zone
Normal 20 July, Late 9 Aug., (20 days)	JKM 189, Asha, BSMR 853	CZ
Normal July 5, Late Aug., 8 (30-35 days)	TTB 7, LRG 30, LRG 38, LRG 41,	SZ
	WRP 1	
Normal 25 July,Late 31Aug., (30-35 days)	Azad	NEPZ
Pre rabi/post rainy season	DA 11, NDA 2, Pusa 9, Bahar	NEPZ
From 1 st to 15 th September		

• Cropping system: The space between the rows could be profitably utilized by growing short duration crops such as urd, moong, cowpea, etc; *Important cropping systems* followed are: MaizeóPigeonpea (Rabi); ii) Pigeonpea-Urd-Wheat; iii) Pigeonpea-Sugarcane; iv) Mung + Pigeonpea-Wheat; v) Pigeonpea (early)-Potato-urdbean.

State-Wise Prevalent Cropping System

Cropping System	States
Medium/late pigeonpea (sole cropping)	U.P., Bihar, Karnataka, M.P.
Pigeonpea ó wheat	Haryana, Punjab, U.P, north
	Rajasthan
Early pigeonpea + groundnut	A.P., Gujarat, T.N.
Late/medium pigeonpea +sorghum/soybean	M.P., Western U.P., Maharastra, T.N
Cotton+ Pigeonpea	Maharashtra, A.P., Gujarat, TN
Maize- rabi pigeonpea	UP., north Bihar, W.B., Assam

- **Intercropping:** Pigeonpea is commonly intercropped with a wide range of crops. In India, it was estimated that 80 ó 90 % of the pigeonpea were intercropped:
 - a) With cereals (sorghum, maize, pearl millet, finger millet and rain-fed rice)
 - b) With legumes (groundnut, cowpea, mung bean, black gram, soybean)
 - c) With long-season annuals (caster, cotton, sugarcane and cassava)

• State-wise recommended Inter-Cropping

States	Inter-cropping
Central & Southern States	Tur+Sorghum/Pearlmillet/G.Nut/Horsegram/ Cpowpea/
	Castor/Urd/Mung/Soybean
Bihar/ UP	Tur+Rice/Pearlmillet/Sorghum/Castor/Maize/Urd/Mung
	Tur + Groundnut
Jharkhand	Tur + G.Nut/Castor
Andhra Pradesh	Tur+ Horsegram/G.Nut/Cowpea/Millets
Karnataka	Tur + Cotton/Soybean/Sorghum/Maize
Gujarat, Telangana, MS	Tur + Maize/Bajra
M.P, A.P, and Gujarat	

• Zone wise varieties suitable for Inter Cropping

Pigeonpea Varieties	Associated Crop	Zone
Paras. AL 1455, Pusa 992	Urdbean, mungbean	NWPZ
Asha (ICPL 87119) JKM 7, GT 101,	Urdbean, mungbean, Soybean,	CZ
BSMR 853, JKM 189, BSMR 736	Groundnut maize, Sorghum	
TTB 7, JKM 186 CO 6, LRG 41,	Urdbean, mungbean, Groundnut,	SZ
WRG 27	Ragi	
NDA1, Pusa 9	Urdbean, mungbean, maize	NEPZ

- Water management: Being a deep rooted crop, it can tolerate drought. In crop planted in June, one or two pre-monsoon irrigations should be given as per requirement. After the start of monsoon, there is no need to irrigation but in case of prolonged drought during the reproductive period of growth, one or two irrigations may be needed. A pre-requisite for the success of arhar is proper drainage. *Ridge planting* is effective in areas where *sub-surface drainage is poor*. This provides enough aeration for the roots during the period of excess rainfall. During rainy season, water should not stand anywhere in the field.
- Plant Nutrient Management: Apply 25-30 kg N, 50-75 kg P₂O₅, 30 kg K₂O and 10-15 kg ZnSO₄ in one ha area as dose. Apply 20 kg S per ha in addition to NP at the time of sowing. For correcting Zn deficiency, foliar spray of 0.5 kg ZnSO₄ with 0.25 kg lime or soil application of ZnSO₄ @ 25 kg per ha to one crop on Zn deficient soils is helpful to both the crop of pulse based cropping system. Mo deficiency can be corrected by applying 1 kg sodium molybdate per ha and for boron deficient soils foliar spray of B @ 1.061.5 kg B per ha or soil application of 4 kg borax. Spray 1.0 per cent FeSO₄ to recoup the crop from Fe deficiency. Application of fertilizer is based on soil testing.

- Plant Nutrient Management in intercropping system: Application of full dose of nutrients to cereal component of pigeonpea intercrop ($N_{60}P_{40}$) along with full dose of fertilizers for pigeonpea ($N_{18}P_{40}$), has been found beneficial. In irrigated pigeonpea-cereal intercrop, the N should be split into two doses.
- Weed management: Weeds poses serious problem during rainy season by robbing the crop of precious nutrients and moisture and also give shelter to various insects and pests. The period of early 60 days is very critical for weed management point of view. Therefore, field should be kept free from weeds by giving two weeding through hand or wheel hoe at 25-30 and 45-50 days after sowing, respectively. If manual weeding is not possible either due to continuous rains or non availability of labour etc., weeds can also be manage successfully by using either of any one herbicides @ of 1 kg a.i./ha viz. Metachlor, Oxadiazon and Pendimithalin as pre-emergence spray or Basaline as pre-plant incorporation in soil.
- Weed management in intercropping system: An initial 45 and 30 days after sowing period is found very critical for severe weed crop competition causing a loss of about 46 % and 34% in NWPZ, 73% and 81% in CZ and 43 and 56% in NEPZ for pigeonpea intercropping with cereals and short duration pulses like green gram/black gram/cow pea/soybean, respectively.

Besides manual weeding with hand or wheel hoe, weeds may also be effectively controlled in pigeonpea intercropping system with pre-emergence application of Pendimethalin @ 0.5 \u00e1 1 kg a.i./ha depending upon weed intensity and soil type. Application of Quizalofop ethyl 100 g a.i./ha 15 to 20 DAS controls annual grasses and Imazethapyr 50-100 a.i./ha at 20-25 DAS for wide spectrum of weeds.

• Harvesting, Threshing & Storage: With two third to three fourth pods at maturity judged by changing their colour to brown is the best harvesting time. The plants are usually cut with a sickle within 75-25 cm above the ground.

Harvested plants should be left in the field for sun drying for 3-6 days depending on season. Threshing is done either by beating the pods with stick or using Pull man thresher. The proportion of seed to pods is generally 50-60%.

The clean seeds should be sun dried for 3-4 days to bring their moisture content at 9-10% to safely store in appropriate bins. To avoid further development of bruchids and other storage pests, it is recommended to fumigate the storage material before onset of monsoon and again after the monsoon with ALP @ 1-2 tablets per tonne. The small quantity of the produce can also be protected by mixing inert material (soft stone, lime, ash, etc) or by smearing edible/non-edible vegetable oils or by mixing plant products like neem leaf powder at the rate of 1-2% w/w basis.

• **Yield**: 15-30 qtls of grain (depending upon maturity group of variety and climate) and 50-60 qtls of sticks for fuel, as well.

4.7.4 Plant Protection Measures

A. Disease

The important diseases of Pigeon pea are Wilt, Sterility mosaic disease, Phytophthora blight, Alternaria blight, Powdery mildew.Symptoms of these disease and their suitable control measures are given below:

i) Wilt

Symptoms: Xylem gradually develops black streaks, dark purple bands appear on thestem surface plants extending upwards from the base. Main stem of such plants is split open, intensive blackening of the xylem can be seen. In humid weather, a pinkish mycelial growth is commonly observed the basal portions of the wilted plants. It may be seenin seedling, flowering & vegetative stage.

Control Measures

i) Seed Treatment with Trichoderma viride @ 10 g/kg of seed or Thirum (2 gm) + Carbendazim (1gm)/kg of seed; ii)

Soil application-T. virideó2.5 kg/ha + 50 kg of well decomposed FYM or sand at 30 days after sowing; iii) Mixed cropping with sorghum; iv) Uproot wilted plants; v) Avoid over or under watering plants; vi) Grow resistant varieties like Amar, Azad, Asha (IPCL-87119), Maruthi, C-11, BDN-1, BDN-2, NP-5, JKM-189, C-11, JKM-7, BSMR-853 & BSMR-736 etc.

ii) Sterility mosaic disease

Symptoms: It is caused by mosaic virus & spread from plant to plant under field conditions through *Eriophyid* mite. Leaves become small and cluster near branch tips & reduced in size. Plants are pale green and bushy in appearance, without of flowers and pods. Diseased plants are usually in groups. It may be seenin Vegetative growth & Pre-flowers stage.



Control Measures

Spray Fenazaquin 10 EC (Magister) @ 1 ml/liter of water on 45 and 60 DAS; ii) Rogue out the infected plants in the early stages of growth; iii) Crop rotation with non host crop like,tobacco, sorghum, pearl millet, cotton; iv) Grow resistant varieties like Pusa-885, Asha, Sharad (DA11),Narendra Arhar1, Bahar, BSMR-

853, BSMR 736, Rajeev Lochan, BDN-708.

iii) Phytophthora blight

Symptoms: Foliage blight symptoms are circular or irregular water soaked lesions on leaves. The lesions on stems and branches increases rapidly, girdles, cracks and dries the stem. Infected stem and branches break easily in the wind.

Control Measures

i) Seed treated with Metalaxyl 35 WS @3 g/kg of seed; ii) Good drainage in the fields and the plants should be protected from stem injury; iii) Crop rotation should be followed; iv) Grow resistant varieties like ICPL 7916/12055/12114/12161, JKM-189, JA-4 etc.



iv) Alternaria blight

Symptoms: Symptoms appear on all aerial part of plants are small, circular, necrotic spots that develop quickly, forming typical concentric rings. The spots are initially light brown and later turn dark brown. In severe infection, defoliation and drying of infected leaves, branches and flower buds.

Control Measures

Spray the crop with Mancozeb 75 WP @ 2 g/liter or Carbendazim 50 WP @ 1g/liter of water; ii) Cultivation of pigeonpea on ridges with proper drainage system and



avoiding the sowing in heavy soil helpful in disease management; iii) Grow resistant varieties like DA- 2,MA 128-1, MA 128-2.

B. Insect-Pest

i) Pod borers

Nature of Damage: It is widely distributed and is the most injurious pest of early and medium maturing varieties. The larvae, after hatching, feed on tender leaves and twigs but a pod formation they puncture pods and feed on developing grains. It may be seen in vegetative & podding stage.

Control Measures

Use *H. armigera* pheromone trap @ 12/ha; ii) Spray the crop with Emamectin benzoate 5% SG @220 g/ha. Or Indoxacarb 15.8% SC@333 ml/ha; iii) The caterpillar should be picked by hand after shaking the plants and destroyed in the early stages of attack.




ii) Tur Pod fly

Nature of damage: Stripes can be seen on the surface of the affected grains, while the attacked pods are somewhat twisted or deformed. In case of severe damage, as many as 80 *per cent* pods and 60 *per cent* grains may be damaged.

Control Measures

Spraying Neem seed kernel extract (NSKE) 5 *per cent* at 50% flowering stage to manage the insectøs populations; ii) Pest can be controlled by spraying the crop with Lemda Cyhalothrin 5 EC 400-500 ml in 800-1000 litres of water per hectare.



iii) Plume Moth

Nature of damage: The larvae damaged seeds as well cause flowers, buds and pods to drop. The caterpillar is greenish-brown in color and fringed with short hairs and spines. It also enters into the pod and feeds on developing grains.

Control Measures

i) Apply the Neem oil 2%; ii) Spray the crop with Azadirachtin 0.03% WSP 2500-5000 g/ha or Emamectin benzoate 5% SG @ 220 g/ha or Indoxacarb 15.8% SC@ 333 ml/ha.




iv) Pod-sucking bugs

Nature of damage: Damaged seeds become shrivelled and develop dark patches. Shedding of green pods.

Control Measures

Soil application of carbofuran 3G @ 15 kg/ha at sowing; ii) Spray the crop with HaNPV 3 x1012 POB/ha in 0.1% teepol; ii) Immature bugs can be hand picked and destroyed; iii) The main natural enemies of bugs are egg parasitoids, ants and birds reported reduce feeding by green shield bugs; iv) Spraying with aromatic plants (e.g. gums, lantana, Neem-based pesticides).



4.8 Recommendation to achieved higher production

- Deep summer ploughing once in 3 years.
- Application of fertilizer should be based on soil test value.
- Seed treatment should be done before sowing.
- Use wilt and Sterility Mosaic disease (SMD) resistant /tolerant varieties BSMR 736, 853, 846, ICPL 96053, BDN 2010, ICPL 43, 44, IPA 203, 204, 234 and IPH 09-5 as per suitability of region. (IIPR AICRP Pigeonpea).
- Wilt resistant varieties VL Arhar -1, Vipula, JKM -189, G.T.-101, Pusa 991, Azad (K-91-25), BSMR-736, MA-6 etc.
- Use hybrid varieties PPH -4, ICPH-8, IPH 09-5, ICPH -2740 as per suitability of region.
- Weed control should be done at right time&Adopt integrated approach for plant protection.

4.9 Hybrid Pigeonpea

- In pulses, self-pollination is a rule and pigeonpea is no exception. However under field conditions, occurrence of hybrid plants in the populations is not uncommon; and it happens due to some degree of natural out-crossing that took place in the preceding generation.
- Consequently the crop is considered as partially cross pollinated. Traditionally high yielding purelines *vis-a-vis* single plant selections of different maturity groups have been developed through exploitation of pedigree selection.
- But in pigeonpea, genetics of yield and yield governing traits offer new niches for exploitations of hybrid-vigour. For commercial hybrid seed production in crop, there are two main prerequisites: (i) *efficient mass pollen transfer mechanism* (ii) *stable male-sterile source*.
- The hybrid pigeonpea R&D program is supported by Department of Agriculture, Cooperation & Farmer's Welfare, Govt. of India, under National Food Security Mission and ICRISAT's Hybrid Parents Research Consortium.
- A) GMS Based Hybrids: Natural cross-pollination in pigeonpea was witnessed as early as 1919 but could not be exploited in commercial hybrid seed breeding due to non-availability of suitable male sterile source. In the recent past the genetic male system in Pigeonpea has been identified and exploited for commercial hybrid by public sector. The GMS based hybrid developed are given below.

GMS BASED PIGEONPEA HYBRIDS

Hybrid	Source/Public sector Institution
ICPH-8	ICRISAT, Hyderabad
PPH 4	PAU, Ludhiana
COPH1	TNAU, Coimbatore
AKPH 4101	PDKV, Akola
AKPH 2012	PDKV, Akola

- The above hybrids, however, could not be popularized due to seed production constraints, besides problem related to seed purity concern and economic feasibility. As an ongoing programme on agriculture research and development, sincere efforts have been made in 1994, when the work of identification of CGMS system under ICAR and NATP programme has been initiated.
 - a) CGMS Based Hybrids: Stable Cytoplasmic Genetic Male sterile lines (õAö lines) along with their maintainer line (õBö lines) and appropriate fertility restorer lines, with better combining ability for yield, have already been developed (õRö lines). By exploiting A, B and R lines, biotic and abiotic stresses resistant hybrids with yield superiority over best check are being developed.

These hybrids have given better results in terms of yield and earlyness. It is beyond doubt that the area, production and productivity shall enhance by adoption of hybrids

based on CGMS systems. CGMS based Hybrid GTH-1 developed by SKAU, SK Nagar (Gujarat), has recently been released for cultivation.

CMS BASED PIGEONPEA HYBRIDS

Hybrid	Source/Public sector Institution
ICPH-2671	ICRISAT, Hyderabad
ICPH- 2740	ICRISAT, Hyderabad

b) Package and Practices of Hybrid Seed Production

The package manual being developed by Dr. A.N. Tikle, Senior Scientist, Pigeonpea breeder, RVSKVV, Gwalior on Hybrid Seed Production is reproduced below:

- Selection of field: Select field where pigeonpea was not grown in the previous season. Pigeonpea hybrid and parents seed production should be taken in medium to light soil with good drainage facility.
- **Isolation requirement:** Hybrid seed production field must be isolated by about 200 m from other pigeonpea fields.
- Sowing time: Onset of monsoon (IInd fortnight of June óIst week of July) or sowing time may be delayed by 10-15 days as late season planting showed relatively short plant height.

Seed rate

Female -6-8 kg/ha (seed drill) or 1 kg/ha (hand dibbling) for medium maturing hybrid Male -2-3 kg/ha (seed drill) or 1-1.5 kg/ha (hand dibbling) for medium maturing hybrid.

- Seed Treatment: Vitavax 2.5 g + Trichoderma 5 g/kg of seed.
- Plant spacing and depth: 90 cm between rows and 30 cm within rows, whereas in late sown condition the row to row distance can be reduced. Two seeds may be placed 4-5 cm deep in the soil on raised hill. Only one plant should be allowed to grow per hill which enables easy monitoring.
- Row ratio of A-line and R-line: Seed should be sown in the row ratio of 4:2 of A-line and R-line for hybrid seed production. It is advisable to sow 2-4 border rows of R-line in periphery of seed plot for abundance of pollen availability for enhancing pollen load. A-line should be demarketed by sowing of soybean within the rows of A-line.
 - Thinning and re-sowing: It should be completed within 15 days after germination.
 - Inter-culturing & weeding: Crop should be kept weed free by 2-4 inter-culture operation by Kolpa and hand weeding.
 - Fertilizer dose: 25:50:20: 20 NPK & ZnSO4 kg/ha basal dose.
 - Irrigation: If necessary, crop may be irrigated on critical stages of crop growths viz., seedling. However, irrigation at flowering stage increases enhances better pod setting and seed formation.

• Identification of pollen shedders in CMS (A-) line

Anthers of the male sterile plants are totally devoid of pollen grains. The sterility is confirmed at flowering i.e. anthesis. Anthers of A-line are white and translucent type, while B-line and anthers are full of pollen grains which can be seen by naked eye in the form of pollen mass. Pollen shedder plants in A-line may be identified at flower initiation stage and removed by uprooting the plants.

• Identification of Off-types:

- a) Pollen shedder plants may be identified as flowers with full of pollen grain;
- b) A & B plants are morphologically/phenotypically same except for male sterility;
- c) Diagnostic morphological traits are useful in identification of off type plants.
- Rouging: Offótype, rogue and pollen shedder plants are being rouged out regularly at
 different plant growth stages to maintain high genetic purity. Rouging in seed production
 plots should be carried out first at seedling stage, second at flowering stage, third at pod
 setting stage and fourth at physiological maturity stage.
- De-podding:De-podding in male plant rows ensure continuous pollen availability for subsequent flowering flush for pod setting in A-lines. At 75 % physiological maturity, pods can be harvested by picking or cutting pod bearing portion of the branches of A-line to ensure second flowering flush in ratio crop to give additional seed yield.
- Plant protection measures: First spray of insecticide should be done at flower initiation with any safer insecticide for pollinators viz., Profenofos 1.5 l/ha or Quinalphos 1l/ha and second after 10-15 days with Profenofos 1.5 l/ha, Neem kernel extract 5%. If infestation of pod borer persists, third and fourth spray may be done in the interval of 10-15 days. Sex pheromonetrap 25/ha may be installed across the field. Spraying/dusting of Malathion, chlorpyriphos, cypermethrin should be avoided.

Harvesting

- i. Pods of A-lines can be harvested at 75 % physiological maturity by picking or cutting pod bearing portion of the branches.
- ii. R-lines should be harvested before A-lines to avoid admixture in hybrid seed.
- iii. Matured pod in A-line should be harvested and threshing should be made separately.
- Grading &packing: Seed should be cleaned and graded as per seed certification standards.
 Seed samples should be taken for seed testing and grow out test (GOT) before packing and labeling.



CPA 2043 A



Certified seed production of hybrid seed in the Ratio of 4: 2 rows of female (A-lines & R-lines)



Male sterile flower devoid of pollen grains in anther of Field view of Hybrid seed production of ICPH 2671 A-line



4.9.1 Chronology of R&D work in development of hybrid pigeonpea

- In 1991, a milestone in the history of food legume breeding was achieved when the worldøs first pigeonpea hybrid, ICPH 8, was released.
- ➤ ICRISAT and ICAR jointly developed the hybrid using a genetic male-sterility (GMS) system, although high production costs prevented acceptance by seed producers.
- ➤ In 2005, another breakthrough was achieved when a cytoplasmic nuclear male-sterile (CMS) hybrid was developed by crossing a wild relative of pigeonpea (Cajanus cajanifolius) and a cultivar.
- The new hybrid technology is based on a three line system that includes A-line (malesterile); B-line (maintainer), and R-line (restorer).
- > Several experimental hybrids were evaluated at ICRISAT and various ICAR centers, which demonstrated 50-150% superiority in yield over popular varieties.
- ➤ In over 2000 on-farm trials conducted in five states of India the hybrids ICPH 2671 and ICPH 2740 respectively exhibited 47% and 42% yield advantage over the best local variety.
- > Seed production of hybrids, mediated by honey bees, is easy. Under congenial growing conditions, 700-1200 kg/ha of hybrid seed was produce.

4.9.2 Scope

The hybrids having more canopy than traditional varieties, consequently need less per hectare seed rate compensating higher cost of hybrid seed. The hybrids have tremendous scope of popularization in northern and central India and other parts growing wheat crop after harvest of hybrid pigeonpea. Similarly in central and southernpart, early and medium duration hybrids are likely to play important role in increasing the area under pigeonpea. The cost of cultivation would automatically reduced with the higher production and increase in cropping intensity due to hybrids.

4.10 Situations/Season of Cultivation

4.10.1 Rabi Pigeonpea

This is practiced in flood prone areas where fields get flooded or waterlogged during rainy season. The states of U.P. (eastern parts), Bihar, West Bengal, Odisha, Gujarat C.G and M.P. may exploit this potential with following practices/recommendations, for successful cultivation of rabi pigeonpea.

- The sowing must be done in II/III week of September. The crop can also be taken after harvest of early maize or paddy.
- Sowing at closer spacing (30 x 20 cm) having up to 2 lakh plants/ha.
- High seed rate of 40-50 kg/ha should be followed.
- Sowing depth should not exceed 5 cm. The seed should be treated with culture.
- Apply N₃₀P₅₀ (N₂₀P₅₀ basal and N₁₀ top dressing at 30 days after sowing) and also apply 20 kg ZnSO₄ and 10 kg sulphur if previous crop is not supplied with Zn and S. Fertilizer Application is based on soil testing.
- The crop should be irrigated thrice i.e. at branching (30 DAS), pre-flowering (70 DAS) and pod filling (110 DAS) stages.
- Heptachlor 6% @ 25 kg/ha should be mixed in soil at the time of last tillage operation before sowing.
- Spraying of Malathion 0.05% or carbaryl 0.1% at pod formation stage, controls pod borers.

4.10.2 Summer Pigeonpea

An alternate best way for increasing cropping intensity and timely wheat planting under *pigeonpea – wheat cropping system* of Northern India with approximately 2 lakh ha area, is summer sowing of pigeonpea alongwith summer moong. Under this situation, advanced sowing of pigeonpea may be done during mid-April keeping row-to-row spacing of 90 cm, intercropped with 3 rows of greengram at 20 cm row spacing. Greengram become ready for harvest by the end of June after two pickings. Immediately in the space vacated by green gram, inter planting of black gram can be done between pigeonpea rows. While blackgram will be ready for harvest by end of September, pigeonpea attain maturity to be harvested by the mid November. Early harvest of pigeonpea thus facilitates wheat sowing at optimum time to harness the best yield. Thus, summer sown pigeon pea may be harvested alongwith other kharif crops in November and short statured crops of greengram and blackgram will be an additional source of income.

Summer pigeonpea is gaining popularity in Narmadapuram and Jabalpur Division (Hoshangabad and Narsinghpur).

4.10.3 Production Technology

Pigeonpea was usually considered a low-input marginal crop and therefore, not much attention was paid for development of improved production technology except for some nutritional requirement. Based on recent studies, production technology was strengthened which are suited to different zones as given below:

(Table.-4.9): Pigeonpea production technology for different zones.

Zone	Planting time	Seed rate & spacing	Seed inoculation	Fertilizer	Irrigation	Weed management	Cropping system
NWPZ: (Punjab, Haryana, Delhi, Western Uttar Pradesh and North West Rajasthan)	2 nd fortnight of May to1 st fortnight of June.	18-20 kg/ha 60 x 20 cm.	Rhizobium culture @ one packet for 10 kg seed.	N:P:K:S: ZnSO ₄ @ 20:40:20:20:25 kg/ha.	Pre sowing and in absence of rains at flowering and pod development stage.	Two hand weeding at 25 and 45 days after sowing (DAS) or application of Pendimethalin @ 1.25 kg a.i./ha followed by one hand weeding at 45-60 DAS.	Pigeonpea-wheat sequential cropping. Intercropping of mungbean /urdbean with pigeonpea (1:1 row ratio)
NEPZ: (Eastern part of Uttar Pradesh, Bihar, West Bengal, Assam & Jharkhand	Early:1 st fort night of June. Late: 1 st fortnight of July Pre-rabi: 1 st fortnight of Sept.	18-20 kg/ha 60 x 20 cm 12-15 kg/ha 75-90 x 25 cm. 25-30 kg/ha 30 x 10 cm.	-do-	-do-	Early: Pre sowing Pre-rabi: At 45 days after sowing (DAS) and at 100- 110 DAS in absence of rains.	-do-	Early: Pigeonpea-wheat sequential cropping. Late: Pigeonpea sorghum mungbean/urdbean/ Sesamum by pairing pigeonpea rows and planting one rows of intercrop.
CZ: (Gujarat Maharashtra , M.P., Rajasthan & Chattisgarh.	Both early and medium: 1 st fortnight of July/ on set of monsoon	Early:15-18 kg/ha 60 x 20 cm. Medium: 15- 16 Kg/ha 75-90 x 25 cm	-do-	-do-	In absence of rains at flowering and pod development stage.	-do-	Pigeonpea + Groundnut(2:4) Pigeonpea+Soybean (2:4), Pigeonpea + Sorghum (2:1) and pigeonpea+urdbean(1:1)
SZ: (A.P., Karnataka & Orissa)	Early: : 2 nd fortnight of June Medium: 1 st fortnight of July/ on set of monsoon	-do-	-do-	-do-	-do-	-do-	Pigeonpea+ mungbean/ urdbean /cowpea (1:1), Pigeonpea+ groundnut (4:2).

NWPZ= North West Plain Zone, NEPZ= North East Plain Zone, CZ= Central Zone, SZ= South Zone, DAS= Days after sowing.

(Table-4.10): Agronomic recommendations of ICAR-IIPR for increased production under different cropping systems in the country

Agronomic Practices	Impact
Plant Population: Optimum plant density for early maturing pigeonpea genotypes is 25-33 Plants/m². For Pre- <i>Rabi</i> planting 22 plants /m² and for long maturity group 11 plants/ m² are optimum. Cropping system: Pigeonpea + groundnut (first year) followed by Maize + Groundnut (2 nd year)	Optimum plant population significantly contributes towards the yield and critical solar energy harvest, its conversion and higher production per unit of land. Increased the productivity and has potential to provide increased gross return under sequential cropping system.
 Intercropping: Pigeonpea +Sorghum / Pearl millet / Seasmum (2:1 row ratio) and (2:2 row ratio) Pigeonpea + Mungbean /Urdbean /Soybean /Maize /Groundnut Pigeonpea + maize (2:1) supplemented with FYM @ 5 tonnes per hectare. NWPZ: Pigeonpea (Paras, Pusa 992, AL 1455) +Mungbean/Urdbean/soybean/Groundnut CZ: Pigeonpea (Asha, JKM7, JKM189, GT 101, BSMR 736, BSMR 853) + Mungbean/Urdbean/soybean/Groundnut. SZ: Pigeonpea (TTB 7, JKM 186, Co6, LRG 41) + Mungbean /urdbean/ Urdbean/ Soybean/ Groundnut/ fieldbean. 	 Bonus yield of intercrops, highly remunerative as compared to sole crop of pigeonpea both under optimal and sub-optimal fertility levels. Reduced wilt incidence (12-21%) than sole crop of pigeonpea. Reduced pod borer incidence 27% than sole pigeonpea damage 49% with the help of bird perches (Sorghum, maize) Higher pigeonpea equivalent yield (PEY) net return, return /Rs invested and LER. Bonus yield of intercrops Spray (2% urea) at 15 and 45 DAH of intercrop is beneficial. Reduced yield of pigeonpea but higher gross monitory return alongwith additional harvest of fodder sorghum
Ridge and Furrow raised beds planting: • Pigeonpea (ridges) + Rice (Furrow) • Pigeonpea (Ridges) + Intercrops(side of furrow) • Pigeonpea (Ridges) + Urdbean 2:2 (raised beds) Weed management:	Ensured optimum plant population and higher grain yield as a bonus without any adverse effect on grain yield on pigeonpea. Yield advantage (23%) over flat system and sufficient control to seasonal
 Raised bed system (2.7 m width) of planting and one hand weeding of 30 DAS. Pre-emergence application of pendimethalin (1.0-1.5 kg/ha) or Benthiocarb, Metalachlor /Alachlor @ 2.0 kg/ha and hand weeding 25-30 DAS optional second weeding (45-60 DAS) Pre-emergence spray of Pendimethalin @ 0.75 kg/ha followed by post emergence spray of Paraquot of weeds in Pigeonpea. 	weed.

Agronomic Practices	Impact
Integrated Nutrient management	Grain yield advantage as recorded upto 177 kg /ha.
 40 kg N, 22 kg P₂0₅/ha, 20 kg/ha Sulphur(Gypsum& SSP), Sodium molybdate @ 1.0kg/ha and ZnSO₄@ 15 kg/ha Pigeonpea requires fertilizer dose of 20:18:27:20:25 kg N:P:K:S:Zn Soil application of sodium molybdate @ 4g/kg seed, chelated iron @ 1 kg/ha and RDF + Seed treatment borax (4g/kg of seed) enhances the yield. Seed treatment with micronutrients such as Zinc, boron, molybdenum and iron has been found as effective as that of soil application. Application of borax, molybdenum and iron (deficient soil) through soil application, foliar application and seed treatment however, rate will vary depending on extent of deficiency in the soil. Soil amelioration through lime in low pH soil supplement with molybdenum. Application of Vermicompost (2.5t/ha) + organic fertilizer (75% RDF) + biofertilizer 	 Rabi pigeonpea (Sharad), yield increase 25% over control Yield advantage 18-20 kg/ha respectively with NPKS (20-18-27-20 kg/ha) as compared to without sulphur application. No response of Mb in Rabi Pigeonpea. Application of ZnSO₄ with RDF is beneficial.
(Rhizobium+ PSB+ PGPR) enhances the yield and soil physical properties.	
Agronomic Practices for insect Pest management:	• Recorded lower pod borer incidence 20%.
 Pigeonpea +Sorghum (1:2) + optimum dose of fertilizer application than sole pigeonpea with sub-optimum dose of fertilizer <50%. Pigeonpea + Sorghum (1:2)+ RDF+ chemical control (20%). 	• Lower wilt incidence (11%)
Cultural Practice:	• Efficient use of phosphorus through P ótolerant varieties for higher yield.
 In P- deficient soils some varieties viz. Such as GT-1, VRG 17, TTB 7 and VBN 2 were recorded as efficient user of Phosphorus. Under delayed planting upto first week of August- use of suitable genotypes such as Asha, JKM 189, BSMR 853(2), TTB 7, LRG 41 30, 38 WRP 1 (SZ) and Azad (NEPZ). 	
Drought Management:	• Drought management with physiological and agronomic practices.
 Spray of 2% KCl for minimizing the adverse effect of terminal drought. Foliar application of urea, KCl and seed soaking with CaCl2 and K₂HPO₄ in reducing the adverse effect of water stress at the terminal growth of pigeonpea. Cultural mulch and water spray to minimize the adverse effect of terminal drought. 	
 HybridPigeonpea: For hybrid GTH-1 plant density of 75 x 20-30 cm. And application of 20-60-25-20 and 15 Kg NPKS and Zn was optimum. 	• Improved technology through Agronomic management.

Accumulation of run off water in a pond can be successfully utilized for substances of crop during the water scarcity.

(Table-4.11): Recommended Varieties/Characteristics

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. Yield (q/ha)	Days to maturity	Special characteristics
BSMR-175	MAU	1991	Maharashtra	11-12	165-170	White seeded, res. To Sterility Mosaic & Wilt
JA-4	JNKVV	1991	Madhya Pradesh	16-18	180-200	Tolerant to wilt & SMD
Birsa Arhar 1	BAU	1992	Bihar	10-15	180-200	Wilt Resistant
Gujarat Tur-100	GAU	1992	Gujarat	16-18	120-135	Tolerant to wilt &SMD white, bold-seeded.
Vamban 1	TNAU	1993	Tamil Nadu	8-10	95-100	Suitable for inter cropping with Peanut
Asha (ICPL-87119)	ICRISAT	1993	CZ&SZ (M.P., MS, Gujarat, Karnataka, AP, Odisha, Tamilnadu).	16-18	160-170	Resistant to wilt & SMD, Bold seeded., Indeterminate
Pusa-855	Central	1993	NWPZ (Punjab, Haryana, Delhi, North Rajasthan, West U.P).	24-25	145-150	Plant Indeterminate, Medium bold seeded.
Pusa-9	IARI	1993	NEPZ (East Uttar Pradesh, Bihar, West Bengal).	22-26	210-248	Tolerant to Alternaria & SMD, Tall & bold- seeded, Suitable for pre-rabi.
CO-6	TNAU	1993	Tamil Nadu	8-10	170-180	Tolerant to Pod borer. Indeterminate
Sharad (DA 11)	RAU, Dholi	1993	Bihar	18-20	240-250	Alternaria blight & Sterility Mosaic Resistant
Sarita (ICPL 85010)	ICRISAT	1994	A.P.	10-12	130-140	Determinate.
TS-3	UAS, Gulberga	1995	Karnataka	14-16	180-190	White, bold seeded, res. To Wilt
AL-201	PAU	1995	Punjab.	15-16	140-150	Indeterminate variety.
Durga(ICPL84031)	ICRISAT	1995	Andhra Pradesh.	8-10	120-125	Determinate.
Jawahar (KM-7)	JNKVV	1996	CZ (MP,MS & Gujarat). SZ (Odisha, Karnataka, A.P. & Tamilnadu).	18-20	173-180	Tolerant to wilt & Phytophthora blight. Seeds dark brown
BSMR-736	MAU	1996	Maharashtra	12-18	180-185	Resistant to wilt and SMD. Brown seeded.

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. Yield (q/ha)	Days to maturity	Special characteristics
						Indeterminate
Narendra Tur-1	NDUAT	1997	Uttar Pradesh.	20-22	240-260	Resistant to SMD and tolerant to wilt and
(NDA-88-2)						Phythopthora blight.
Amar (KA 32-1)	CSAUAT	1997	Uttar Pradesh.	16-20	250-270	Compact, res. To SMD. Tol. To wilt, Seed brown
H 82-1(Paras)	CCSHAU	1998	Haryana	15-20	133-145	Indeterminate
Malviya Vikalp (MA-3)	BHU	1999	CZ (M.P., Maharashtra, Gujarat).	20-22	178-162	Spreading, Constricted Pod, resistant to pod fly.
Azad (K 91-25)	CSAUAT	1999	U.P. & Bihar	20-22	250-260	Wilt Tolerant, Sterility Mosaic Resistant
AKT-8811	Akola	2000	Maharashtra	13-14	145-150	Indeterminate
Laxmi (ICPL-85063)	ICRISAT	2000	Andhra Pradesh	18-20	160-200	Pre-rabi
Vaishali (BSMR-853)	MAU	2002	Maharashtra	16-17	165.170	Resistant to wilt and SMD.
Sel-31	ARS, Gulbarga	2002	Karnataka	12	100-110	Irrigated command areas wherever double & multiple cropping system is being in practices
Pusa-992	IARI	2002	Haryana, Punjab, U.P., Rajasthan	18-20	130-140	Indeterminate
MA-6	BHU	2002	Central & Eastern U.P.	20-23	248-267	Late, Spreading type
Pusa 991	IARI	2003	Delhi	16-20	140	Tolerant to wilt, Phytophthora blight and SMD
Pusa-992	IARI	2004	Haryana, Punjab, Delhi, Western UP and Rajasthan	17	119-162	Tolerant to SMD and wilt
GT-101	GAU	2004	Gujarat	13	Early	Tolerant to wilt and SMD
Malviya chamatkar (MAL-13)	BHU	2005	UP & W.B.	27-29	226-271 (Kh) 189- 248 (Preó rabi)	Moderately resistant to wilt and SMD, Recommended for Kharif and pre-Rabi seasons.
VL Arhar-1	VPKAS,	2006	Uttarakhand	19	150	Res.to wilt, Alternaria leaf blight and rot

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. Yield (q/ha)	Days to maturity	Special characteristics
	Almora					
CORG-9701	TNAU	2006	Tamil Nadu, Karnataka, A.P., Odisha	11	120-130	Tol. To wilt, Sterility Mosaic & phytophthora blight, Tol. To pod borer & pod fly
Amol (BDN 708)	ARS Badnapur	2007	Maharashtra	15	160-165	Moderate resistant to wilt & sterility mosaic, Tolerant to pod borer & pod fly
Vipula	MPKV	2007	Maharashtra	16	145-160	Resistant to <i>Fusarium wilt</i> , Moderate resistant to sterility mosaic disease
Lam-41	ANGRAU	2007	A.P.	12	Medium	Tolerant to <i>Helicoverpa</i> pod borer
Jawahar (JKM-189)	JNKVV	2007	M.P.	21	116-124	Res. To wilt, Moderately resistant to sterility mosaic and Phytopthora blight
GTH-1* (SKNPCH-10)	SDAU	2007	Gujarat	18	135-145	No incidence of sterility mosaic disease
TT-401	BARC	2007	M.P., MS Gujarat & CG	16	138-156	Tolerant to pod borer & tolerant to wilt
Pusa 2002	IARI	2008	Delhi	17	110-150	Sowing to first week of June, suited for double cropping system, tolerant to moisture stress
Pant Arhar 291 (PA 291)	GBPAUT	2008	Uttrakhand	17	140-150	Early maturing, tolerant to phytophthora blight and pod borer
PAU 881 (AL 1507)	PAU	2008	Pun., Har., Western UP and plains of Uttrakhand	16-18		Early maturing
NDA 2	NDUA&T	2008	U.P., Bihar, WB, Assam & Jharkhand	25-28	240-260	Long duration, resistant to SMD and pod borers
TJT 501	BARC & ZARS, Khargone	2009	CZ	18	135-183	Tol.to SMD, wilt and phytophthro. Tol. To pod borer and pod fly
Surya (MRG-1004)	ARS Madhira	2009	Andhra Pradesh	20-22	166-180	Tolerant to Macrophomina stem canker/wilt
BRG 2	UAS, Bangalore	2009	SZ	12-16	175-185	Moderately tolerant to wilt, SMD and pod borer

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. Yield (q/ha)	Days to maturity	Special characteristics
TS-3R	ARS, Gulbarga	2010	Karnataka	11-17	150-160	Kharif and late sown cropping system res. To wilt
PKV, Tara (TAT-9629)	PDKV, Akola	2011 (SVRC)	CZ	19-20	154-189	Indeterminate, Semi spreading and tolerant to Pod Borer & pod Fly
AGT 2	Anand Agri. Uni., Moel farm Vadodara	2011 (SVRC)	Gujrat	15-16	170-180	Indeterminate, semi spreading, white and bold seeded, medium maturity, yields about 16 q/ha
Rajeev Lochan	IGKVV, Raipur	2011 (SVRC)	Chattisgarh	18-19	178	Resistant to wilt & sterility mosaic disease
WRG-65	ARS,Waran gal	2012 (SVRC)	Andhra Pradesh	14-16	150-160	Indeterminate, spreading medium, Resistant to wilt, tolerant to pod borer
BDN 711	ARS,Badna pur	2012 (SVRC)	Maharastra	15-23	150-160	Indeterminate, spreading, Resistant to wilt & sterility mosaic disease, escape terminal drought
Phule T 0012	MPKV, Rahuri	2012 (CVRC)	CZ	18-20	135-150	Moderately resistant to fusarium wilt, SMD and tolerant to Pod borer and Pod fly
RGT-1	ARS Tandur	2012 (CVRC)	Telangana	14-15	150-160	High yielding, Wilt resistant, semi determinate, suitable to <i>kharif</i> and <i>rabi</i>
Anand grain Tur 2 (AGT 2)	AAU	2012	Gujarat	16	175-180	No severe disease was observed
BDN 711 (BDN 2004-3)	ARS Badanapur	2012	Maharashtra	15-23	150-160	Mod. Resistant to wilt and SMD
ICPH 2671	RAK College Sehore	2013 (SVRC)	Madhya Pradesh	15-16	180-190	Indeterminate, medium maturity, tolerant to wilt and sterility mosaic
BRG 4	UAS, Bangalore	2014 (SVRC)	Karnataka	12-16	135-150	Indeterminate, semi determinate, mid early, suitable for normal and delayed

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. Yield (q/ha)	Days to maturity	Special characteristics
						sowings
IPA 203	IIPR, Kanpur	2014 (CVRC)	NEPZ	16-18	115-120	Resistannt to sterility mosaic disease, tolerant to Phytohthora blight and Fusarium wilt.
GJP-1	JAU, Junagrh	2015 (SVRC)	Gujrat	20-21	177	Indeterminate, semi spreading, medium maturity, light colour foliage, white seeded, bold seed size, moderately resistant to wilt and sterility mosaic and resistant to blight
TDRG 4	ARS Tandur	2015 (SVRC)	Telangana	20-22	170-180	Indeterminate, spreading, resistant to wilt and moderately resistant to sterility mosaic disease and highly tolerant to pod borer
PRG 176	RARS, Palem	2015 (SVRC)	Telangana	20-22	130-135	Indetrminate, semi spreading, , suitable to low railfall areas and chalka soils of Telangana
ICPH 2740	ICRISAT	2015 (SVRC)	Telangana	15-16	180-185	Indeterminate, semi spreading, resistant to sterility mosaic and wilt.
BRG 5	UAS, Bangalore	2015 (SVRC)	Karnataka	10-12	160-170	Indeterminate, semi spreading, red flowers, brown seeded (12.5 g/100 seed), resistant to wilt and moderately resistant to sterility mosaic disease
LRG 52	RARS, Lam	2015 (SVRC)	Andhra Pradesh	15-18	170	Indeterminate, semi spreading, dark purple pods, brown and large seeded and moderately resistant wilt
Rajendra Arhar-1	Dr. RPCA, Pusa, Bihar	2015 (SVRC)	Bihar	28-30	260 - 370	Indeterminate, erect and compact, dark brown and oval seeds, resistant to sterility mosaic and moderately resistant to wilt.

Pulses in India: Retrospect and Prospects -2018

Variety	Source	Release/	Area of adoption	Ave. Yield	Days to	Special characteristics
		Notification Year	(Zone/State)	(q/ha)	maturity	
GT 103	SDAU SK	2015 (SVRC)	Gujrat	15-16	140-150	Indeterminate, spreading, cream colour
	Nagar					and boldseeds matures in about 175 days,
						resistant to SMD and tolerant to wilt.
GRG 881	ARS	2016 (SVRC)	Karnataka	13-16	160	Indeterminate, semi spreading, , brown
	Gulbarga					seeded, resistant to fusarium wilt
						moderately resistant to sterility mosaic
						disease, 100 seed weight 10 grams
BDN 716	ARS	2016 (SVRC)	Maharastra	21-22	160-165	Indeterminate, spreading, resistant to wilt
	Badnapur					and sterility mosaic, red and large seeded
						(12g/100 seed)
CORG 8	TNAU	2016 (SVRC)	Tamil Nadu	18-20	170-180	Indeterminate, bold seeded (100 seed wt.
	Coimbatore					11g), resistant to sterility mosaic disease
						and tolerant to Helicoverpa armigera &
						Maruca vitarata

^{*} Hybrid; Res. = Resistant, Tol. = Tolerant, Mod. = Moderately, SMD= Sterility Mosaic Disease SVRC- State Variety Release Committee, CVRC-Central Variety Release Committee CZ- Central Zone, NWPZ- North Western Plain Zone, NEPZ-North Eastern Plain Zone, SZ-South Zone, NHZ-Northern Hills Zone.

Source: - Project Coordinator's Report, Annual Group Meet on Pigeonpea, 2016-17, AICRP, ICAR, IIPR, Kanpur.

GREENGRAM (MUNGBEAN)





Chapter-5

Green Gram (Mungbean)

Botanical Name : *Vigna radiate* (L.)
Origin : India and Central Asia

Chromosome : 2n = 24Synonym : Mung

5.1 Economic Importance

Green gram is an excellent source of high quality protein with easy digestibility, consumed as whole grains, dal and sprouted in variety of ways. As value addition, splitand dehusked, fried in fat, fetch good value as snacks. After harvesting the pods, green plants are fed to the cattle. The husk of the seed also used as cattle feed. It is also used as green manuring crop.

In India these crops are cultivated in three different seasons, viz., kharif, rabi and summer. Summer moong can be grown after harvesting of pea, gram, potato, mustard, linseed. Cultivation of Jayad Moong is important to increase soil fertility in these areas where paddy ówheat crop rotation is used.

5.2 Nutritive Value

Protein	24-25%	Calcium	124 mg/100 g
Fat	1.3%	Phosphorus	326 mg/100 g
Minerals	3.5%	Iron	7.3 mg/100 g
Dietary Fiber	16.3%	Calorific value	334 Kcal/100g
Carbohydrate	56%	Moisture	10%

5.3 Agronomic Importance

Short duration and photo insensitive varieties fit well in many intensive cropping systems across the country. Spring/Summer mungbean especially help in sustaining the productivity levels of *rice-wheat* cropping system of Indo-Gangetic belt of northern India without any competition to rice or wheat, with additional yield of 10-15 qtls/ha.

5.4 Growth Rate

From 1980-81 to 2002-03, the total acreage under pulses has almost slightly (\pm) being showed, however, a sudden jump in production and productivity was observed in 2003-04 and 2010-11 for production with 96% and 160% growth rate whereas, for productivity it was 66% and 127% (Table 5.1).

(Table-5.1): Growth rate of mungbean

(Area- Million ha, P- Million tones, Y-kg/ha, Growth Rate (GR)- %)

Voor			Mungbo	ean		
Year	Area	GR	Prod.	GR	Yield	GR
1980-81	2.84		0.98		344	
1985-86	3.00	5.6	1.18	20.1	392	13.7
1990-91	3.36	11.7	1.38	17.7	413	5.4
1995-96	2.72	-19.1	1.01	-27.1	372	-10.0
1996-97	3.06	12.8	1.32	30.7	431	15.9
1997-98	3.02	-1.5	0.95	-28.0	315	-26.9
1998-99	3.05	1.2	1.16	22.3	380	20.8
1999-00	2.91	-4.9	1.08	-7.2	371	-2.5
2000-01	3.01	3.5	1.02	-5.0	340	-8.3
2001-02	3.09	2.6	1.11	8.5	360	5.8
2002-03	3.01	-2.3	0.87	-21.9	288	-20.0
2003-04	3.55	17.7	1.70	96.2	480	66.7
2004-05	3.34	-5.8	1.06	-37.9	317	-34.0
2005-06	3.11	-7.0	0.95	-10.5	304	-3.9
2006-07	3.19	2.8	1.12	17.9	349	14.7
2007-08	3.73	16.7	1.52	36.5	409	17.0
2008-09	2.84	-23.7	1.03	-32.1	364	-10.9
2009-10	3.07	8.0	0.69	-33.1	226	-38.0
2010-11	3.51	14.3	1.80	160.0	513	127.6
2011-12	3.39	-3.5	1.63	-9.2	483	-6.0
2012-13	2.72	-19.7	1.19	-27.4	436	-9.6
2013-14	3.38	24.4	1.61	35.3	475	8.8
2014-15	3.02	-10.7	1.50	-6.4	498	4.9
2015-16	3.83	26.8	1.59	6.0	416	-16.4
2016-17	4.33	13.0	2.17	36.4	501	20.4
2017-18*	4.26	-1.6	2.01	-7.4	472	-5.8

Source: DES, Ministry of Agri. & FW (DAC&FW), Govt. of India; 2017-18*- IVth Adv. Est.

(Table -5.2): Per cent share to total pulses

(Area- lakh ha, Production-Lakh Tones, Yield-kg/ha)

Year		Mung		Total Pulses			% Share to Total		
	1]	Pulses	
	A	P	Y	A	P	Y	A	P	YI
1980-81	28.44	9.80	344	224.45	106.24	473	12.7	9.2	73
1990-91	33.55	13.84	413	372.55	203.68	547	9.0	6.8	75
1995-96	27.15	10.09	372	331.77	170.10	513	8.2	5.9	72
2000-01	30.08	10.23	340	203.48	110.75	544	14.8	9.2	62
2001-02	30.87	11.11	360	220.08	133.68	607	14.0	8.3	59
2002-03	30.15	8.67	288	204.96	111.25	543	14.7	7.8	53
2003-04	35.48	17.02	480	234.58	149.05	635	15.1	11.4	76
2004-05	33.41	10.58	317	227.63	131.30	577	14.7	8.1	55
2005-06	31.09	9.46	304	223.91	133.84	598	13.9	7.1	51
2006-07	31.94	11.15	349	231.92	141.98	612	13.8	7.9	57
2007-08	37.27	15.23	409	236.33	147.62	625	15.8	10.3	65

Year		Mung		Total Pulses			% Share to Total Pulses		
	A	P	Y	A	P	Y	A	P	YI
2008-09	28.43	10.35	364	220.94	145.66	659	12.9	7.1	55
2009-10	30.70	6.92	226	232.82	146.62	630	13.2	4.7	36
2010-11	35.08	18.00	513	264.02	182.41	691	13.3	9.9	74
2011-12	33.87	16.34	483	244.62	170.89	699	13.8	9.6	69
2012-13	27.19	11.86	436	232.57	183.42	789	11.7	6.5	55
2013-14	33.83	16.05	475	252.18	192.55	764	13.4	8.3	62
2014-15	30.19	15.03	498	235.53	171.52	728	12.8	8.8	68
2015-16	38.28	15.93	416	249.11	163.48	656	15.4	9.7	63
2016-17	43.27	21.65	500	294.47	231.31	785	14.7	9.3	63
2017-18*	42.57	20.09	472	299.93	252.35	841	14.2	8.0	56

Source: DES, Ministry of Agri. &FW (DAC&FW), Govt. of India; 2017-18*- IVth Adv. Est.

5.5 Production Trends

5.5.1 National Scenario: Plan Periods

Plan-wise performance revelaed, the highest area VII^{th} Plan and production & Productivity increased during XII^{th} plan over the previous plan period. The details plan-wise performance is given below:

(Table-5.3): Plan-wise national scenario

(Area-Mha, Production-MTonnes, Yield-kg/ha)

Plan	Area	%COPP	Prod.	% COPP	Productivity	%COPP
Sixth Plan (1980-85)	2.88		1.12		390	
Seventh Plan (1985-90)	3.13	8.77	1.24	10.42	396	1.51
Eighth Plan (1992-97)	2.92	-6.71	1.21	-2.71	413	4.28
Ninth Plan (1997-02)	3.01	3.19	1.06	-11.84	353	-14.57
Tenth Plan (2002-07)	3.24	7.53	1.14	6.89	351	-0.60
Eleventh Plan (2007-2012)	3.31	2.02	1.34	17.50	404	15.16
Twelfth Plan (2012-2017)	3.46	4.53	1.61	20.14	465	15.09

Source:DES, Twelfth Plan (2012-2017); % COPP is percentage Changeover previous plan




Fig.- 5.1: Mungbean: Plan-wise Scenario

- *Tenth Plan (2002-2007):* The total area under Moong during ninth plan was 32.41 lakh hectares with production of 11.38 lakh tones. Rajasthan stands first in respect of area (22.54%) followed by Maharashtra (17.06%) and A.P. (14.73%). The maximum contribution of production was in the state of Maharashtra (21.61%) followed by Rajasthan (20.23%) and A.P. (15.81%). The highest yield was recorded by the state of Punjab (800 kg/ha) followed by Bihar (548 kg/ha) and Jharkhand (538 kg/ha) with the overall National yield average of 351kg/ha. The lowest yield was recorded in Karnataka (146 kg/ha) followed by Odisha (238 kg/ha) and Chhattisgarh (235 kg/ha).
- *Eleventh Plan (2007-2012):* The total area covered under moong in India was 33.32 lakh hectares with a total production of 13.52 lakh tonnes. Moong is a common crop grown in most of the states. The share of area and production are some what homogeneous in the Moong growing states. However, during the Plan Period, the coverage of area and its production was maximum in Rajasthan (31.21% & 31.68%) followed by Maharashtra (15.26% & 18.57%) and Andhra Pradesh (10.36% & 10.62%). The highest yield was recorded by the state of Punjab (778kg/ha) followed by Bihar (617 kg/ha) and Jharkhand (571 kg/ha). The National yield average was of 406 kg/ha. The lowest yield was observed in Karnataka (202 kg/ha) followed by Odisha (270 kg/ha) and C.G. (250 kg/ha).
- Twelfth Plan (2012-2017): The total area covered under moong in India was 34.55 lakh hectares with a total production of 16.11 lakh tonnes. The coverage of area and its production was maximum in Rajasthan (33.02% &30.95%) followed by Maharashtra (11.52 %& 10.34%) of the total area and production. Karnataka ranked third in area (8.83%) and Madhya Pradesh is on third position for production (9.17%). The highest yield was recorded by the state of Punjab (848 kg/ha) followed by Jharkhand (703 kg/ha) and Andhra Pradesh (699 kg/ha). The lowest yield observed in the state of Karnataka (226 kg/ha) followed by C.G. (326 kg/ha) and Odisha (347 kg/ha).

During the last three Plan Period area fluctuating, however, production and productivity showed increasing trend.

(Table-5.4):Plan-Wise StatesøScenario ó Major Stateøs

(Area- lakh ha, Production-Lakh Tones, Yield-kg/ha)

State		X th Plan	% to AI	XI th Plan	% to AI	XII th Plan	% to AI
	A	4.77	14.73	3.45	10.36	1.56	4.51
Andhra Pradesh	P	1.80	15.81	1.44	10.62	1.09	6.76
	Y	377		417		699	
	A	1.86	5.74	1.67	5.01	1.64	4.75
Bihar	P	1.02	8.92	1.03	7.59	1.04	6.44
	Y	548		617		632	
	A	0.17	0.53	0.16	0.49	0.16	0.48
Chhattisgarh	P	0.04	0.38	0.04	0.30	0.05	0.33
	Y	235		250		326	
	A	1.73	5.35	2.20	6.62	1.45	4.19
Gujarat	P	0.70	6.12	1.06	7.84	0.76	4.72
	Y	405		482		526	
	A	0.22	0.67	0.27	0.80	0.44	1.27
Haryana	P	0.07	0.58	0.12	0.86	0.27	1.66
	Y	318		444		611	
	A	0.13	0.40	0.21	0.63	0.24	0.68
Jharkhand	P	0.07	0.63	0.12	0.86	0.17	1.03
	Y	538		571		703	
	A	4.11	12.69	3.76	11.27	3.05	8.83
Karnataka	P	0.60	5.26	0.76	5.61	0.69	4.29
	Y	146		202		226	
	A	0.85	2.62	0.86	2.58	2.89	8.38
Madhya Prd.	P	0.27	2.39	0.28	2.07	1.48	9.17
	Y	318		326		510	
	A	5.53	17.06	5.08	15.26	3.98	11.52
Maharashtra	P	2.46	21.61	2.51	18.57	1.67	10.34
	Y	445		494		418	
	A	2.23	6.87	2.59	7.76	2.89	8.36
Odisha	P	0.53	4.67	0.70	5.16	1.00	6.21
	Y	238		270		347	
	A	0.15	0.46	0.09	0.26	0.38	1.10
Punjab	P	0.12	1.04	0.07	0.50	0.32	1.99
	Y	800		778		848	
	A	7.31	22.54	10.40	31.21	11.41	33.02
Rajasthan	P	2.30	20.23	4.28	31.68	4.99	30.95
	Y	315		412		437	
	A	1.25	3.87	1.54	4.63	1.90	5.49
Tamilnadu	P	0.54	4.73	0.54	3.99	1.09	6.79
	Y	432		351		577	
	A			0.72		1.24	3.59
Telangana	P			0.39		0.71	4.41
	Y			545		572	
	A	0.74	2.30	0.79	2.38	0.90	2.59
				0.45	3.36	0.48	2.97
Uttar Pradesh	P	0.35	3.05	0.45	3.50	0.40	2.71
Uttar Pradesh		0.35 473	3.05	570	3.30	533	2.51
Uttar Pradesh All India	P		3.05		3.30		2.51
	P Y	473	3.05	570	3.30	533	2.71

Source: DES, DAC&FW,ND (*Twelfth plan - Avg. 2012-13 to 2016-17)




Fig.-5.2: Mungbean Area: State@s Contribution (XIIth Plan)




Fig.-5.3: Mungbean Production: State@s Contribution (XIIth Plan)

5.5.3 District Scenario (2016-17) – Potential Districts

Analysing the Intra-state, status of the moong crop, district Nagaur with 12.90% area and 15.51% production tops in the country. District-wise area, production and yield of top ten district of India in respect of production, are presented below which contributed 46.56% and 43.49% percent of area and production of the country (Table 6 5.5).

The yield of most of the potential districts were below the National average yield need to be adopt Improved Package of Practices of the greengram to increase the production in districts as well as country.

(Table-5.5): Top potential districts: Mungbean

{Area-Lakh ha, Production-Lakh Tonnes, Yield-kg/ha}

Name of	State	A	rea	Prod	uction	Yie	ld
districts		Area	% to India	Prod.	% to India	Yield	YI
Nagaur	Rajasthan	5.58	12.90	3.36	15.51	602	120
Jodhpur	Rajasthan	2.77	6.40	1.10	5.07	396	79
Pali	Rajasthan	2.79	6.44	1.07	4.92	382	76
Ajmer	Rajasthan	1.05	2.43	0.74	3.41	701	140
Jaipur	Rajasthan	1.07	2.46	0.73	3.35	681	136
Ganjam	Odisha	1.08	2.50	0.53	2.45	490	98
Churu	Rajasthan	1.76	4.08	0.50	2.29	281	56
Ganganagar	Rajasthan	0.84	1.94	0.48	2.23	577	115
Tonk	Rajasthan	0.67	1.54	0.48	2.22	719	144
Jalore	Rajasthan	1.24	2.87	0.44	2.04	355	71
Total above		18.85	43.56	9.42	43.49	500	
All India		43.27		21.65		500	

Source: State Department of Agriculture




Fig.-5.4: Mungbean: Potential District

(Table-5.6): State-wise yield gap and additional return

(Yield: kg/ha; Return: Rs./ha.)

State	Yi	eld (kg/ł	ıa)	Gap over	r FP	Gap o	ver	Yield		tional	
				SAY		Y	2016-17	return by			
									_	ng yield	
			~		0.1		0.1			gap (Rs/ha)	
	IP	FP	SAY	Actual	%	Actual	%		FP	SAY	
Kharif											
Tripura	1016	782	513	234	30	503	98	0	13104	28168	
Gujarat	766	606	501	160	26	265	53	454	8960	14840	
Karnataka	655	563	214	92	16	441	206	277	5152	24696	
Maharasthra	546	430	418	116	27	128	31	585	6496	7168	
Rajasthan	773	600	436	173	29	337	77	495	9688	18872	
Tamil Nadu	1000	796	588	204	26	412	70	808	11424	23072	
Manipur	642	479	0.00	163	34	642	0	0	9128	35952	
WB	1229	915	838	314	34	391	47	893	17584	21896	
Average	781	608	435	173	28	345	79	585	9682	19339	
Rabi											
AP	1398	1228	704	170	14	694	99	490	9520	38864	
Rice fallow											
AP	1094	947	704	147	16	390	55	490	8232	21840	
Odisha	825	498	359	327	66	466	130	471	18312	26096	
Average	960	723	532	237	33	428	80	484	13272	23968	
Summer/Spr	ing Mun	gbean									
Bihar	939	735	603	204	28	336	56	548	11424	18816	
Uttar	1176	052	700	222	22	467	"	C21	12400	26152	
Pradesh	1176	953	709	223	23	467	66	631	12488	26152	
Haryana	524	487	576	37	8	-52	-9	735	2072	-2912	
Punjab	1143	0.00	857	1143	0	286	33	986	64008	16016	
Assam	875	622	624	253	41	251	40	686	14168	14056	
Average	931	559	674	372	66	257	38	717	20832	14414	

Source-Annual Report- 2016-17, GoI, DPD, Bhopal (Ave. 2013-14 to 2015-16)

State Average Yield – E&S (Ave. 2011-12 to 2015-16)

IP: Improved Practise FP: Farmers Practise SAY: State Avergae Yield

5.6 Mungbean Agronomy

5.6.1 Botanical Description

It is a small herbaceous annual plant growing to a height of 30 to 100 cm with a slight tendency to twining in the upper branches. Depending upon the plant type and nature of crop being grown, central stems are more or less erect while side branched are semi erect, leaves are trifoliate with long petioles. Both the stem and leaves are covered with short hairs, generally shorter than urd. Flowers are various shades of yellow colour produced in cluster of 10-20 in axillary racemes. Crop is fully self-fertile. Pods are 6-10 cm long, hairy and round having 7-10 seeds inside. Hilum is white and flat. Germination type epigeal and colour of cotyledons is yellow.

5.6.2 Production Technology

- Climate: The crop needs high temperature, less humidity and moderate rainfall of about 60-80 cm. Water logging is fatal for root development and nitrogen fixation during early vegetative stage. Crop is generally grown as rain fed but under assured irrigation during summer in Indo Gangetic plains of Northern India.
- Soil and Field Preparation: Best soil for its cultivation is loam soil with good drainage. The crop should not be raised on alkaline, saline or waterlogged soils. A well prepared seedbed is required for proper germination and establishment of the crop. For this give 2 ó 3 ploughings followed by planking to make the seedbed free from clods and weeds. For the summer/spring cultivation after the harvesting of last crops, the tillage should be done after irrigation.
- Sowing Time: Mungbean should be sown during the last week of June to mid or first week of July. For the summer or spring crop, mungbean should be sown after the harvest of last crop (potato, sugarcane, mustard and cotton, etc). The first fortnight of March is most suitable for spring/summer cultivation. Late sown mungbean takes more loss at the time of flowering stage due to high tempreture and yield affected.
- Growing seasons: Mungbean is cultivated in all the three seasons viz., kharif, rabi and zaid (spring/summer). As a catch crop, in between rabi and kharif crops, after the harvest of rabi crops, like wheat, potato, mustard, sugarcane, etc., it is grown under irrigated conditions/rice fallow areas.

During 2016-17, mungbean recorded the ever highest production at 20.70 lakh tons (kh-15.28 + rabi- 5.42 lakh tons), which is 38% higher over the Normal and 30% higher than the last year. Rajasthan, with 48% of total all India production during kharif and >31 % collectively in a crop year, ranks Ist in the country. During, rabi, TN with 19 % of national production stands at Ist position.

More than 90% of Mungbean production comes from 10 states, namely Rajasthan, Maharashtra, TN, AP, Bihar, MP, Odisha, Gujarat, Telangana and Karnataka.

- **A. Kharif greengram-** The kharif crop is grown both as a sole and as an intercrop. Since it is largely grown as a rainfed crop, the sowing time usually depends on the onset of the monsoon in the different regions. *Normally the sowing is done from mid-June to second week of July*, which is considered optimum. Delayed sowings would result in progressive decline in yields.
- **B.** Rabi greengram- Rabi greengram is grown in an area of approx. > 9.54 Lha (Avg. 2011-12-2015-16). The major states are Odisha, Tamil Nadu, Bihar, Andhra Pradesh, Madhya Pradesh Haryana, Uttar Pradesh, Gujarat, Punjab and West Bemngal. There is a great scope of increasing the area under rabi greengram in rice fallows on residual moisture or under irrigated conditions in the southern states.

An area of approximetly 12-20 lakh ha is covered during spring summer season, however, this is not recorded seperatly by the DES and is finally a part of total area coverage during rabi season.

C. Spring/Summer greengram-The optimum sowing time for mungbean in the northern plains ranges from 15 March to 15 April, as this also enables the crop to be harvested safely before the onset of the monsoon. A fine balance has to be achieved in choosing the correct sowing time which will avoid the relatively low temperature of winter and at the same time would not delay harvest for timely sowing of the kharif crop. Also, a late-sown crop could be caught in the pre-monsoon showers towards the end of June. Late March or early April sowings are most appropriate for north-Indian conditions. However, for optimum yields the sowing is to be advanced to the early part of March. Sowing time for the summer crop is late January in Tamil Nadu, Andhra Pradesh and Karnataka; February in Odisha and West Bengal; March in Bihar, Madhya Pradesh and Rajasthan; and first fortnight of April in Uttar Pradesh, Haryana and Punjab.

• Advantages of spring/summer cultivation

- The crop has very little or no infestation of insect-pest and diseases due to high temperature and desicating winds.
- The crop/varieties take lesser time to mature (normally 60-65 days).
- It suits well after wheat, mustard, potato and late rice in West Bengal.
- The cropping intensity can be increased.
- The area and production can be increased under pulses without eliminating other crop to be grown during kharif season.
- It utilizes the residual soil fertility when grown after heavily fertilized crops like potato, wheat and winter maize.
- In return, it adds at least 30-35 kg available nitrogen/ha through Rhizobium fixation which may be adjusted while applying fertilizers in following kharif season crop.
- After picking pods, the foliage can be incorporated into soil as green manure *in-situ* to add organic matter into the soil as bonus for boosting soil fertility and improving physical conditions of the soil.
- It controls the weeds and checks wind erosion during summer.

• Recommendations for successful cultivation of Rabi Mungbean

- Select high yielding varieties resistant to YMV, leaf curl, powdery mildew and drought.
- Use only duel inoculated (Rhizobium + PSB) seeds for better root development and harnessing maximum ÷Nø fixation.
- Treat the seed with Emidacloprid @ 5 ml/kg followed by Mancozeb @ 3g/kg, two days before seed inoculation, as protection against incidence of seedling pest and diseases.
- Use a seed rate of 12-15 kg/ha for upland and 30 kg/ha for rice fallow areas.
- Use basal application of 20 kg N + 50 kg P_2O_5 + 200 kg Gypsum/ha during field preparation, 3-4 cm below and side of the seeds. Balanced Fertilizer application is

- based on soil testing.
- Keep the field free from weeds up to 30 DAS by one hand hoeing.
- In Rice fallow area, Echinochloa (barn yard grass) is the major weed, can be control by mixing Benthiocarb @ 5 L in 50 kg dry sand as broadcast, 3 to 4 days before harvest of paddy.
- For control of Cuscuta spp; post emergence, sand mix application of Pendimethalin and flucholarlin at 2.0 L + 1.5 L respectively in 50 kg sand gives best results.
- One irrigation at 35 DAS and 2% spray with urea or DAP at pre-flowering, flowering and pod development, is often associated with high jump in grain yield.
- Seed Rate & Spacing & Method: During Kharif season 15-20 kg seed/ha should be sown in rows 45 cm apart while during Rabi and Summer 25-30 kg seed /ha sown in rows 30 cm apart. 30 cm x 10 cm. row spacing is considered optimum, for modern varieties. As a companion crop with sugarcane seed rate should be 7-8 kg/ha. The plant-to-plant distance should be maintained (atleast 5 cm). Sowing can be done behind the local plough or with the help of seed drill.
- Seed Treatment: Treat the seed with Thirum (2 gm.) +Carbendazim (1 gm.) or Carbendazim & Kepton (1 gm + 2 gm) to control the soil & seed germinated disease. For sucking pest control seed treatment with Imidacloprid 70 WS @ 7g/kg seed. It is also desirable to treat the seed with Rhizobium and PSB culture (5-7gm/kg seed).
- Varieties: The variety may be selected per the adaptability to the region, agro-eco sitiation, sowing time and the cropping system.
- Cropping systems: The important crop rotations with moongbean are given as under:

Rice-Wheat-Moong (<u>summer</u>)
Maize+Moong-Wheat-Moong
Maize(early)-Potato(early)-wheat-Moong,
Sugarcane+Moong (<u>summer 1:2</u>)
Cotton + Greengram (1:3 in Central India 60/90 cm paired row)

Rice-Rice-Greengram(<u>south India</u>)
Moong-Wheat/Barley
Sunflower+Moong (<u>summer 2:2</u>)
Moong+Pigeonpea (2:1)

- Intercropping: During kharif, mungbean are grown generally with maize, pearl millet, pigeonpea and cotton as intercrop. During spring, mungbean is grown in 2:1 ratio with sugarcane with row to row distance of 90 cm. & from mungbean 30 cm distance maintain. Intercropping with sunflower is also suitable with ratio (2:6).
- Water Management: Generally the kharif crop requires one life saving irrigation, which may be applied during the early pod formation stage. For the summer/spring mungbean, 364 irrigations are required. Apply first irrigation after 20-25 days of sowing and repeat after 10-15 days as per need. One irrigation before flowering and another at pod-filing stage would ensure healthy seeds. Water logging in the field should be avoided at all cost. No irrigation should be given when the crop is in full bloom stage.

- **Plant nutrient management**: The response to phosphorus is highest on red and laterite soils. Application of P₂O₅ @ 30-40 kg/ha along with a starter dose of 10-15 kg nitrogen is adequate. Phosphorus application has always a significant effect in increasing the yields. Seeds should be treated with an efficient Rhizobium culture for obtaining higher yields. Rhizobial inoculation may reduce the nitrogen requirement of the crop. Fertilizer application is based on soil testing.
- Weed management: The highest crop yield was obtained when weeds were removed 35 days after sowing. Any further delay in weed removal results in a corresponding decrease in yield. A maximum of 2 hand-weedings in the initial stages of crop growth up to 30-35 days, is adequate to take care of the weed problem. However, whenever labour is in short supply or the rainfall pattern does not allow early hand-weeding, herbicides need to be used. Pre-emergence application of Lasso or Tok E-25 @ 2kg ai/ha in 1,000 litres of water ensures complete weed control. Application of Pendimethalin (PI) + Imazethapyr (POE) 1250+100 g a.i./ha at 0-3 (PI) and 20-25 (POE) DAS control weeds.
- Harvesting and threshing: Mung should be harvested when more than 80 per cent pods mature. One or two rounds of picking of pods are also recommended to avoid losses due to shattering. The plants are cut with the sickle and dried on the different floor. These are then threshed by beating with sticks or by trampling with bullocks.
- Yield: A well-managed crop may yield about 15-20 quintals of grain per ha.

5.7 Plant Protection Measures

There are several important disease of mung, yellow mosaic, leaf crinkle, leaf curl, anthracnose, cercospora leaf spot are important one.

A. Diseases

i) Yellow Mosaic Virus

Symptoms: This disease is caused by the mung bean yellow mosaic virus (MYMV) belong into Gemini group of viruses, which is transmitted by the whitefly (*Bemisia tabaci*). The tender leaves show yellow mosaic spots, which increase with time leading tocomplete yellowing. Yellowing leads to less flowering and pod development. Early infection often leads to death of plants.

Control Measure

Diseased plants should be rogued out to prevent further spread of the disease; ii) In order to prevent whitefly (Bemisia spp.) infestation spray with triazophos 40 EC @ 2.0 ml/l or malathion 50 EC @ 2.0 ml/l or oxydemeton methyl 25


EC @ 2.0 ml/l at 10-15 days intervals if required; iii) Grow tolerant/resistant varieties like Narendra Mung1, Pant Mung 3, PDM 139 (Samrat), PDM 11, MUM 2, ML 337, IPM 02-14, MH 421, SML 832 etc.

ii) Leaf Curl

Symptoms: The symptoms are visible first in third leaf after three to four weeks of sowing. These are characterized by enlargement of leaves followed by their crinkling. Later the leaves become thicker an leathery. The affected plants, however, do not die till the harvest of the crop.

Control Measures

i) Treat the seeds with imidacloprid 70 WS@ 5 ml/kg; ii) Foliar spray of insecticide (dimethoate 30 EC @ 1.7 ml/ha) on 30 days after sowing; iii) Rogue out the infected plants and Field sanitation; iv) Use resistant varieties like D-3-9, K 12, ML 26, RI 59, T-44.



iii)Anthracnose

Symptoms: The fungus Colletotrichum spp. Is the causal organism affecting aerial plant parts, however, the leaves and pods are more vulnerable. The characteristic symptoms of this disease are circular brown sunken spots with dark centers and bright red orange margins on leaves and pods. Infection just after germination causes seedling blight.

Control Measures

Hot water seed treatment at 58°C for 15 minutes has been found effective in checking the seed-borne infection and increasing



proportion of seed germination; ii) Seed treatment with carbendazim 50 WP @ 2g/kg of seed helps in eliminating the seed borne infection; iii) Spray the crop with carbendazim @ 2 g/liter of water with first appearance of symptoms on the crop and repeat after 15 days (if necessary).

iv)Cercospora leaf spot

Symptoms: It is most important fungal disease of mungbean. Leaf spots are circular and irregular in shape with brown to greyishwhite centres and reddish-brown to dark brown margins. Such spots are also visible or pods are the affected pods become blackened.

Control Measures

i) Field sanitation, crop rotation, destruction of infected crop debris; ii) Opt for resistant varieties as per recommendation of local agricultural authorities (Like- LM 113, LM 168, LM 170,



JM 171, Ganga 1, HUM 12, Pusa 06722); iii) Treat the seeds with thiram or captan @ 2.5g/kg of seed; iv) On appearance of the symptoms spray with carbendazim 50 WP @1.0 g/liter or mancoze 75 WP @ 2.0 g/liter or hexaconazole (contaf 5 % EC) @ 1 ml/ liter of water. Subsequent spray should be done after 10 to 15 days, if required. Spraying with copper oxychloride @ 3 to 4 g /liter water has also been found effective in management of the disease.

B. Insect-Pest


Numerous insect pests attack the mungbean. The loss in the production caused by them may reach up to 70% depending upon the severity of attack. Some common insect pests of mungbean and their management are as follow:

i) White fly

Nature of damage:The infested plants become very weak showing downward cupping of the leaves giving a sickly look and the plant may die Insect secretes honey dew on leaves results blackening of leaves, drastically reducing photosynthetic rate and drying of leaves. Whitefly is a vector of number of viral diseases especially mungbean yellow mosaic virus (MYMV).

Control Measures

i) Seed treated with Dimethoate 30 EC @ 5ml/kg; ii) Foliar spray of Triazophos 40 EC @ 2.0 ml/l or Malathion 50 EC @ 2.0 ml/l $\,$





Nature of damage: The thrips nymphs and adults feed on stigma inside the flower, flower sheds before opening and there is elongation of terminal shoot. Plants attain a bushy growth and the crop looks dark-green in colour, bearing few pods with shrivelled grains.

Control Measures

i) Seed treatment with Thiomethoxam 70 WS @ 2 ml /kg seed + foliar spray of Thiomethoxam 25 WG 0.2 g/liter of water of is quite effective in controlling bean thrips; ii) Spray neem seed

kernel extract (50 g/l) and neem oil 3000 ppm @ 20 ml/l; iii) Opt for resistant varieties (Mungbean: PIMS 2, PMS 3, 12-333, Co 3, ML 5, ML 337; iv) Timely irrigation at an interval of 15 days results in low build up of thrips; v) The incidence of bean thrips can be minimized by intercropping mungbean with cotton; vi) Spray of Triazophos 40 EC

@ 2.0 ml/liter, Ethion 50 EC @ 2 ml/liter of water.

iii) Stem fly

Nature of Damage: Stem fly (*Ophiomyia phaseoli*) maggots mine the leaves or bore into the leaf petiole or tender stem resulting in withering, drooping and death of plant. The characteristic symptoms of damage include drooping of the first two leaves and yellowing of plants. It can cause 5 ó 20% damage in mungbean.

Control Measures

i) Follow clean cultivation, crop rotation, earthing up, growing trap crop, destroying alternative hosts like *Solanum nigrum* to minimize the stem fly incidence; ii) Opt for resistant varieties





(Mungbean: CoGG 912 & CoGG 917; Urdbean: CoBG 671 & AC 222); iii) Seed soaking either in imidacloprid 17.8 SL @ 5.0 ml/kg seed in 100 ml water for one hour or thiomethoxam 25 WG @ 5.0 g/kg seed in 100 ml water to avoid early incidence of stem fly is recommended; iv) Spray either Imidacloprid 17.8 SL @ 0.2ml/l or thiomethoxam 25 WG @ 0.3g/lit at 15 days after sowing.

iv) Bihar Hairy Caterpillar

Nature of damage: Female moths lay eggs on plants in a field. Young caterpillar eat away all the green matter of the leaves and it can be easily recognized by perforated, dusty white coloured leaves in the field. The grown-up caterpillars feed voraciously on leaves, soft stems and branches. The insect totally denude the crop within few days resulting in total failure of the crop.



Control Measures

i) Uproot the damaged plants along with the young larvae at the gregarious phase and burry under the soil; ii) Spray of Quinalphos 25 EC @ 2.5 ml/liter or Dichlorvos 10 EC @ 1.0 ml/liter or Fenvalerate 20 EC @ 1.87 ml/liter of water or dusting with Fenvalerate 0.4% @ 15 kg/ha.

5.8 Recommendation to achieved higher production

- Deep summer ploughing once in 3 years.
- Seed treatment should be done before sowing.
- Application of fertilizer should be based on soil test value.
- In kharif season sowing should be done by ridge & furrow method.
- Yellow mosaic resistant/ tolerant varieties Narendra Mung1, Pant Mung 3, PDM 139 (Samrat), PDM 11, MUM 2, ML 337, IPM 02-14, MH 421, SML 832 etc choose as per suitability of region.
- Weed control should be done at right time.
- Adopt integrated approach for plant protection.

(Table-5.7):Recommended mungbean varieties/characteristics

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. Yield (q/ha)	Days to maturity	Special characteristics
ADT-3	TNAU	1991	Tamil Nadu	10.7	65-70	Tolerant to YMV. Seed small
Co-5	TNAU	1991	Tamil Nadu	9.0	70-75	Tolerant to YMV. Seed small
MUM-2	Meerat University	1992	Punjab, Haryana, Delhi & West UP	12.0	60-70	Res. To YMV, small seeded, early
BM-4	MAU	1992	M.P., Maharashtra, Gujarat.	10-12	65	Early, Tol.to YMV and PM, Bold Seeded
Phule M 2	MPKV	1992	Maharashtra	6.9	65	Tolerant to YMV, early, small seed
AKM-8803	PKV	1992	Maharashtra.	10.5	65-70	Tolerant to YMV. Seed small
Narendra Mung-1	NDUAT	1992	Uttar Pradesh.	10.0	60-70	Tolerant to YMV.
AKM-8803	PKV	1992	Maharashtra.	10.5	65-70	Tolerant to YMV.
Asha	CCSHAU	1993	Haryana.	12.0	75-80	Tolerant to YMV.
TARM-2	BARC/PKV	1994	Maharashtra.	9.5	65	Tolerant to PM.
Pusa-9072	IARI	1995	SZ (KN, A.P., Odisha, TN (Rabi).	8-10	65-75	Tolerant to Powdery Mildew. Rabi
Warangal-2 (WCG-2)		1995	Andhra Pradesh.	14.0	65-70	Suitable for all Season, Tolerant to YMV
Madhira-295	ANGRAU	1995	Andhra Pradesh.	14.0	65-70	Tolerant to YMV
LGG-407 (Lam 407)		1995	Andhra Pradesh.	14.0	70-75	Tolerant to YMV.
JM-721	JNKVV	1996	Madhya Pradesh.	12.4	70-75	Tolerant to PM.
ML-613	PAU	1996	Punjab.	13.0	84	Res. To YMV, Bacterial leaf spot and Pod-leaf spot. Seed med. Bold
SML-134	PAU	1996	Punjab.	11.0	68	For summer/spring.
PDM-84-178		1996	Andhra Pradesh.	8.1	65-70	Tol. To YMV & PM, suitable for summer and early kharif.
TARM-1	BARC/PKV	1997	Maharashtra.	8-12	85	Res.to PM, Suitable for Rabi. Small seed

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. Yield (q/ha)	Days to maturity	Special characteristics
Pant Mung-4	GBPUAT	1997	Eas. UP, Assam, Bihar, W.B	7.5	68	Resistant to YMV.
HUM-1	BHU	1999	Gujarat, MS, MP, TN, KN	8-9	60-65	Res. To YMV, Summer season
CO-6	TNAU	1999	Tamil Nadu	10.0	65	Suitable for all season, Resistant to YMV.
Pusa-9531	IARI	2000	M.P., MS, Gujarat, Punjab, Haryana, Delhi, West UP	10-12	60	Res. To YMV, Tolerant to Jassids and whitefly, suitable for summer.
Pusa Vishal	IARI	2000	NWPZ (Punjab,Haryana,Delhi, West UP, North Rajasthan)	11.0	62	Res. To YMV, Tol.to Jassids and whitefly, suitable for summer,very bold seeded (6 g/100 seed)
LAM-460	ANGRAU	2001	Andhra Pradesh	12.0	70-75	Tolerant to YMV
PDM 139	IIPR	2001	Uttar Pradesh.	12-15	50-60	Summer season, Mod.Res. to YMV
Ganga-8 (Gangotri)	RAU, Sri Ganga Nagar	2001	NWPZ (Punjab,Haryana, West UP, North Rajasthan)	9.2	72	Kharif, tolerant to stem fly and pod borer.
OUM-11-5	OUAT	2002	SZ (Karnataka, AP, Odisha, TN).	7.0	62	Kharif, Moderately resistant to diseases
Malviya Jagriti (HUM-12)	BHU	2003	U.P., Bihar, Jharkhand, W.B.	11-12	66	Mod. Res. YMV, CLS, Summer Season
IPM 99-125	IIPR	2004	NEPZ (Eas. UP, Bihar, W.B.).	10.0	66	Res. To YMV, Summer Season
TM 99-37	BARC	2005	NEPZ (Eastern UP, Bihar, W.B.).	11.0	65	Mod. Res. To YMV, Summer
COGG 912	TNAU	2005	SZ (Karnataka, A.P, Odisha, TN).	8.0	62	Res. To YMV, CLS, Kharif
Kamdeva (OUM 11-5)	OUAT	2004	SZ (Karnataka, A.P, Odisha, TN)	8.0	46-69	Mod. Rest. To PM, MYMV & CLS
Muskan (MH-96-1)	CCS HAU	2004	Haryana	15.0	70-75	Resistant to YMV, Anthracnose and Leaf Crinkle

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. Yield (q/ha)	Days to maturity	Special characteristics
Ganga-1 (Jamnotri)	ARS, Sri Ganga Nagar	2004	Rajasthan	14	76	Mod. Res. To YMV, CLS, PM, anthracnose, Bacterial leaf blight, Macrophomina & web blight & Rhizopus, Moderate tolerant to white fly and jassids
Shalimar Moong-1	SKUAST	2005	J & K	9	105-115	Res. To leaf spot, pod blight
HUM 16 (Malviya Jankalyani)	BHU	2006	NEPZ (Eastern UP, Bihar, W.B.).	14-16	55-58	Summer, Resistant to YMV, Root Knot and Leaf Crinkle
Tromday pesara (TM-96-2)	ANGRAU	2006	Andhra Pradesh	6	69-73	Rabi & summer, Res. To PM and Cercospora leaf spot
Tromday Jawahar M-3 (TJM-3)	JNKVV	2006	MP	8-10	61-75	Kharif & summer, Resistant to YMV, PM and Rhyzoctonia root rot
ML 818	PAU, Ludhiana.	2007	NWPZ (HR, PN, UP, UK)	10-10.50	72-78	Resistant to MYMV and CLS Moderately resistant to anthracnose & web blight diseases.
PKV Green gold (AKM 9911)	Dr.PDKV, Akola	2007	Vidarbh Region of MS	10-12	64-72	Moderate resistant to pearl millet Podding on top and easy to harvest
BM 2002-1	ARS, Badnapur	2007	Maharashtra	10-12	65-70	Mod. resistant to PM disease.
Trombay Pesara (TM 96-2)	BARC and ANGRAU, Lam	2007	AP	9-10	65-70	Resistant to Powdery mildew
TJM 3	BARC and JNKVV	2007	MP	7.50-10.65	61-75	Resistant to powdery mildew, YMV and CLS.
MH 2-15 (Sattya)	Hisar	2008	NWPZ/CZ(DL, HR, RJ, W. UP)	10.55-13.20	67-72	Resistant to MYMV and CLS
Pant Mung 6	GBPUA& T, Pantnagar	2008	NHZ(TR, MN, J&K, HP)	10.52	69 -101	Resistant to MYMV, CLS, BLS and Lea Crinkle

Pulses in India : Retrospect and Prospects -2018 DPD , Govt. of India, Bhopal\

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. Yield (q/ha)	Days to maturity	Special characteristics
PAU 911(ML 1265)	PAU, Ludhiana	2008	Punjab	12-13	75-80	Fairly resistant to MYMV, BLS and CLS
SML 668	PAU, Ludhiana	2008	Punjab	10-11.33	60	Resistant to MYMV
KM 2241 (Sweta)	CSAU, Kanpur	2009	NHZ(TR, MN, J&K, HP)	10-11	65-70	Resistant to MYMV.
IPM 02-3	IIPR, Kanpur	2009	NWPZ (DL, HR, HP, J&K, PN RJ)	11	70-72	Large seeded, Resistant to MYMV Relatively Tolerant To Major Pests
MGG 207	ARS, Madhira	2009	Andhra Pradesh	12-14	75-80	Tolerant to stress and YMV
KKM-3	ARS, Davangere, Karnataka	2009	Karnataka	8-9	62	Moderately tolerant to PM and YMV and Pod Borer.
WBM-29	Pulses & Oilseed Research Station, Berhampore -WB		West Bengal	9.87-11.14	65-70	Tolerant to root knot nematode, mod resistant to YMV and Leaf Crinkle Virus.
PKV AKM 4 (AKM- 9904)	PKV, Akola	2011	CZ (MH, MP) SZ (KA,OD,TN)	10-11	57-80	Resistant to MYMV
Pusa 0672	IARI, New Delhi	2010	NHZ (J&K, MN, TR)	10	64	Resistant to MYMV
Pant Mung-5	GBPUAT, Pant Nagar	2010	Uttar Pradesh	10-12	62-79	Res. to mungbean yellow mosaic virus CLS and anthracnose.
Madhira Pesara-347 (MGG 347)	ARS, Madhira	2010	AP & and Telanagana	13-15	65-70	Tolerant to Thrips, Stem Fly, YMV Cercospora Leaf Spot
VBN (Gg) 3	NPRC, Vamban, Tamil Nadu	2010	Tamil Nadu	9-10	65-70	Moderately resistant to YMV
Basanti(MH 125)	CCSHAU, Hisar, Haryana	2010	Haryana	12-15	64-70	Posses for high protein content, Resistanto MYMV
TM 2000-2 (Pairy Mung)	BARC, and IGKV, Raipur	2010	CG	10-11	65-70	Resistant to Powdery Mildew
IPM 02-14	IIPR, Kanpur	2011	SZ (AP, KN, TN, OD)	11	62-70	Resistant to MYMV

Pulses in India: Retrospect and Prospects -2018

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. Yield (q/ha)	Days to maturity	Special characteristics
BM 2003-2	ARS, Badnapur	2012	Maharashtra	8-11	65-70	
KM 2195 (Swati)	CSAUAT,Kanpur	2012	UP	10-12	65-70	Resistant to MYMV
SML 832	PAU, Ludhiana	2013	Punjab	11-12	60-65	Tolerant to Thrips and, MYMV
DGGV 2	UAS Dharwad	2014	Karnataka	11-14	70-75	Mod. resistant to Powdery mildew
MH 421	CCSHAU, Hissar	2014	NWPZ/CZ (DL, HR, RJ, W.UP)	10-12	60-61	Resistant to MYMV
BGS 9 (Somnath)	UAS, Raichur	2014	Karnataka	12-13	65-68	Moderately resistant to PM, Bold seed and long pod
Shalimar Mung 2	Srinagar centre, SKUASTA	2015	Kashmir valley	10-12	99-100	Resistant to Cercospora leaf spot and moderate resistant to aphid
CO. (Gg) 8	TNAU, Coimbatore	2015	Tamilnadu	10-12	60-65	Resistant to YMV
MH 318	CCSHAU, Hisar	2015	Haryana	14 in Kharif and 10 in Spring/Summer	60-65	Resistant to MYMV, well under rice wheat rotation
Utkarsh KM 11-584	MS State Seed Corporation	2016	Maharashtra	12-13	60-65	Moderately tolerant to MYMV
Yadadri (WGG 42)	PJTSAU, Hyderabad	2016	Telangana	10-12	55-60	Resistant to MYMV
Sri Rama (MGG 351)	PJTSAU, Hyderabad	2016	Telangana	12-14	60-65	Moderately tolerant to MYMV
RMG 975 (Keshwanand Mung 1)	RARI, Dugapura	2016	Rajasthan	8-9	65-70	Moderately tolerant to MYMV and tolerant to root knot nematode
MSJ 118 (Keshwanand Mung 2)	RARI, Dugapura	2016	Rajasthan	7-8	60-65	Moderately tolerant to MYMV
ML 2056	PAU, Ludhiana	2016	Punjab	11-12	70-75	Resistant to MYMV
GBM 1	NAU, Gujarat	2016	Gujarat	11-12	102-105	Mod. tolerant to MYMV

Pulses in India: Retrospect and Prospects -2018 DPD, Govt. of India, Bhopal\

Variety	Source	Release/	Area of adoption	Ave. Yield	Days to	Special characteristics
		Notification	(Zone/State)	(q/ha)	maturity	
		Year				
IPM 410-3	IIPR, Kanpur	2016	NWPZ/CZ (RJ, PN, HR, DL	11-12	65-70	Resistant to MYMV
(Shikha)			HP, UK, J&K, MP, MH & GJ.			
IPM 205-7 (Virat)	IIPR, Kanpur	2016	A, TN) CZ (MP, & GJ)	10-11	52-56	Resistant to MYMV
SML 1115	PAU, Ludhiana	2016	NEHZ (TR, MN, & MZ)	11-12	65-70	Moderately resistant to MYMV
DGG-1	UAS, Dharward	2017	Karnataka	12-14	70-74	Resistant to powdery mildew
Pant Mung 8	GBPUAT, Pant	2017	Uttarakhand	10-11	78-83	Resistant to MYMV, CLS & PM
(PM 9-6)	Nagar					
Pusa 1371	IARI, New Delhi	2017	NHZ (TR, MN, J&K, HP)	9-10	80-90	Multiple resistant to MYMV, root rot,
						web blight and Anthracnose

Note: AS- Assam, BH-Bihar, DL-Delhi, GJ-Gujarat, HP-Himachal Pradesh, HR-Haryana, JH-Jharkhand, J&K — Jammu & Kashmir, KA-Karnataka, MP-Madhya Pradesh, MH- Maharashtra, MN-Manipur, MZ- Mizorum, OD-Odisha, PN-Punjab, RJ-Rajasthan, TN-Tamil Nadu, TL-Telangana, TR-Tripura, UP-Uttar Pradesh, UK- Uttarakhand, W.UP- Western Uttar Pradesh, WB- West Bengal,

- Res. = Resistant, Tol. = Tolerant, Mod. = Moderately, YMV= Yellow Moasaic Virus, CLS= Cercospora leaf Spot, PM= Powdery Mildew, BLS- Bacterial leaf spot, BLB-Bacterial leaf blight.
- CZ- Central Zone, NWPZ- North Western Plain Zone, NEPZ-North Eastern Plain Zone, SZ-South Zone, NHZ-Northern Hills Zone.

Source: - Project Coordinator's Report, AICRP on MULLaRP, ICAR, IIPR, Kanpur. 2017-18

BLACKGRAM (URDBEAN)





Chapter-6

Black Gram (Urdbean)

Botanical Name : *Vigna mungo L*.

Origin : India Chromosome : 2n = 24

Synonym : Urd, Biri, Mash

6.1 Economic Importance:

Black gram is one of the important pulse crops grown throughout India. It is consumed in the form of \pm dalø(whole or split, husked and un-husked) or perched. Urd differs from other pulses in its peculiarity of attaining a mucilaginous pasty character when soaked in water. It is consumed in variety of waysdiffer the *North to South* in preparation of different regular and popular dishes like *vada*, *idli*, *dosa*, *halwa*, *imarti* in combination with other foodgrains. Also used as a nutritive fodder for milch cattle.

6.2 Nutritive Value

Protein	24%	Calcium	154 mg/100 g
Fat	1.4-1.6%	Phosphorus	385 mg/100 g
Minerals	3.2%	Iron	9.1 mg/100 g
Dietary Fiber	16%	Calorific value	347 Kcal/100 g
Carbohydrate	59.69-63.0%	Moisture	10-11%

It is the richest source of phosphoric acid among pulses (5-6% richer than others)

6.3 Agronomic Significance

Short duration and photo insensitive varieties fit well in different cropping situations, especially intensive crop rotations, including intercropping. The crop is also used for green manuring after picking the pods with its characteristics to fix the atmospheric nitrogen. The plant with deep tap roots binds soil particles and helps in conservation of soil.

6.4 Growth Rate

From 1985-86 to 1990-91, the total Area, production and productivity has almost slightly increased with (9%, 32% & 21%) however, a good result in production and productivity was observed during 2010-11 for production and productivity with 42% and 29% growth rate. (Table 6.1).

(**Table-6.1**): Growth rate of urdbean

(Area-Million ha, P- Million tones, Y-kg/ha, Growth Rate (GR)- %)

V /	Urdbean									
Year	Area	GR	Prod.	GR	Yield	GR				
1980-81	2.83		0.96		339					
1985-86	3.19	12.6	1.24	29.2	389	14.7				
1990-91	3.48	9.3	1.65	32.8	473	21.5				
1995-96	2.80	-19.6	1.32	-19.7	472	-0.1				
1996-97	3.01	7.4	1.35	2.0	448	-5.1				
1997-98	3.06	1.9	1.38	2.2	450	0.3				
1998-99	2.92	-4.8	1.35	-2.0	463	3.0				

Year		Urdbean									
1 car	Area	GR	Prod.	GR	Yield	GR					
1999-00	2.94	0.8	1.33	-1.4	453	-2.2					
2000-01	3.01	2.4	1.30	-2.6	431	-4.9					
2001-02	3.30	9.7	1.50	15.6	454	5.4					
2002-03	3.55	7.5	1.47	-1.7	415	-8.5					
2003-04	3.42	-3.5	1.47	-0.2	430	3.5					
2004-05	3.17	-7.4	1.33	-9.8	419	-2.6					
2005-06	2.97	-6.3	1.25	-6.2	419	0.2					
2006-07	3.07	3.3	1.44	15.9	470	12.2					
2007-08	3.19	4.0	1.46	1.0	457	-2.9					
2008-09	2.67	-16.3	1.17	-19.4	440	-3.7					
2009-10	2.96	10.8	1.24	5.2	418	-5.1					
2010-11	3.25	9.8	1.76	42.4	542	29.7					
2011-12	3.22	-1.0	1.77	0.4	549	1.4					
2012-13	3.15	-2.0	1.97	11.6	625	13.8					
2013-14	3.06	-2.9	1.70	-13.8	555	-11.3					
2014-15	3.25	6.0	1.96	15.3	604	8.8					
2015-16	3.62	11.6	1.95	-0.7	537	-11.1					
2016-17	4.48	23.7	2.83	45.1	632	17.6					
2017-18*	5.44	21.4	3.56	25.8	655	3.6					

Source: DES, Ministry of Agri. &FW (DAC&FW), Govt. of India; 2017-18*- IVth Adv. Est.

(Table- 6.2): Per cent share to total pulses

(Area- lakh ha, Production-Lakh Tones, Yield-kg/ha)

Year		Urd		To	otal Pulses		% share to Total Pulses			
	A	P	Y	A	P	Y	A	P	YI	
1980-81	28.30	9.59	339	224.45	106.24	473	12.6	9.0	72	
1990-91	34.83	16.46	473	372.55	203.68	547	9.3	8.1	86	
1995-96	28.00	13.22	472	331.77	170.10	513	8.4	7.8	92	
2000-01	30.11	12.96	431	203.48	110.75	544	14.8	11.7	79	
2001-02	33.03	14.99	454	220.08	133.68	607	15.0	11.2	75	
2002-03	35.50	14.74	415	204.96	111.25	543	17.3	13.2	76	
2003-04	34.24	14.71	430	234.58	149.05	635	14.6	9.9	68	
2004-05	31.69	13.27	419	227.63	131.30	577	13.9	10.1	73	
2005-06	29.69	12.45	419	223.91	133.84	598	13.3	9.3	70	
2006-07	30.67	14.43	470	231.92	141.98	612	13.2	10.2	77	
2007-08	31.88	14.57	457	236.33	147.62	625	13.5	9.9	73	
2008-09	26.70	11.75	440	220.94	145.66	659	12.1	8.1	67	
2009-10	29.58	12.36	418	232.82	146.62	630	12.7	8.4	66	
2010-11	32.48	17.60	542	264.02	182.41	691	12.3	9.6	78	
2011-12	32.16	17.66	549	244.62	170.89	699	13.1	10.3	79	
2012-13	31.53	19.71	625	232.57	183.42	789	13.6	10.7	79	
2013-14	30.62	16.99	555	252.18	192.55	764	12.1	8.8	73	
2014-15	32.46	19.59	604	235.53	171.52	728	13.8	11.4	83	
2015-16	36.24	19.45	537	249.11	163.48	656	14.5	11.9	82	
2016-17	44.78	28.32	632	294.47	231.31	785	15.5	12.2	80	
2017-18*	54.39	35.62	655	299.93	252.35	841	18.1	14.1	78	

Source: DES, Ministry of Agri. &FW (DAC&FW), Govt. of India; 2017-18*- IVth Adv. Est.

6.5 Production Trends

6.5.1 National Scenario: Plan Periods

Plan-wise performance revelaed, the highest area and production increased during XIIth plan over the previous plan period. The details plan-wise performance is given below:

(Table-6.3): Plan-Wise National Scenario

(Area-Mha, Production-MTonnes, Yield-kg/ha)

Plan	Area	%COPP	Prod.	% COPP	Productivity	%COPP
Sixth Plan (1980-85)	2.86		1.06		373	
Seventh Plan (1985-90)	3.21	12.26	1.39	30.80	434	16.51
Eighth Plan (1992-97)	2.94	-8.42	1.35	-3.09	459	5.82
Ninth Plan (1997-02)	3.05	3.76	1.37	1.60	450	-2.08
Tenth Plan (2002-07)	3.24	6.20	1.39	1.52	430	-4.40
Eleventh Plan (2007-2012)	3.06	-5.56	1.48	6.24	484	12.50
Twelfth Plan (2012-2017)	3.51	14.7	2.08	40.5	592	22.3

DES, Twelfth Plan (2012-2017); % COPP is percentage Changeover previous plan




Fig.-6.1: Urdbean: Plan-wise Scenario

6.5.2 States' Scenario: Plan Analysis (Xth -XIIth)

- *Tenth Plan (2002-2007)*: The total area was 32.39 lakh ha with a total production of 13.96 lakh tonnes. States showed that Maharashtra state stands first in area and second in production (17.15% and 19.53%), whereas A.P. ranked first in Production (20.56%). U.P ranked second in area and third in Production (16.14% and 15.46%) and M.P. stands third in area fourth in production (15.95% and 12.77%) respectively. The highest yield was recorded by the state of Bihar (760 kg/ha) followed by Andhra Pradesh (579 kg/ha) and Jharkhand (548 kg/ha) with the overall National yield average of (431 kg/ha). The lowest yield was observed in the state of Karnataka (187 kg/ha) followed by Odisha (268 kg/ha) and C.G. (286 kg/ha).
- *Eleventh Plan (2007-2012)*: The total production was 14.81 lakh tonnes on an area of 30.58 lakh hectares. As regards the total contribution from states, Madhya Pradesh stand first in respect of area (17.08%) followed by U.P. (16.53%) and Andhra Pradesh (14.87%), whereas in production U.P. stands first (19.00%) followed by Andhra Pradesh (18.37%) and Maharashtra (15.35%). The highest yield was recorded by the state of Bihar (842 kg/ha) followed by Jharkhand (663 kg/ha) and Gujarat (650 kg/ha). The lowest yield was observed in the state of C.G. (294 kg/ha) followed by Odisha (298 kg/ha) and Karnataka (311 kg/ha).
- *Twelfth Plan (2012-2017)*: The total production was 20.82 lakh tonnes on an area of 35.13 lakh hectares. As regards the total contribution from states, Madhya Pradesh stand first in respect of area (24.32%) followed by U.P. (16.73%) and Andhra Pradesh (10.95%), whereas in production M.P. stands first (22.96%) followed by Andhra Pradesh (15.26%) and U.P (14.66%). The highest yield was recorded by the state of Bihar (889 kg/ha) followed by Jharkhand (875 kg/ha) and A.P. (826 kg/ha) the National yield average was (593 kg/ha). The lowest yield was recorded in the state of Odisha (358 kg/ha) followed by C.G. (309 kg/ha).

The overall trend during last three plan period was shown increasing trend in Production and Productivity front but, area is fluctuating in the same period.

(Table-6.4):Plan-Wise StatesøScenario ó Major Stateøs

{Area- lakh ha, Production-Lakh Tones, Yield-kg/ha}

State		X th Plan	% to AI	XI th Plan	% to AI	XII th Plan	% to AI
A.P	A	4.96	15.31	4.55	14.87	3.85	10.95
71.1	P	2.87	20.56	2.72	18.37	3.18	15.26
	Y	579	20.50	598	10.57	826	13.20
Assam	A	0.38	1.17	0.43	1.39	0.53	1.51
7 1334111	P	0.2	1.4	0.24	1.63	0.33	1.59
	Y	526	1.1	558	1.03	624	1.57
Bihar	A	0.25	0.77	0.19	0.62	0.14	0.41
Dillai	P	0.19	1.34	0.16	1.08	0.13	0.61
	Y	760	1.54	842	1.00	889	0.01
Chhattisgarh	A	1.19	3.66	1.09	3.57	1.00	2.86
Cimatingain	P	0.34	2.4	0.32	2.14	0.31	1.49
	Y	286	2.1	294	2.11	309	1.17
Gujarat	A	1.02	3.16	1	3.28	1.03	2.93
Gujurui	P	0.47	3.34	0.65	4.4	0.63	3.04
	Y	461	3.34	650	7,7	615	3.04
Jharkhand	A	0.73	2.25	0.83	2.71	1.09	3.10
Jimikimi	P	0.4	2.84	0.55	3.7	0.95	4.58
	Y	548	2.01	663	3.7	875	1.50
Karnataka	A	1.39	4.29	1.22	4.00	0.92	2.62
Tarriatana	P	0.26	1.83	0.38	2.55	0.38	1.84
	Y	187	1.05	311	2.55	416	1.01
Madhya	A	5.17	15.95	5.22	17.08	8.54	24.32
Pradesh	P	1.78	12.77	1.85	12.50	4.78	22.96
	Y	344	12.77	354	12.50	559	22.70
Maharashtra	A	5.56	17.15	4.26	13.93	3.19	9.08
1120110100	P	2.73	19.53	2.27	15.35	1.51	7.25
	Y	491		533		473	
Odisha	A	1.27	3.93	1.31	4.28	0.92	2.63
	P	0.34	2.4	0.39	2.60	0.33	1.59
	Y	268		298		358	
Rajasthan	A	1.78	5.5	1.56	5.11	2.78	7.92
J	P	0.58	4.18	0.75	5.05	1.46	6.99
	Y	326		481		523	
Tamilnadu	A	2.18	6.73	2.88	9.43	3.54	10.09
	P	0.9	6.47	1.11	7.48	2.59	12.45
	Y	413		385		731	
U.P.	A	5.23	16.14	5.06	16.53	5.88	16.73
	P	2.16	15.46	2.81	19.00	3.05	14.66
	Y	413		555		520	
All India	A	32.39		30.58		35.13	
	P	13.96		14.81		20.82	
	Y	431		484		593	
Sayyaa DES E				2012 12 to 2		0,0	

Source: DES, DAC&FW,ND (*Twelfth plan - Avg. 2012-13 to 2016-17)




Fig.6.2 - Urdbean Area: State & Contribution (XIIth Plan)




Fig.6.3 - Urdbean Production: State & Contribution (XIIth Plan)

6.5.3 District Scenario (2016-17) – Potential Districts

Analysis of the districts within the country revealed that the Krishna district of A.P. contributed with 3.55 % of area and 4.35 % of production. Lalitpur of U.P. ranked second with repect to production (3.59%) and third in production (3.10%). Whereas, Tikamgarh of M.P. ranked second in production 3.63%. Area, production and yield of top ten districts of India is contributing about 21.47 % & 24.11% area and production of the country, presented below

(Table –6.5): Top Potential Districts: Urdbean

{Area-lakh ha, Production-Lakh Tones, Yield-kg/ha}

Name of	State		Area	Pro	duction	Yield	
districts	•	Area	% to India	Prod.	% to India	Yield	YI
Krishna	AP	1.59	3.55	1.23	4.35	775	123
Tikamgarh	MP	1.16	2.59	1.03	3.63	885	140
Lalitpur	MP	1.61	3.59	0.88	3.10	545	86
Chattarpur	MP	1.00	2.22	0.64	2.27	644	102
Bundi	Raj.	0.91	2.03	0.57	2.01	626	99
Villupuram	TN	0.95	2.13	0.53	1.88	558	88
Damoh	MP	0.65	1.46	0.52	1.85	800	127
Guntur	AP	1.00	2.23	0.51	1.81	513	81
Kurnool	AP	0.34	0.76	0.49	1.74	1445	229
Mandsaur	MP	0.40	0.90	0.42	1.49	1052	166
Total above		9.61	21.47	6.83	24.11	710	112
All India		44.78		28.32		632	

Source: State Department of Agriculture




Fig.-6.4: Urdbean: Potential District

(Table-6.6): State-wise yield gap and additional return: urdbean

(Yield: kg/ha; Return: Rs./ha.)

State	Yi	eld (kg/	ha)	Gap ove	er FP	Gap o	ver	Yield	Addition	nal return
		()	,	•		-	SAY		by brid	ging yield
								17	-	(Rs/ha)
	IP	FP	SAY	Actual	%	Actual	%		FP	SAY
Kharif										
Assam	868	701	0	167	24	868	0	-	9018	46872
J&K	576	403	358	173	43	218	61	-	9342	11772
Tamil Nadu	907	724	781	183	25	126	16	781	9882	6804
Karnataka	795	672	386	123	18	409	106	482	6642	22086
M.P.	570	393	475	177	45	95	20	660	9558	5130
MH	668	535	506	133	25	162	32	542	7182	8748
UP	1060	700	536	360	51	524	98	539	19440	28296
West Bengal	1196	927	622	269	29	574	92	692	14526	30996
Manipur	613	466	0.00	147	32	613	_	_	7938	33102
Rajasthan	875	702	474	173	25	401	85	641	9342	21654
Average	813	622	368	191	31	445	121	620	10287	24019
Rabi	010	022	200	1/1	01	110	121	020	10207	21017
AP	1437	1188	831	249	21	606	73	636	13446	32724
Tamil Nadu	969	784	716	185	24	253	35	609	10175	13915
Average	1203	986	774	217	22	429	55	623	12152	24024
Rabi & rice		, , ,				>		020		
Tamil Nadu	927	764	716	163	21	211	29	609	8802	11394
AP	1442	1240	831	202	16	611	74	636	11110	33605
Average	1185	1002	774	183	18	411	53	623	10220	22988

Source-Annual Report- 2016-17, GoI, DPD, Bhopal (Ave. 2013-14 to 2015-16)

State Average Yield - E&S (Ave. 2011-12 to 2015-16)

IP: Improved Practise FP: Farmers Practise SAY: State Avergae Yield

6.6 Urdbean Agronomy

6.6.1 Botanical Description

It is an annual herbaceous plant attaining a height of 30 to 100 cm. The leaves are large, trifoliate and hairy, generally with a purplish tinge. The inflorescence consists of a cluster of 05 to 06 flowers at the top of a long hairy peduncle. The flowers of urd start opening early in the morning and are completely open between 7 to 8 am. Self-fertilization is the general rule. The pods are long and cylindrical being about 4-6 cm in length. There are four to ten seeds in a pod. The seeds are generally black or very dark brown. The split seed of black gram is white in clolour. The germination of seed is of epigeal type.

Black gram is divided into two sub species namely i) *V. mungo var. niger*- matures early, having bold seeded with black colour; ii) *V. mungo var. viridis*- a group of longer maturity duration having small seeded size with green colour.

6.6.2 Production Technology

- Climate: It a crop of tropical region, requires hot and humid climate for best growth. Due to this reason it is grown as summer and rainy season crop in Northern India and in both the main seasons in Eastern and Southern India where winter temperature are quite high. Water logging is fatal for root development and nitrogen fixation during early vegetative stage. Crop is generally grown as rain fed but under assured irrigation during spring in Indo Gangetic Plains of Northern India.
- Soil and Field Preparation: Black gram can be grown on variety of soils ranging from sandy soils to heavy cotton soils. The most ideal soil is a well drained loam with pH of 6.5 to 7.8. Black gram cannot be grown on alkaline and saline soils. Land is prepared like any other kharif season pulse crop. However during summer it requires a thorough preparation to give a pulverized free from stubbles and weeds completely.

• Sowing Time

- **Kharif:** In kharif season sowing is done with the onset of monsoon in later part of June or early part of July.
- Rabi: Second fortnight of October (upland) second fortnight of November (Rice fallow).
- Summer: The sowing could be done from the third week of February to first week of April. Sowing should be done in furrows opened at a distance of 20-25 cm. seed drill could be used for this purpose.

• Recommendation for successful cultivation of Rabi Urdbean:

- Select high yielding varieties resistant to YMV, leaf curl, powdery mildew and drought.
- Use only dual inoculated (Rhizobium+PSB) seeds for better root development and harnessing maximum ÷Nø fixation.
- Treat the seed with Emidacloprid @ 5 ml/kg followed by Macozeb @ 3 g/litre, two days before seed inoculation, as protection against incidence of seedling pest and diseases.
- Use a seed rate of 18-20 kg/ha for upland and 40 kg/ha for rice fallow areas.
- Use basal application of 20 kg N + 50 kg P_2O_5 + 200 kg Gypsum/ha during field preparation, 3-4 cm below and side of the seeds.
- Keep the field free from weeds up to 30 DAS by one hand hoeing.
- In Rice fallow area, Echinochloa (barn yard grass), a major weed can be controlled by mixing Benthiocarb @ 5 L in 50 kg dry sand and applied it as broadcast 3 to 4 days before harvest of paddy.
- For control of Cuscuta spp; post emergence sand mix application of Pendimethalin and flucholarlin at 2.0 L + 1.5 L respectively in 50 kg sand gives best results.
- One irrigation at 35 DAS and 2% spray with urea or DAP at pre-flowering, flowering and pod development is often associated with high jump in grain yield.
- Monitor the crop on field bund cercospora leaf spot (a major problem during rabi in rice fallows) to take effective corrective measures by spraying the crop with mancozeb or copper oxychlorid @ 3 g/liter at 35 and 45 DAS.

• Seed Rate and Spacing

- **Kharif:**During kharif season 12-15 kg seed/ha. The crop should be sown at a distance of 30-45 cm. with 10 cm. plant spacing.
- Rabi:About 18-20 kg seed/ha for upland and 40 kg/ha for Rice fallows with a crop geometry of 30 cm x 15 cm. Higher seed rate in rice fallow is used due to delayed in sowing.
- Summer: About 20-25 kg seed is required per ha. Plant to plant spacing should be kept at 5-8 cm depending upon sowing time and varietal behaviour.

• Seed Treatment

Treat the seed with Thirum (2g) + Carbendazim (1g) or Carbendazim @2.5 g/kg seed to control the soil & seed germinated disease. For sucking pest control seet treatment with Imidacloprid 70 WS @ 7g/kg seed. It is also desirable to treat the seed with Rhizobium & PSB culture (5-7 gm/kg seed).

- Varieties: The variety may be selected as per the adaptability to the region, agro-eco sitiation, sowing time and the cropping system (Table 6 6.7).
- Cropping systems: The important crop rotations with urd are (i) Maize+urd-wheat, (ii) Maize-potato-urd, (iii) Maize-Mustard-urd, (iv) Sorghum+urd-chickpea (Central & South India), (v) Maize-wheat-urd, (vi) Rice-urdbean (Rice fallow of Central & Southern Area), (vii) Paddy-wheat-urd (Summer) North India, (viii) Pigeonpea+Urd-wheat-urd (Summer) North India and, ix) Sugarcane + urdbean (1:2) (Spring) North India.

• Intercropping:

```
Kharif ó Urdbea + Pigeonpea (1:1)
Spring ó Urdbean+ Sugarcane (2:1); Urdbean + Sunflower (2:6)
```

- Water management: In kharif season irrigation not required, if rainfall is normal & if moisture deficit at pod formation stage irrigation should apply. In summer 3-4 irrigation required according to crop requirement. Generally, the crop should get irrigation at an interval of 10-15 days. From flowering to pod development stage, there is need of sufficient moisture in the field.
- Plant nutrient management: For sole crop 15-20 kg/ha Nitrogen, 40-50 kg/ha Phosphorus, 30-40 kg/ha Potash, 20 kg/ha Sulphur is should be applied at the time of last ploughing. However phosphatic and potassic fertilizer should be applied as per soil test value. Fertilizer should be applied by drilling either at the time of sowing or just before sowing in such a way that they are placed about 5-7 cm below the seed. Use of gypsum @ 100 kg/ha would ensures availability of calcium and sulphur at economical rates.
- Weed management: One or two hand weedings should be done up to 40 days of sowing depending upon the weed intensity. Weeds can be controlled by the use of chemicals too use Basalin 1 kg *a.i.* per ha in 800-1000 liters of water as pre-planting spray. It should be well incorporated in the soil before sowing. Application of 100-125g *a.i.*/ha at 0-3 DAS controls wide spectrum of weeds. One or two hand weedings should be done up to 40 days of sowing

depending upon the weed intensity. Weeds can be controlled by the use of chemicals too. Use Pendimethalin 0.75-1.00 kg *a.i.* per ha in 400-600 liters of water as pre-emergence application.

- Harvesting and threshing: Urd should be harvested when most of the pods turn black. Over
 maturity may result in shattering. Harvested crop should be dried on threshing floor for few
 days and then threshed. Threshing can be done either manually or by trampling under the
 feet of bullocks.
- Yield:15-20 quintals of grain per ha.

6.6.3 Plant Protection Measures

A. Diseases

i) Yellow Mosaic Virus

Symptoms: This disease is caused by the mung bean yellow mosaic virus (MYMV) belonging to Gemini group of viruses, which is transmitted by the whitefly (Bemisia tabaci). The tender leaves show yellow mosaic spots, which increase with time leading to complete yellowing. Yellowing leads to less flowering and pod development. Early infection often leads to death of plants.



i) Diseased plants should be rogued out to prevent further spread of the disease; ii) In order to prevent whitefly (Bemisia spp.) infestation spray with triazophos 40 EC @ 2.0 ml/lit, or malathion 50 E

infestation spray with triazophos 40 EC @ 2.0 ml/lit. or malathion 50 EC @ 2.0 ml/lit. or oxydemeton methyl 25 EC @ 2.0 ml/lit. at 10-15 days intervals if required; iii) Grow tolerant/resistant varieties like IPU 94-1 (Uttara), shekhar 3(KU 309), Ujala(OBJ 17), VBN(Bg) 7, Pratap urd 1 etc.



Symptoms: The disease appears on all the part of plants above soil surface. Disease initiates as faint dark spots, which develop into small white powdery spots, coalesceing to form white powdery coating on leaves, stems and pods. At the advance stages, the color of the powdery mass turns dirty white. The disease induces forced maturity of the infected plant causing heavy yield losses and its intensity increases in stress condition.

Control Measures

i) Adopt clean cultivation by destroying diseased plant refuge;ii) Delayed sowing of mungbean and urdbean with wider spacings

considerably reduce the disease severity; iii) Opt for resistant varieties as per recommendation of local agricultural authorities Urdbean: COBG10, LBG 648, 17, Prabha, IPU 02-43, AKU 15 and UG 301); iv) Spray with NSKE @ 50 g/liter of water or neem oil 3000 ppm @ 20 ml/lit. Twice at 10 days interval from initial disease appearance. Spray with eucalyptus leaf extract 10% at initiation of the disease and 10 days later also if necessary; v) Spray with water soluble sulphur 80 WP @ 4 kg/liter or Carbendazin 50 WP @ 1 g/lit.



iii)Leaf Blight

Symptoms: In pre-emergence stage, the fungus causes seed rot and mortality of germinating seedlings. In post emergence stage, seedling blight disease appears due to soil or seed-borne infection. The fungus attacks the stem at ground level, forming localized dark brown patches which coalesce and encircle the stem. Black dot like sclerotia are formed on the surface and below the epidermis on the outer tissue of the stem and root. The pathogen is most favoured at a temperature of 30°C and 15% moisture.

Control Measures

i) Basal application of zinc sulphate @ 25 kg/ha or neem cake
 @ 150 kg/ha or soil application P. fluorescens or T. viride @ 2.5

kg/ha + 50 kg of well decomposed FYM at the time of sowing helps in prevention of the disease; ii) The diseased plants should be uprooted and destroyed so that the sclerotia do not form or survive; iii) Spray with Carbendazim 50 WP @ 1 g/liter of water at an interval of 15 days with the appearance of the symptoms.

B. Insect-Pest Management

i) Aphids

Nature of Damage: Nymphs and adults are seen in large numbers on young plants, leaflets, stem and pods. Young leaves of seedlings become twisted. Excretion of honey dew attracts sooty mold. The adults are black and shiny, upto 2 mm long and some are winged. Nymphs are covered with waxy coating that makes them grey and dull.



Control Measures

i) Spray with 5% crude neem extract or 2% neem oil 3000 ppm;

ii) Spray with Dimethoate 30 EC (1.7 ml/ lit.) or Imidacloprid 17.8 SL @ 0.2 ml/liter of water; iii) Conserve coccinellid beetles, their grubs and Chrysoperla.

ii) Tobaco Caterpillar

Nature of damage: Newly hatched tobacco caterpillar (Spodoptera litura) feed gregariously on the leaf surface for about 2-3 days and leave behind the whitish membranous leaf only. The larvae makes irregular holes on the leaf surface and in severe infestation, they skeletonize the foliage. The maximum damage is caused to the young plants, which are often totally destroyed



i) Collection and destruction of egg masses and newly hatched larvae along with skeletonised leaves can reduce infestation; ii) Spray of microbial pesticides like SINPV [500 LE/ha or or



Bacillus thurengenesis formulations in synchrony with early larval instars is effective against the pest; iii) Spray Malathion 50 EC @ 2.0 ml/lit. or foliar application of Novaluron 10 EC @ 0.75 ml/lit., chitin synthesis inhibitor against eggs of S. litura.

iii) Spotted Pod Borer

Nature of Damage: The larva webs the leaves, inflorescence and feed inside the flowers, flower buds and pods. Eggs are laid on or in the flowers (inserted between the petals). Young larvae feed inside the flowers before moving to developing pods when mid-sized. A larva may consume 4-6 flowers before larval development is completed. Third to fifth instar larvae are capable of boring into the pods and feeding the developing grains. Seeds in damaged pods are totally or partially eaten out by larvae.



Control Measures

i) Spray Bacillus thuringiensis 5 WG @ 1.0 g/liter of water; ii) foliar spray of Profenophos 50 EC @ 2.0 ml/liter of water; iii) Spray of spinosad 45 SC @ 0.2 ml/liter of water is most effective in controlling this pest; iv) Physical shaking of the infested plants over the vessels of oil and water or oily cloth help reduce the population.

iv)Pod Bug

Nature of Damage:Pod bug (Claivgralla gibbosa) adults and nymphs damage leaves, flower buds, stem and pods by sucking cell sap. Major damage is done to the green pods before the maturity of the crop. The attacked pods show pale yellow patches. The grains in the pods become shrivelled and small in size resulting in considerable yield losses.



i) Physical shaking of the infested plants over the vessels of oil and water or oily cloth help reduce the population; ii) Spray



Monocrotophos 36 SL @ 1.0 ml/liter water during flowering and at pod formation stage.

6.7 Recommendation to achieved higher production

- Deep summer ploughing once in 3 years.
- Seed treatment should be done before sowing.
- Application of fertilizer should be based on soil test value.
- In kharif season sowing should be done by ridge & furrow method.
- Yellow mosaic resistant/ tolerant varieties IPU 94-1 (Uttara), shekhar 3(KU 309), Ujala(OBJ 17), VBN(Bg) 7, Pratap urd 1 etc choose as per suitability of region.

(Table-6.7): Recommended urdbean varieties/characteristics

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
Teja (LBG-20)	ANGRAU	1991	Andhra Pradesh	14.0	70-75	Tol. To YMV
Vamban-1	TNAU	1991	Tamil Nadu	8.0	65-70	Tol. To YMV
ADT-4	TNAU	1991	Tamil Nadu	8-9	65-70	Tol. To YMV
ADT-5	TNAU	1991	Tamil Nadu	8.0	65-70	Tol. To YMV, dwarf & erect
Basant Bahar (PDU-1)		1991	All India except South& HillZone	12-13	70-80	Spring, Tolerant to YMV
Prabha (LBG 402)	ANGRAU	1991	Karnataka, Andhra Pradesh, Odisha, T.N.	10.8	78	Rabi, seed bold & dull black
TPU-4	BARC/MAU	1992	MP, Maharashtra & Central part of Rajasthan	7.5	75	Plant erect, medium tall. seed bold & dull black
TAU-2	BARC/PKV	1993	Maharashtra	10.0	70	seed bold & purplish black
Narendra Urd-1 (NDU-88-8)	NDUAT	1993	Uttar Pradesh.	10.0	60-70	Resistant to YMV, Black, medium bold seeded.
LBG-611	ANGRAU	1995	Andhra Pradesh.	14.0	85-90	Resistant to wilt.
WBU-108	BCKV	1996	Punjab, West UP, Rajasthan, Karnataka, A.P. TN).	12	85	Tolerant to YMV, kharif
Mush-338	PAU	1996	Punjab.	9.0	85-90	Tolerant to YMV. seed bold
Mash-414	PAU	1996	Punjab.	9.6	72	Tolerant to root rot. Spring
Birsa Urd-1	BAU	1996	Bihar.	11.0	80	Tolerant to YMV.
Melghat (AKU-4)	PKV	1996	Maharashtra.	10.0	93	Tolerant to stress, for rabi season .
KBG-512	TNAU	1997	Tamilnadu.	7-8	70-75	Tolerant to Stemfly, pods hairy.
Vamban-2	TNAU	1997	Tamilnadu.	12	70	Tolerant to YMV & drought.
KU-301	CSAUAT	1998	TN, Odisha, A.P. & Karnataka	12	70	Res. To YMV, Rabi Season
TU-94-2	BARC	1998	Karnataka, Andhra Pradesh, Odisha, Tamilnadu.	15.0	69	High yielding & YMV resistant early, rabi season
Azad Urd-1 (KU-92-1)	CSAUAT	1999	UP, Bihar, WB.	10.0	80	Spring, Res. To YMV
WBG-26	ANGRAU	1999	Karnataka, A.P. Odisha, TN	10	70	Res. to PM

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
Barkha	RAU,	1999	MP, Maharashtra & Central part	12.0	75	Bold seeded ,Res. to Cercospora leaf
(RBU-38)	Bansawar		of Rajasthan			spot
IPU-94-1 (Uttra)	IIPR	1999	Punjab, Haryana, West UP, North Rajasthan, Gujarat, Bihar, W.B.	11-12	85	Resistant to YMV, kharif season.
Shekhar 2 (KU-300)	CSAUAT	2001	Punjab, Haryana, Delhi, West UP & North Rajasthan)	11-12	70	Resistant to YMV, spring season.
NDU 99-3	NDAUT	2003	NHZ	9.5	85	Res. To YMV, Kharif Season
KU 96-3	CSAUAT	2003	CZ (MP, MS & Gujarat)	8.0	73	Res. To YMV, Kharif Season
Goutam (WBU-105)	Resea.station, Berhanpur	2004	West Bengal	13-15	69-90	Resistant to YMV, Mod. Res. To Cercospora leaf spot
Shekhar 3 (KU 309)	CSAUAT	2004	U.P	10	66-84	Kharif, Resistant to YMV, leaf crinkle, CLS
Mash 1008	PAU	2004	Punjab	12	72	Early, Resistant to MYMV & leaf Crinkle virus
Gujarat urd-1	SDAU	2004	Gujarat	12	late	Late, Moderately resistant to PM & CLS
AKU-15	PDKV	2006	Maharashtra	10-12	65-83	Kharif, Tolerant to PM
Lam 709	ANGRAU	2006	Andhra Pradesh	14	Medium	Tolerant to YMV
Himachal Mash 1	CSK HPKV, HAREC, Dhaulakaun	2007	Low hills of Himachal Pradesh	14-16	75-80	Resistant to leaf crinkle virus, powdery mildew, Anthracnose and YMV
DU-1	UAS, Dharwad	2008		14-15	80-85	High yielded bold seeded and erect growth habit, moderately susceptible to Cercospora leaf spot & powdery mildew
Mash 114	PAU, Ludhiana	2008	Punjab	9-10	83-85	Moderately susceptible to Cercospora leaf & powdery mildew

150 Pulses in India: Retrospect and Prospects -2018 DPD, Govt. of India, Bhopal\

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
Sulata (WBU109)	PORS Beahanpur	2008	UP, Bihar, WB, Assam & Jharkhand	15-16	80-83	Resistant to MYMV, spring season
Pant Urd 31	Central	2008	UP, Tripura, Rajasthan, Odisha,CG, Bihar,AP,Uttrakhand	15	75-80	Resistant to YMV,
Pant Urd 40	Central	2008	Rajasthan, Uttrakhand	14-15	70-75	Short duration variety
Prasad	Central	2008	UP, T N, Odisha	12-14	60-65	Short duration variety,
Mash 114	PAU, Ludhiana	2008	Punjab	9-10	83-85	
VBN (BG)5	Tamil Nadu	2009	TN,	14	60-65	Short duration variety,
Madhra Minumu 207	ARS, Madhira	2009	MS,MP& AP	13	75-80	Tolerant to YMV & stress. Suitable for Kharif, Rabi & Summer
LBG 752	RARS, Lam, Guntur	2009	Andhra Pradesh	18-20	75-80	Photo insensitive, resistant to YMV
CO 6/ COBG 653	TNAU, Coimbatore	2009	TN, AP, Orissa	8-9	60-65	Moderately resistant to YMV
IPU 02-43	IIPR, kanpur	2009	AP, Odisha, Karnataka, Tamil Nadu,Assam	9-11	75	Resistant to MYMV, and PM, kharif season.
KU 99-21	CSAUT, Kanpur	2009	Punjab, Haryana, Western UP and plains of Uttarakhand	10-11	70-75	Kharif season.
Mash 479 (KUG 479)	PAU, Ludhiana	2010	Punjab, Haryana, Western UP and plains of Uttrakhand	12.00	82	Resistant to MYMV and PM spring season
UPU 00-31 (Himachal Mash 1)	CSKHPKV, Palampur	2010	Low hill subtropical zone in kharif season (H.P.)	14-16	75	Resistant to Anthracnose, YMV and Leaf Crinkle and Tolerant to CLS and PM, lister Beetle and Hairy Caterpillar.
LAM Minimum752	ANGRAU	2010	Andhra Pradesh	15	75-82	Resistant to wilt and YMV
VBN 6	NPRC, Vamban	2011	Tamilnadu	8-9	65-70	Resistant to YMV

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
Mash 391 (LU 391)	PAU, Ludhiana	2011	AP, Odisha, Karnataka, Tamil Nadu	8.00	71	Resistant to MYMV, Leaf Crinkle virus, CLS, Anthracnose and PM, spring season
UH 1 (UH 04-06)	CSSHAU	2011	Haryana	11.0	73	Resistant to YMV, kharif season.
VBN (BG) 7 (VBG04-008)	TANU, Coimbatore	2012	AP, Odisha, Karnataka, Tamil Nadu	8.00	63-90	Resistant to MYMV and PM
Vishwas (NUL-7)	Nirmal seeds, pachora (MS)	2012	Chhattisgarh, UP,& Rajasthan		Tolerant to major disease	
Pratap Urd-1 (KPU 07-08)	ARS Kota	2013	Rajasthan	10-11	74-76	It is tolerant to moisture stress, moderately resistant to MYMV, leaf crinkle, anthracnose, bacterial leaf blight
SBC 40	RARS, Shillongini	2014	Assam	15-16	75-80	Resistant to Cercospora leaf spot and YMV
MDUI	AC&RI, Madurai	2014	Tamilnadu	7-8	70-75	Bold seeded
DBGV-5	UAS, Dharwad	2014	Karnataka	14-15	82-85	Moderately susceptible to Cercospora leaf spot and moderately resistant to YMV
Vallabh Urd 1	SBVP UAT, Meerut	2015	Uttar Pradesh	10-11	70-75	Tolerant to MYMV
Indira Urd Pratham	IGKV, Raipur	2016	Chhattisgarh	12-13	70-75	Resistant to powdery mildew upto podding,
Tirupati Minumu- 1 (TBG 104)	ARS, Tirupati	2016	Andhra Pradesh	15-16	75-80	Tolerant to MYMV
PDKV Blackgold (AKU 10-1)	Dr.PDKV, Akola	2016	Maharashtra	10-11	70-75	Tolerant to MYMV and powdery mildew,
LBG 787 (Tulasi)	ARS, Lam	2016	TN , AP, KA, Telangana and Andaman & Nicobar	13-14	70-75	Resistant to Powdery Mildew

Pulses in India: Retrospect and Prospects -2018 DPD , Govt. of India, Bhopal\

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
ADT 6	TNAU, Adhuthurai	2017	Tamilnadu	7-8	65-70	Resistant to leaf crinkle virus, powdery mildew.
KKM-1	SAU,Research institute, Kilikulam(TNAU)	2017	Tamilnadu	6-7	65-70	Erect and determinate plant type with lanceolate leaf, yellow color flower, hairy pods with dull black coloured seeds.

NHZ- North Hilly Zone (H.P.,J.K& U.P.hills),CZ- Central Zone (MP.,Maharashtra, Chhattisgarh, Gujarat), SZ- South Zone (A.P., Karnataka, TN, Odisha) NEPZ-North East plane Zone (East Uttar Pradesh, Bihar, Jharkhand, West Bengal). NWPZ- North West Plane Zone (Punjab, Haryana, Delhi, West UP & North Rajasthan) Res.- Resistant, Tol.= Tolerant, Mod.= Moderately, YMV= Yellow Moasaic Virus, CLS= Cercospora leaf Spot, PM = Powdery Mildew.

Source: - Project Coordinator's Report, AICRP on MULLaRP, ICAR, IIPR, Kanpur. 2017-18

LENTIL (MASUR)





Chapter-7

Lentil (Masur)

Botanical Name : Lens culinaris Medikus subsp. culinaris

: Masur, Malka (bold seeded), lentille (French) linse Synonym

(German), Lenteja (Spanish) and Mercimek (Turkish).

Origin : Turkey to South Iran

: 2n = 14Chromosome

7.1 Economic Importance

It is a valuable human food, mostly consumed as dry seeds (whole decorticated, seed decorticated and split). In Indian sub continent it is mostly consumed as -Daløby removal of outer skin and separation of cotyledons, snacks and soup preparation etc. It is easy to cook and easily digestible with high biological value, hence also referred to patient. Dry leaves, stems, empty and broken pods are used as valuable cattle feed. Bold seeded, attractive shaped grains have high demand for export at premium prices.

7.2 Nutritive Value

Protein	24-26%	Phosphorus	300 mg/100 g
Fat	1.3%	Iron	7mg/100 g
Dietary Fibre	11-12 %	Calcium	69 mg/100g
Carbohydrate	57 ó 60%	Calorific value	343-346 Kcal/100 g
Vitamin C	10-15 mg/100 g	Vitamin A	(450 IU) and Riboflavin

7.3 Agronomic Significance

The crops leaves a reasonable good amounts of atmospheric :Nø in readily available form (upto 30-40 kg/ha) to the succeeding crop. Associated intercrop (other than legume) also gets benefited by Nø transfer from lentil roots up to some extent. It also contributes to sustain production system through physical, chemical and biological improvements of soil properties, as a rotation effect.

It offers good scope in late vacated paddy fields either as *Utera* or succeeding crop as delayed sowing does not affect as adversely as in chickpea and pea due to its high cold tolerant nature. By this reason, this crop is preferred over gram in the regions having cold winters like plains of North and lower Himalayan Hills. It is also a good substitute of chickpea in areas which may be too dry due to shorter duration. The crop is also used as cover crop to check soil erosion in problem areas.

7.4 Growth rate of lentil

From 1985-86 to 1990-91, growth rate observed for APY during both years for APY. however, the maximum growth rate in producton and productivity was recorded with 34% & 23% during 1996-97 (Table 7.1).

(Table-7.1): Growth rate of lentil

(A- Million ha, P- Million tones, Y-kg/ha, Growth Rate (GR)-%)

Year		(Le	entil	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	(== 9
	Area	GR	Prod.	GR	Yield	GR
1980-81	0.93		0.47		498	
1985-86	1.09	16.7	0.66	42.4	607	22.1
1990-91	1.19	8.9	0.85	28.5	717	18.0
1995-96	1.26	5.7	0.71	-16.2	569	-20.7
1996-97	1.37	9.1	0.96	34.8	702	23.6
1997-98	1.29	-5.8	0.80	-16.4	623	-11.3
1998-99	1.39	7.7	0.94	16.7	675	8.3
1999-00	1.46	5.3	1.08	15.0	738	9.3
2000-01	1.48	1.1	0.92	-15.2	619	-16.1
2001-02	1.47	-0.8	0.97	6.5	664	7.3
2002-03	1.38	-6.1	0.87	-10.4	634	-4.6
2003-04	1.40	1.4	1.04	18.9	743	17.2
2004-05	1.47	5.5	0.99	-4.2	675	-9.2
2005-06	1.51	2.2	0.95	-4.8	628	-6.9
2006-07**	1.47	-2.4	0.91	-3.5	621	-1.1
2007-08	1.31	-11.2	0.81	-11.1	622	0.1
2008-09	1.38	5.4	0.95	17.5	693	11.5
2009-10	1.48	7.5	1.03	8.2	697	0.6
2010-11	1.60	7.9	0.94	-8.5	591	-15.2
2011-12	1.56	-2.2	1.06	12.2	678	14.7
2012-13	1.42	-8.9	1.13	7.1	797	17.6
2013-14	1.34	-5.8	1.02	-10.3	759	-4.8
2014-15	1.47	9.5	1.04	1.7	705	-7.1
2015-16	1.28	-13.1	0.98	-5.7	765	8.5
2016-17	1.46	14.0	1.22	24.4	835	9.1
2017-18*	1.55	6.2	1.61	32.0	1034	23.8

Source: DES, Ministry of Agri. &FW (DAC&FW), Govt. of India; 2017-18*- IVth Adv. Est.

(Table-7.2): Per cent share to total pulses

{Area- lakh ha, Production-Lakh Tones, Yield-kg/ha}

Year		Lentil		T	otal Pulse	es	% Sha	re to Total	Pulses
	A	P	Y	A	P	Y	A	P	YI
1980-81	9.35	4.65	498	224.45	106.24	473	4.17	4.38	105
1990-91	11.88	8.51	717	372.55	203.68	547	3.19	4.18	131
1995-96	12.55	7.14	569	331.77	170.10	513	3.78	4.20	111
2000-01	14.78	9.15	619	203.48	110.75	544	7.26	8.26	114
2001-02	14.66	9.74	664	220.08	133.68	607	6.66	7.29	109
2002-03	13.77	8.73	634	204.96	111.25	543	6.72	7.85	117
2003-04	13.96	10.38	743	234.58	149.05	635	5.95	6.96	117
2004-05	14.73	9.94	675	227.63	131.30	577	6.47	7.57	117
2005-06	15.06	9.46	628	223.91	133.84	598	6.73	7.07	105
2006-07	14.70	9.13	621	231.92	141.98	612	6.34	6.43	101
2007-08	13.06	8.12	622	236.33	147.62	625	5.52	5.50	100
2008-09	13.76	9.53	693	220.94	145.66	659	6.23	6.54	105
2009-10	14.80	10.32	697	232.82	146.62	630	6.36	7.04	111
2010-11	15.97	9.44	591	264.02	182.41	691	6.05	5.17	86
2011-12	15.62	10.59	678	244.62	170.89	699	6.39	6.20	97
2012-13	14.23	11.34	797	232.57	183.42	789	6.12	6.18	101
2013-14	13.41	10.17	759	252.18	192.55	764	5.32	5.28	99
2014-15	14.69	10.35	705	235.53	171.52	728	6.24	6.04	97
2015-16	12.76	9.76	765	249.11	163.48	656	5.12	5.97	117
2016-17	14.61	12.24	837	294.47	231.31	785	4.96	5.29	106
2017-18*	15.54	16.07	1034	299.93	252.35	841	5.18	637	123

Source: DES, Ministry of Agri. &FW (DAC&FW), Govt. of India; 2017-18*- IVth Adv. Est.

7.5 Import & Export

Import: The import of pulses in India during April, 2014 to March, 2015 was 8.16 lakh tonnes worth Rs.3418.48 crores against the value of Rs.17196.87 crore for total foodgrains, Rs.121319.02 crore for total agricultural imports and against Rs.2737086.58 crore for total National Import. The import during April, 2015 to March, 2016 was 12.60 lakh tonnes worth Rs.6713.00 crore against the import value of Rs. 26841.87 crore for total foodgrains, Rs.140288.69 crore for total agricultural import and Rs.2490298.08 crore for total National import respectively during this period. The share of Lentil import to Agricultural import was 2.82% and 4.79% respectively during April, 2014 to March, 2015 and April, 2015 to March, 2016.

Export: The pulses export of the country during April,2014 to March, 2015 was 0.080 lakh tonnes worth Rs.49.85 crore against the value of Rs. 59500.54 crore for total foodgrains, Rs.239681.04 crore for total agricultural exports and against Rs.1896445.47 crore for total National export. The export during April, 2015 to March, 2016 was 0.118 lakh tonnes worth

Rs.83.05 crore against the export value of Rs. 42622.29 crore for total foodgrains, Rs.215395.68 crore for total agricultural export and Rs.1716378.05 crore for total National export respectively during this period. The share of Lentil export to Agricultural export was 0.021% and 0.0.039% respectively during April, 2014 to March, 2015 and April, 2015 to March, 2016.

(Table-7.3): Importing & Exporting countries

(Quantity-Thousand Tonnes)

S.	I	MPORT			EXPORT	
No.	Country	Aveg.*	% Share	Country	Aveg.*	% Share
1	Canada	675.60	81.97	Sri Lanka DSR	1.76	23.50
2	USA	82.97	10.07	Bangladesh PR	0.98	13.08
3	Australia	46.26	5.61	Myanmar	0.93	12.37
4	Argentina	18.31	2.22	U Arab EMTS	0.62	8.27
5	Myanmar	0.26	0.03	USA	0.39	5.20
6	Nepal	0.23	0.03	UK	0.39	5.14
7	Turkey	0.16	0.02	Pakistan IR	0.38	5.13
8	Sri Lanka DSR	0.10	0.01	Iraq	0.30	4.04
9	Mozambique	0.08	0.01	Nepal	0.25	3.28
10	Afghanistan TIS	0.07	0.01	Singapore	0.21	2.75
11	Korea RP	0.06	0.01	Malaysia	0.20	2.72
12	Vatican City	0.03	0.003	Australia	0.14	1.86
13	Uzbekistan	0.03	0.003	Turkey	0.10	1.34
14	Kenya	0.03	0.003	Egypt A RP	0.09	1.25
15	Tanzania Rep	0.01	0.002	Kuwait	0.09	1.25
16	Ukraine	0.01	0.001	Bhutan	0.08	1.07
17	Unspecified	0.01	0.001	Saudi Arab	0.07	0.97
18	Pakistan IR	0.005	0.001	Jordan	0.07	0.97
19	Madagascar	0.004	0.001	Netherland	0.07	0.92
20				Others	0.37	4.91
	Total	824.23		Total	7.50	

Source: Ministry of Commerce and Industry; Aveg. *- 2012-13 to 2016-17.

7.6 Production Trends

7.6.1 Global Scenario

India ranked first in the area (18.00 lakh ha) and second in the production (11.00 lakh tonnes) with 39% and 22% of world area and production respectively. The highest productivity is recorded in Croatia (2862 kg/ha) followed by New Zealand (2469kg/ha). Canada rank first in production (41%) due to very high level of productivity (1633 kg/ha) as compared to India (611 kg/ha).

(Table-7.4): Global Ranking: Major Countries

{Area- lakh ha, Production-Lakh Tones, Yield-kg/ha}

Country	Area	%	Country	Prod.	%	Country	Yield
		Cont.			Cont.		
India	18.00	39.79	Canada	19.87	41.16	Croatia	2862
Canada	12.17	26.90	India	11.00	22.79	New Zealand	2469
Turkey	2.43	5.38	Turkey	3.45	7.15	Armenia	2263
Nepal	2.06	4.55	Australia	2.38	4.93	China	2083
Iran	1.68	3.71	Nepal	2.27	4.70	Egypt	2056
Australia	1.62	3.59	Bangladesh	1.57	3.25	Canada	1633
Bangladesh	1.25	2.75	USA	1.56	3.24	Iraq	1566
Syrian Arab	1.11	2.46	Ethiopia	1.37	2.85	USA	1491
Republic							
USA	1.05	2.32	China	1.25	2.59	Australia	1466
Ethiopia	0.99	2.19	Iran	0.85	1.76	Lebanon	1456
World	45.24		World	48.27		World	1067
						India	611

Source: FAO Stat. 2014




Fig.-7.1: Global Area and Production: Lentil

7.6.2 National Scenario: Plan Periods

Plan-wise performance revelaed, the highest area during IXth Plan. However, production & Productivity increased during VIIth plan over the previous plan period. The details plan-wise performance is given below:

(Table-7.5): Plan-wise national scenario

(Area-Mha, Production-MTonnes, Yield-kg/ha)

Plan	Area	%COPP	Prod.	% COPP	Productivity %	6COPP
Sixth Plan (1980-85)	0.96		0.51		527	
Seventh Plan (1985-90)	1.08	12.69	0.68	35.20	632	19.97
Eighth Plan (1992-97)	1.23	13.98	0.80	16.83	648	2.50
Ninth Plan (1997-02)	1.42	14.85	0.94	17.81	665	2.58
Tenth Plan (2002-07)	1.44	1.95	0.95	1.16	660	-0.78
Eleventh Plan (2007-2012)	1.46	1.37	0.96	0.72	655	-0.64
#Twelfth Plan (2012-2017)	1.39	-4.79	1.08	12.5	773	18.01

Source: DES, Twelfth Plan (2012-2017; % COPP ispercentagechange over previous plan.




Fig.-7.2:Plan-wise Scenario: Lentil

7.6.3 States' Scenario: Plan Analysis (Xth -XIIth)

- *Tenth Plan (2002-2007)*: The area under lentil was 14.44 lakh hectares with the total production of 9.53 lakh tonnes. The highest area and production contribution was made by U.P. (41.27% and 48.79%) followed by M.P. (35.04% and 25.50%) and Bihar (11.91% and 14.17%). The highest yield was recorded by the state of Rajasthan (1000 kg/ha) followed by Haryana (833 kg/ha) and Bihar (785 kg/ha). The National yield average was (660 kg/ha). The lowest yield was recorded in the state of C.G. (294 kg/ha) followed by Maharashtra (429 kg/ha) and M.P. (480 kg/ha).
- *Eleventh Plan (2007-2012)*: The country¢s area under Lentil was 14.64 lakh hectares with a production of 9.60 lakh tonnes. The highest area and production contribution was made by U.P. (37.98% and 46.25%) followed by M.P. (37.57% and 24.27%) and Bihar (12.36% and 16.56%). The highest yield was recorded by the state of Rajasthan (893 kg/ha) followed by U.P. (799 kg/ha) and Bihar (878 kg/ha). The National yield average was (656 kg/ha). The lowest yield was observed in the state of C.G. (313 kg/ha) followed by M.P. (424 kg/ha) and Maharashtra (429 kg/ha).
- Twelfth Plan (2012-17): The country area under Lentil was 13.94 lakh hectares with a production of 10.77 lakh tonnes. Madhya Pradesh is on first ranked with respect to acerage 41.01% (5.58 lakh ha) followed by UP 31.65 % and Bihar 11.58% respectively. While in terms of production MP is on first ranked 35.79% (3.86 lakh tonnes) followed by Uttar Pradesh (29.65%) and Bihar (15.98%). The highest yield was recorded by the state of Bihar (1066 kg/ha) followed by Haryana (976 kg/ha) and W.B. (969 kg/ha). The National yield average was (773 kg/ha). The lowest yield was observed in the state of Maharashtra (385 kg/ha) followed by C.G. (389 kg/ha) and M.P. (691 kg/ha).

The overall trend of area, production and yield during the last three plan period shows increasing trend in production and productivity however, area decline during XII plan period is a major concern.

(Table-7.6): Plan-wise statesøscenario ó major states

{Area-lakh ha, Production-Lakh Tones, Yield-kg/ha}

State		X th Plan	% to AI	XI th Plan	% to AI	XII th Plan	
State	A						
	A	0.20	1.39	0.22	1.50	0.28	2.01
Assam	P	0.11	1.15	0.11	1.15	0.19	1.80
	Y	550		500		693	
	Α	1.72	11.91	1.81	12.36	1.61	11.58
Bihar	P	1.35	14.17	1.59	16.56	1.72	15.98
	Y	785		878		1066	
	Α	0.17	1.18	0.16	1.09	0.16	1.16
Chhattisgarh	P	0.05	0.52	0.05	0.52	0.06	0.58
	Y	294		313		389	
	Α	0.06	0.42	0.04	0.27	0.04	0.30
Haryana	P	0.05	0.52	0.03	0.31	0.04	0.38
	Y	833		750		976	
Modhro	A	5.06	35.04	5.5	37.57	5.58	40.01
Madhya Pradesh	P	2.43	25.5	2.33	24.27	3.86	35.79
Tradesii	Y	480		424		691	
	A	0.07	0.48	0.07	0.48	0.01	0.09
Maharashtra	P	0.03	0.31	0.03	0.31	0.01	0.05
	Y	429		429		385	
	A	0.03	0.21	0.01	0.07	0.01	0.04
Punjab	P	0.02	0.21	0.01	0.10	0.00	0.04
	Y	667		1000		667	
	Α	0.19	1.32	0.28	1.91	0.51	3.66
Rajasthan	P	0.19	1.99	0.25	2.6	0.48	4.44
	Y	1000		893		937	
	Α	5.96	41.27	5.56	37.98	4.41	31.65
Uttar Pradesh	P	4.65	48.79	4.44	46.25	3.19	29.65
	Y	780		799		724	
	Α	0.16	1.11	0.15	1.02	0.11	0.80
Uttarakhand	P	0.08	0.84	0.09	0.94	0.09	0.81
	Y	500		600		784	
	A	0.65	4.5	0.55	3.76	0.75	5.37
West Bengal	P	0.45	4.72	0.44	4.58	0.73	6.73
	Y	692		800		969	
	A	14.44		14.64		13.94	
All India	P	9.53		9.6		10.77	
	Y	660		656		773	

Source: DES, DAC&FW,ND (*Twelfth plan - Avg. 2012-13 to 2016-17)




Fig.-7.3: Lentil Area: State Contribution (XIIth Plan)




Fig.-7.4: Lentil Production: State & Contribution (XIIth Plan)

7.6.4 District Scenario (2016-17) – Potential Districts

Analysis of the intra-state status of Lentil crop, is presented in table 7.7. Inter district analysis revealed that district Satna of M.P. with 5.39% of production has the highest share followed by Sagar (5.18%), Narsinghpur (4.71%) and Raisen (4.62%) of M.P. District-wise area, production and yield of top ten district of India in respect of production are presented below which contributed 29.56 *per cent* and 42.53 *per cent* of area and production of the country.

(Table-7.7): Top Potential Districts: Lentil

{Area-Lakh ha, Production-Lakh Tonnes, Yield-kg/ha}

Name of	State		Area	Pr	od.	Yie	ld
District		Area	% to India	Prod.	% to	Yield	ΥI
					India		
Satna	MP	0.54	3.87	0.58	5.39	1075	139
Sagar	MP	0.46	3.32	0.56	5.18	1205	156
Narsinghpur	MP	0.42	3.02	0.51	4.71	1206	156
Raisen	MP	0.30	2.15	0.50	4.62	1656	214
Vidisha	MP	0.39	2.80	0.50	4.61	1272	165
Bahraich	UP	0.51	3.64	0.43	4.03	856	111
Jalaun	UP	0.30	2.16	0.40	3.75	1338	173
Banda	UP	0.48	3.46	0.38	3.49	779	101
Dindori	MP	0.37	2.65	0.37	3.40	989	128
Rajgarh	MP	0.34	2.47	0.36	3.35	1048	136
Total above		4.12	29.56	4.58	42.53	1439	186
All India		13.94		10.77		773	

Source: State Department of Agriculture.




Fig.-7.5: Lentil: Potential District

(Table-7.8): State-wise yield gap and additional return

(Yield: kg/ha; Return: Rs./ha.)

State	Yi	eld (kg/	ha)	Gap	over	Gap	over	Yield	Additio	nal
				FP		SPY		2016-	return	by
								17	bridging	g yield
									gap (Rs.	/ha)
	IP	FP	SAY	Actual	%	Actual	%		FP	SAY
Assam	1013	770	668	243	32	345	52	658	9720	13800
HP	642	396	567	246	62	75	13	619	9840	3000
Jharkhand	670	520	911	150	29	-241	-26	864	6000	-9640
Bihar	1805	1295	1068	510	39	737	69	1005	20400	29480
Chhattisgarh	644	465	378	179	38	266	70	374	7160	10640
M.P.	918	721	594	197	27	324	55	846	7880	12960
Rajasthan	1788	1443	942	345	24	846	90	1003	13800	33840
UP	1369	1112	755	257	23	614	81	761	10280	24560
Tripura	933	712	690	221	31	243	35	667	8840	9720
West	1899	1254	922	645	51	977	106	976	25800	39080
Bengal										
Average	1289	966	777	323	33	512	66	777	12920	20480

Source-Annual Report- 2016-17, GoI, DPD, Bhopal (Ave. 2013-14 to 2015-16)

State Average Yield - E&S (Ave. 2011-12 to 2015-16)

IP: Improved Practise FP: Farmers Practise SAY: State Avergae Yield.

7.7 Lentil Agronomy

7.7.1 Botanical Description

Lentil is an herbaceous annual plant mostly erect and bushy type with four to six primary branches, plant height not exceeding 50-60 cm in general. It has a well developed root system including a central tap root with several lateral branches, spreading in all directions. Root nodules, the site of atmospheric :Nø fixation, are mainly concentrated on primary root. The stem is weak and quadrangular and ends of leaflets some time forms tendrils. Inflorescence is a raceme of two to four flowers. Flowers are small and white with blue, violet or pink tinged. Ovary is short with one or two ovules hence, pods are one to two seeded. Anthesis takes place in buds sometimes before opening of flowers in the next morning. Hence, self pollination is a general rule. Pods are one to one and half cm in length with a curved beak. Grains are often light brown in colour with lens shaped.

Based on the seed size and test weight, it is classified into two main groupsó

i) **Bold seeded:** Includes sub sp. *macro-sperma*, with the test weight of >25 g. also known locally as *Masur* or *Malka Masur* and mainly cultivated in Bundelkhand region of UP/MP and Maharashtra state.

ii) Small seeded: Sub sp. *micro-sperma*, test weight, <25 g, locally known as *masuri* and primarily grown in Indo Gangtic plains of NEPZ (UP, Bihar, West Bengal and Assam).

7.7.2 Production Technology

- Climate: Lentil requires cold climate. It is very hardy and can tolerate frost and severe winter to a great extent. It require cold temperature during its vegetative growth and warm temperature at the time of maturity. The optimum temperature for growth is 18-30°C. Unlike Bengalgram, it can thrive well under stress conditions of frost and winter rains, even at flowering and fruiting stage.
- Soil and Field preparation: Well drained, loam soils with neutral reaction are best for lentil cultivation. Acidic soils are not fit for growing lentil. The soil should be friable and weed free so that seeding could be done at uniform depth. On heavy soils, one deep ploughing followed by two to three cross harrowing should be done. After harrowing, the field should be levelled by giving a gentle slope to ease irrigation. Like gram, it also require good aeration for nodule development, achieved by one deep ploughing followed by one cross harrowing.
- Sowing Time: *Rainfed* Ist fortnight of October in Central and South India and IInd fortnight of October in North India; under irrigated condition first fortnight of November in North India and for **Late sowing** ó First week of December in rice fallows of NEPZ or on fields vacated very late by kharif crops under irrigated condition.
- Seed rate & Spacing: For small seeded: 40-45 kg/ha; Bold seeded: 45-60 kg/ha; Late sown condition: 50-60 kg/ha; Utera cropping: 60-80 kg/ha seed is recommended. Sowing should be done in rows 30 cm. apart and it should be sown at a lower depth (3-4 cm). This could be done either by using a Ferti-seed-drill or by seeding behind desi plough.
- Seed treatment: Fungicide: Thirum (2 gm) + Carbendazim (1gm) or Thirum @ 3 gm or Carbendazim @2.5 g per Kg. of seed; Insecticide: Chlorpyriphos 20E.C. @8 ml./Kg. of seed; Culture: Rhizobium + PSB, one packet each for 10 kg seed.
- Varieties:Based on the region, time of sowing and purpose of cultivation, recommendations etc, selection of variety from Table 6.9.However situation specific varieties for Rice-lentil cropping system ar i) *Utera cultivation* PL-406, PL-639, Arun and, ii) *Late sowings* ó PL-406, PL-639, IPL-15, Narendra Masoor-1.

• Cropping systems

Sequential cropping: The most common rotations under sequential cropping are:

i) Kharif fallow	Lentil (Rainfed areas)	Paddy	- Lentil
iii) Maize	Lentil	Cotton	- Lentil
v) Bajra	Lentil	Jowar	- Lentil

- Intercropping: Most common inter cropping systems are:
- i. Lentil + Sugarcane (Autumn) with two rows of lentil at 30 cm row spacing in between two rows of sugarcane.
- ii. Lentil + Linseed (2:2)
- iii.Lentil + Mustard (2:6)
- Water Management: Most critical stage for moisture stress is pod formation followed by flower initiation. First irrigation should be given at 40-45 days of planting and second at pod filling stage. Most critical stage for moisture stress is pod formation followed by flower initiation. In absence of winter rains and where contribution of soil moisture is negligible *viz*. in Central India, two light irrigations may be applied for significant yield improvement. More irrigation may affect the crop performance adversely.
- Plant nutrient management: Being a legume it does not respond to nitrogen except for some types for initial boosting of growth whereas response to potash is inconsistent due to good :Køsupply status of most of the Indian soils. However, phosphorus definitely plays a vital role in root development, nodulation and growth and yield of the crop. General recommendation is 15-20 kg N and 50-60 kg :Pøas basal placement at soil depth of 10-15 cm during sowing/last ploughing could be met easily through 100 kg DAP/ha. Lentil also respond positively to :Sø(20-40 kg/ha) giving an average nutrient use efficiency of 10-15 kg grain/kg S especially in light textured sandy loam soils of Northern India. SSP is the best source of :Pøfollowed by Gypsum and :Pyriteø
- Among micro-nutrient, Zn is most critical in intensive Rice-Wheat cropping system areas of Punjab, Haryana, Rajasthan (Eastern) U.P. and Bihar General recommendation is 25 kg zinc sulphate as basal, a foliar spray of 0.5% ZnSO4 + 0.25% lime (5 kg zinc sulphate + 2.5 kg lime in 1000 lt. of water per ha). Moø and Feø are the integral components of enzyme introgenousø for Nø fixation. Mo deficiency may create twin deficiency of Nø and Moø Boronø and Moø is found deficient in acidic soil of Eastern India hence 10 kg borax and 1 kg ammonium molybdate as soil application and foliar spray of 2% each of DAP and KCLø at pre flowering and pod development enhance yield by 10-15% along with increasing its ability to resist terminal drought.

• Tips for low input INM

- Application of 2-2.5 tonnes -vermicompostø or 5 t FYM to the -kharifø crop in rotation and seed inoculation with efficient strain of Rhizobia takes care about initial nitrogen requirement and no need to apply +Nø as booster (required especially in low fertile and paddy soils).
- Dual inoculation with Ahizobiumø and PSBø takes care of Nø as well as reduces 25-

- 30% of phosphorus requirement by making available the initial fixed soil #Pø to the plants.
- Rhizobium inoculation is must after paddy as it is aerobic bacteria and most of its population dies during flooding and compaction for want of oxygen.
- In-situ management of rice straw/residues takes care of Zinc and other micronutrient and no need to apply them separately.
- Weed Management: Major weeds are *Chenopodium spp*. (bathua), *Fumaria parviflora* (gajri), *Lathyrus aphaca* (chatri matri), *Vicia sativa* (ankari), *Crisium arvense* (kateli), *Melilotus alba* (senji), *Asphodelus enuifolius* (jungli piaji), *Convolvulus arvensis*, *Phalaris minor* and *Avena ludoriciana*. Orobanche, a parasitic weed is also seen as major problem at some places. Similarly *V sativa* adultrate the grain due to its size, shape and colour. One hand weeding/inter-culture at 30 DAS and another at 55-60 DAS, depending upon the intensity of weed infestation, provides efficient soil oxygen environment to rhizobium bacteria along with soil moisture conservation breaking soil capillaries, creating dust mulch. Application of Metolachlor 1000-1500 g *a.i.*/ha at 0-3 DAS controls many annual grasses and broad leaf weeds.

7.7.3 Plant Protection Measure

A. Disease

i) Seedling Mortality

Symptoms: It is caused by fungi. It appears within a month of sowing when the seedlings start drying up. The drying is mainly two types. (Seedling wilt)- Seedling first turn yellow and dry up. Collar rot- The seedling collapse while still green and then dry out.

Control Measures

i) It can be reduced by delay planting until mid-November; ii) Treat the seed with systemic fungicide Carbendazim @ 2.5 g/kg of seed; iii) Plant resistant varieties like Pant L-406 etc.



ii) Wilt

Symptoms: This is serious disease of lentil in which the growth of the plant is checked, the leaves start yellowing, plant start drying and finally die. The roots of affected plants remain under developed and look light brown in colour.

Control Measures

i) Keep the field clean and follow a three year crop rotation. This will help in reducing the disease incidence; ii) Use tolerant and resistant varieties like Pant Lentil 5, IPL-316, RVL-31, Shekhar Masoor 2, Shekhar Masoor 3 etc; iii) Seed treatment.



iii) Rust

Symptoms: The disease symptoms start as yellowish pustules on the leaflets and pods. Later; light brown pustules appear on both the surfaces of the leaves and other aerial parts of the plant. The pustules finally become dark brown. The plants give dark brown or blackish appearance visible as patches in the field.

Control Measures

i)After harvest, the affected plant trash should be burnt; ii) In NEPZ, normal and early sowing reduces intensity of rust disease; iii) Grow resistant/tolerant varieties like DPL-15,



Narendra Lentil-1, IPL 406, Haryana masur 1, Pant L-6, Pant L-7, LL-931, IPL 316 etc.; iv) Spray the crop with Mancozeb 75 WP@ 0.2 % (2g/liter). 1-2 spray at 50 days after sowing are good for controlling rust.

iv) Stemphylium Blight

Symptoms: The disease causes angular tan leaf lesions; when it is humid (early mornings or after rainfall events), diseased leaves may appear gray due to sporulation by the causal pathogen. Diseased leaves often fall from plants, leaving plants defoliated except for the youngest leaves at the top of the plant. Red lentils are generally more susceptible to the disease than green lentils.



Control Measures

i) After harvest, the affected plant trash should be burnt; ii) Spray the crop with Mancozeb 75 WP@ 0.2~% (2 g/liter). Two spray may be

given at 15 days interval; ii) Grow resistant varieties like Pant L-639, DPL-15, Narendra Lentil-1 etc.

B. Insect-Pest

i) Pod borer

Nature of damage: The caterpillar defoliates the tender leaves and also bores the green pods and feeds upon the ripening grains. It damages almost all the pods in case of severe damage, but causes nearly 25-30% annual yield losses in India.

Control Measures

i)Sprayneem seed extract (5%) @ 50 ml/ liter of water; ii) Spray of Profenphos 50 EC @ 2 ml/ liter or Emammectin benzoate 5 SG @ 0.2 g/liter of water.



ii) Aphids

Nature of Damage: Aphids suck the sap and in case of severe damage the growth is suppressed.

Control Measures

i) Spray of Dimethoate 30 EC @ 1.7 ml/liter or Imidaclopid 17.8 SL @ 0.2 ml / liter of water.



• Harvesting, threshing, storage and yield

Crop become ready for harvest when leaves begin to fall, stem and pod turn brown or straw in colour and seeds are hard and rattle with 15% moisture inside them. Over ripening may lead to fall of pods as well as shattering and seed cracking if seed moisture fall below 10% due to delay in harvesting.

The crop should be allowed to dry for 4-7 days on threshing floor and threshed by manually or bullock/power drawn thresher. The clean seed should be sun dried for 3-4 days to bring their moisture content at 9-10%. The seed should be safely stored in appropriate bins and fumigated to protect them from bruchids.

• Yield:10-15 q/ha.

7.8 Recommendation to achieved higher production

- Deep summer ploughing once in 3 years.
- Seed treatment should be done before sowing.
- Application of fertilizer should be based on soil test value.
- Wilt resistant/ tolerant óRVL-31, IPL81 (Noori), IPL -316, Sekhar masoor-2, Sekhar masoor-2.
 Rust resistant/tolerantóIPL-406, WBL-77, Pant L-6, Pant L-7, Sekhar masoor-2, Sekhar masoor-2, IPL-316.
- Adopt integrated approach for plant protection.
- Weed control should be done at right time.

(Table-7.9): Recommended varieties/characteristics

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
JL 1	JNKVV	1991	MP	8.0	120-125	Early, Tolerant to wilt, Seed bold
Sapana (LH 84-8)	CCSHAU	1991	NWPZ (Punjab, Haryana, Delhi, West UP)	15.0	135-140	Tolerant to Rust & Bold Seeded
VL Masoor 4	VPKAS	1991	Uttrakhand	12.5	168	Tolerant to wilt & Rust, Small seeded &black.
Pant lentil-4 (PL-81-17)		1993	NWPZ (Punjab, Haryana, Delhi, West UP, North Rajasthan)	16.0	140-145	Resistant to Rust & tolerant to wilt.
Lens-4076	IARI	1993	NWPZ (Punjab, Haryana, Delhi, UP) CZ (MP, Maharashtra)	14.0	130-135	Tolerant to wilt & Rust. Seed bold
DPL-15 (Priya)	IARI	1995	NWPZ (Punjab, Haryana, Delhi, West UP)	15-18	130-135	Tolerant to wilt & Rust, bold seeded.
Pusa Vaibhav (L-4147)	IARI	1996	NWPZ (Punjab, Haryana, Delhi, West UP.)	20-24	130-135	Resistant to Rust & Tolerant to wilt, small seeded.
Garima (LH-84-6)	CCSHAU	1996	Haryana.	15-20	135-140	Tolerant to Rust, wilt & Blight.bold seeded.
Narendra Masoor-1	NDAUT	1997	Uttar Pradesh.	14.0	125-130	Resistant to Rust & Tol. to wilt.
DPL-62 (Sheri)	IIPR	1997	NWPZ (Punjab, Haryana, Delhi, West UP.)	17.0	130-135	Resistant to Rust & wilt, bold seeded.
Subrata	BCKV	1998	West bengal	12-18	120-125	Tolerant to Rust, bold seeded.
JL-3	JNKVV	1999	CZ (MP, Maharashtra)	15-19	115-120	Tolerant to wilt, bold seeded .
VL Masoor 103	VPKAS	2000	Uttrakhand	12-14	1645	Tolerant to Rust, small seeded.
Noori (IPL-81)	IIPR	2000	CZ (MP, Maharashtra)	17-18	110-120	Tolerant to Rust, wilt, bold seeded
Pant Lentil-5	GBPUAT	2001	Uttrakhand	15-18	135	Resistant to Rust, bold seeded .
Malaviya Vishwanath (HUL 57)	BHU	2005	Eastern and Central U.P., Bihar, Jharkhand, West Bengal and Assam	14.0	130	Resistant to rust & wilt, small seeded.

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
KLS 218	CSAUAT	2005	NEPZ (East Uttar Pradesh, Bihar, West Bengal).	14-15	125-130	Tolerant to Rust, wilt, small seeded
VL-Masoor-507	VPKAS, Almora	2006	J&K, H.P., Uttrakhand, North Eastern Hills	10-12	140-209	Resistant to wilt
Haryana Masoor-1 (LH-89-48)	CCSHAU	2006	Haryana	14	138	Moderate resistant to all disease
VL Masoor 125	VPKAS, Almora	2006	Uttrakhand	19-20	115-117	Resistant to wilt
VL Masoor 126 (VL-126)	VPKAS, Almora	2007	Uttrakhand, H.P., J&K and North Eastern Hills	12-13	126-212	Resistant to GM and Moderately resistantto wilt and rust
IPL-406 (Angoori)	IIPR	2007	Punjab, Haryana, North Rajasthan, Plains of Uttrakhand and Western UP	17	120-155	Resistant to rust and wilt
Pusa Masoor 5 (L-45994)	IARI	2008	Delhi	17-18	120-128	Resistant to rust moderately resistant to pod borer
Moitree WBL 77	PORS, Berhampore	2009	East UP, Bihar, Jharkhand, Assam & WB	15	117	Resistant to wilt and grey mould
Shekhar Masoor 2 (KLB-303)	Shekhar Masoor 2 (KLB-303)	2009	Uttar Pradesh	14	128	Moderately resistant to wilt and rust
Shekhar Masoor 3 (KLB-320)	Shekhar Masoor 2 (KLB-303)	2009	Uttar Pradesh	14	128	Moderately resistant to wilt and rust
Pant Lentil 7 (PL 024)	GBPUAT	2010	Punjab, Haryana, UP	15	147	Resistant wilt to rust & pod borer
Pant Lentil 8 (PL 063)	GBPUAT	2010	Punjab, Haryana, Plains of Uttrakhand, Western UP, Delhi and Rajasthan	15	135	Mod. Resistant to rust and wilt. Resistant to pod borer

Variety	Source	Release/	Area of adoption	Ave. yield	Days to	Special characteristics
		Notification	(Zone/State)	(q/ha)	maturity	
		Year				
Pant Lentil-6	GBPUAT	2010	Uttrakhand	11	125-145	Resistant to rust. Wilt, Ascochyta
(PL-02)						Blight and Tolerant to pod borer
VL Masoor -129	VPKAS,	2010	Uttrakhand	9.0	151	Resistant to wilt and root rot and no
	Almora					infestation of pod borer
VL Masoor 133	VPKAS,	2011	Uttrakhand	11	150	Resistant to wilt, root rot and rust
(VL133)	Almora					
VL Masoor 514	VPKAS,	2011	Uttrakhand	10	149-159	Moderately resistant to wilt and root rot
(VL514)	Almora					disease. Tolerant to pod borer
LL 931	PAU	2012	Punjab	12-13	146-147	Resistant to lentil rust. Tolerant to pod
						borer

CZ- (MP.,Maharashtra, Chhattisgarh, Gujarat), SZ- (A.P., Karnataka, Tamil nadu, Odisha) NEPZ-North Eastplane Zone (East Uttar Pradesh, Bihar, Jharkhand, West Bengal). NWPZ- North West Plain Zone (Punjab, Haryana, Delhi, West UP & North Rajasthan) Res.= Resistant, Tol.= Tolerant, Mod.=Moderately.

PEAS (MATAR)







Chapter-8

Fieldpeas (Matar)

Botanical Name : Pisum sativum (L.)

Synonym : Matar, Pea

Origin : Mediterranean Region of Southern Europe and Western Asia

Chromosome nos. : 2n = 14

8.1 Economic Importance

Pea is the third most important pulse crop at global level, after dry bean and chickpea and third most popular rabi pulse of India after chickpea and lentil. It provides a variety of vegetarian diet hence liked throughout the world. The mature seeds are used as whole or split into dal and put to use in various ways for human consumption. Beside vegetable purposes, it is also grown as a forage crop for cattle and cover crop to prevent soil erosion but mainly for matured seed for human consumption.

8.2 Nutritive Value

Protein	22-25%	Calcium	64 mg/100g
Fat	0.8- 1%	Iron	4.8 mg/100g
Dietary Fiber-	13.4%	Moisture	11%
Carbohydrate	62.1%		

8.3 Agronomic Significance: Being leguminous crop leaving 25-30kg N/ha to the succeeding crops.

8.4 Growth Rate

Maximum growth for acerage reported during1995-96 (38%) and 2013-14 (25%) and also increasing trend for production and productivity was recorded during 1990-91 with (40% and 27%).

(Table-8.1): Growth rate of fieldpea

(A-Million ha, P-Million tones, Y-kg/ha, Growth Rate (GR)-%)

Year	Fieldpea							
	Area	GR	Prod.	GR	Yield	GR		
1980-81	0.42		0.29		688			
1985-86	0.50	18.6	0.43	47.4	855	24.3		
1990-91	0.55	10.3	0.60	40.9	1092	27.7		
1995-96	0.76	38.0	0.64	5.7	836	-23.4		
1996-97	0.74	-3.8	0.72	13.0	982	17.5		
1997-98	0.75	1.4	0.61	-15.6	818	-16.8		
1998-99	0.84	12.4	0.71	15.8	842	3.0		
1999-00	0.80	-5.0	0.82	15.6	1025	21.8		
2000-01	0.66	-17.2	0.54	-33.7	821	-19.9		

Year			Fie	eldpea		
	Area	GR	Prod.	GR	Yield	GR
2001-02	0.68	2.3	0.61	12.5	903	9.9
2002-03	0.67	-0.5	0.60	-1.3	896	-0.8
2003-04	0.71	6.2	0.73	20.9	1019	13.8
2004-05	0.80	11.7	0.79	8.9	993	-2.6
2005-06	0.78	-1.8	0.72	-9.0	920	-7.3
2006-07**	0.76	-3.0	0.62	-14.3	814	-11.6
2007-08	0.63	-16.9	0.48	-21.6	768	-5.6
2008-09	0.72	13.9	0.66	35.9	916	19.3
2009-10	0.76	6.4	0.67	2.5	883	-3.7
2010-11	0.73	-4.9	0.59	-12.1	816	-7.6
2011-12	0.76	4.3	0.71	19.5	935	14.6
2012-13	0.77	0.8	0.84	18.6	1099	17.6
2013-14	0.96	25.9	0.92	10.0	960	-12.6
2014-15	0.98	1.5	0.89	-3.6	912	-5.0
2015-16	0.90	-7.6	0.74	-16.8	821	-10.0

Source: Agricultural Statistics at a Glance, 2016.DES, Ministry of Agri.&FW(DAC&FW), Govt. of India.

(Table-8.2): Per Cent share to total pulses

{Area- lakh ha, Production-Lakh Tones, Yield-kg/ha}

Year		Peas		Т	Total Pulses			% share to Total Pulses		
	A	P	Y	A	P	Y	A	P	YI	
1980-81	4.23	2.91	688	224.45	106.24	473	0.04	2.74	145	
1990-91	5.54	6.05	1092	372.55	203.68	547	0.06	2.97	200	
1995-96	7.70	6.43	835	331.77	170.10	513	0.08	3.78	163	
2000-01	6.60	5.42	821	203.48	110.75	544	0.07	4.89	151	
2001-02	6.73	6.07	903	220.08	133.68	607	0.07	4.54	149	
2002-03	6.72	6.01	895	204.96	111.25	543	0.07	5.40	165	
2003-04	7.13	7.27	1019	234.58	149.05	635	0.07	4.88	160	
2004-05	7.99	7.92	992	227.63	131.30	577	0.08	6.04	172	
2005-06	7.83	7.20	920	223.91	133.84	598	0.08	5.38	154	
2006-07	7.60	6.18	813	231.92	141.98	612	0.08	4.35	133	
2007-08	6.30	4.84	768	236.33	147.62	625	0.06	3.28	123	
2008-09	7.19	6.59	916	220.94	145.66	659	0.07	4.52	139	
2009-10	7.61	6.75	888	232.82	146.62	630	0.08	4.60	141	
2010-11	7.27	5.93	816	264.02	182.41	691	0.07	3.25	118	
2011-12	7.56	7.06	933	244.62	170.89	699	0.08	4.13	134	
2012-13	7.67	8.43	1099	232.57	183.42	789	0.08	4.60	139	
2013-14	9.63	9.25	960	252.18	192.55	764	0.10	4.80	126	
2014-15	9.75	8.89	912	235.53	171.52	728	0.10	5.18	125	
2015-16	9.03	7.42	821	249.11	163.48	656	0.09	4.54	125	

Source: Agricultural Statistics at a Glance, 2016.DES, Ministry of Agri.&FW(DAC&FW), Govt. of India.

8.5 Import/Export

Import: The import of pulses in India during April, 2014 to March, 2015 was 19.52 lakh tonnes worth Rs. 4970.16 crores against the value of Rs.17196.87 crore for total foodgrains, Rs.121319.02 crore for total agricultural imports and against Rs.2737086.58 crore for total National Import. The import during April, 2015 to March, 2016 was 22.45 lakh tonnes worth Rs.5466.94 crore against the import value of Rs. 26841.87 crore for total foodgrains, Rs.140288.69 crore for total agricultural import and Rs.2490298.08 crore for total National import respectively during this period. The share of Fieldpea import to Agricultural import was 4.10% and 3.90% respectively during April, 2014 to March, 2015 and April, 2015 to March, 2016.

{Dry Peas contributes the single largest share in India's import basket of pulses registering in the total pulses import}.

Export: The pulses export of the country during April, 2014 to March, 2015 was 0.039 lakh tonnes worth Rs.13.63 crore against the value of Rs. 59500.54 crore for total foodgrains, Rs.239681.04 crore for total agricultural exports and against Rs.1896445.47 crore for total National export. The export during April, 2015 to March, 2016 was 0.064 lakh tonnes worth Rs.24.32 crore against the export value of Rs. 42622.29 crore for total foodgrains, Rs.215395.68 crore for total agricultural export and Rs.1716378.05 crore for total National export respectively during this period. The share of Fieldpea export to Agricultural export was 0.006% and 0.011% respectively during April, 2014 to March, 2015 and April, 2015 to March, 2016.

(Table-8.3): Importing & exporting countries

(Quantity-Thousand Tonnes)

S. No.		IMPORT]	EXPORT	
	Country	Aveg.*	% Share	Country	Aveg.*	% Share
1	Canada	1256.96	62.40	Sri Lanka DSR	3.13	80.55
2	Russia	219.60	10.90	Myanmar	0.21	5.39
3	USA	158.37	7.86	Nepal	0.20	5.21
4	France	95.41	4.74	Pakistan IR	0.10	2.66
5	Australia	89.57	4.45	Ukraine	0.06	1.42
6	Lithuania	75.47	3.75	Bangladesh PR	0.05	1.32
7	Ukraine	73.20	3.63	Argentina	0.04	1.13
8	Estonia	12.51	0.62	Bhutan	0.03	0.84
9	Argentina	10.23	0.51	USA	0.03	0.65
10	Romania	8.86	0.44	Maldives	0.02	0.52
11	Germany	4.61	0.23	Kuwait	0.003	0.07
12	Bulgaria	3.24	0.16	South Africa	0.002	0.06
13	Myanmar	1.71	0.08	U Arab EMTS	0.001	0.03
14	Moldova	1.32	0.07	Denmark	0.001	0.03
15	Turkey	0.80	0.04	Singapore	0.001	0.02
16	Others	2.37	0.12	Australia	0.001	0.02
	Total	2014.26		Total	3.88	

Source: Ministry of Commerce and Industry; Aveg. *- 2012-13 to 2016-17

8.6 Production Trends

8.6.1 Global Scenario: Fieldpea

Canada rank first in area (21%) and production (35%) at Global level. China stands second position in area (13.70%) followed by Russian Fed. (12.94 %) respectively. India occupy forth position in area (10.53 %) and 5th position in production (5.36 %). Highest productivity is recorded in Ireland (5000 kg/ha) followed by Netherland (4766 kg/ha), and Denmark (4048 kg/ha). While, Indias productivity is only 822 kg/ha.

(Table-8.4): Global Ranking: Major Countries

{Area-lakh ha, Production-Lakh Tones, Yield-kg/ha}

Country	Area	%Cont.	Country	Prod.	%Cont.	Country	Yield
Canada	14.67	21.16	Canada	34.45	30.80	Ireland	5000
China	9.50	13.70	Russian Fed.	15.03	13.44	Netherlands	4766
Russian Fed.	8.97	12.94	China	13.50	12.07	Denmark	4048
India	7.30	10.53	USA	7.78	6.96	U.Kingdom	4000
Iran	4.75	6.85	India	6.00	5.36	Belgium	3873
USA	3.64	5.25	France	5.39	4.81	Switzerland	3744
Australia	2.45	3.53	Ukraine	3.59	3.21	Germany	3724
Tanzania	2.39	3.45	Ethiopia	3.43	3.06	France	3710
Ethiopia	2.31	3.33	Australia	3.42	3.06	Sweden	3218
Ukraine	1.54	2.21	Iran	2.00	1.79	Armenia	3167
World	69.32		World	111.86		World	1614
						India	822

Source: FAO 2014






Fig.-8.1: Global Area and Production: Fieldpea

8.6.2 National Scenario: Plan Periods

Plan-wise performance revelaed, the highest area & production increased during VIIIth plan followed by XIIth Plan over the previous plan period. The details plan-wise performance is given below:

(Table-8.5): Plan-Wise National Scenario

(Area-Mha, Production-MTonnes, Yield-kg/ha)

			(2)	1. 000 1.1.000, 1 . 0		s, 1000 ng/110/
Plan	Area	%COPP	Prod.	% COPP	Productivity	%COPP
Sixth Plan (1980-85)	0.44		0.32		740	
Seventh Plan (1985-90)	0.47	7.73	0.42	27.95	879	18.76
Eighth Plan (1992-97)	0.71	50.56	0.64	54.90	905	2.88
Ninth Plan (1997-02)	0.74	4.46	0.66	2.06	884	-2.30
Tenth Plan (2002-07)	0.74	0.22	0.69	5.30	929	5.06
Eleventh Plan (2007-2012)	0.72	-3.36	0.62	-9.80	867	-6.67
Twelfth Plan (2012-2017)	0.93	29.1	0.89	43.5	957	10.3

Source: DES, Twelfth Plan (2012-2017); % COPP is percentage Change over previous plan




Fig.-8.2: Fieldpea: Plan-wise Scenario

8.6.3 States' Scenario: Plan Analysis (Xth -XIIth)

- *Tenth Plan (2002-2007):* The area and production during the plan were 7.45lakh ha and 6.92 lakh tonnes respectively. The state of Uttar Pradesh ranked first in area and production (53.75% and 69.25%) followed by Madhya Pradesh (27.93% and 13.92%). Bihar stood in third position with area and production both 3.16% & 3.08% respectively. Rajasthan has recorded highest yield (2190 kg/ha) followed by Haryana (1214 kg/ha) and UP (1197 kg/ha) which is greater than the National productivity (929 kg/ha). Lowest yield was observed in Chhattisgarh (352 kg/ha) followed by Maharashtra (395 kg/ha) and M.P. (463 kg/ha).
- *Eleventh Plan (2007-2012):* During eleventh plan, the area and production were 7.20 lakh hectares and 6.24 lakh tonnes respectively. U.P. stands first in respect of area and production (44.72 % and 62.19 %) followed by M.P. (32.46 % and 15.41%) and Jharkhand (4.28 % & 5.04%). Rajasthan ranked first in yield (1327 kg/ha) followed by U.P. (1205 kg/ha) and Punjab (1192 kg/ha). The lowest yield was observed in C.G. (352 kg/ha) followed by Maharashtra (371 kg/ha) and Madhya Pradesh (412 kg/ha).
- *Twelfth Plan (2012-2017):* The country¢s area under Lentil was 9.34 lakh hectares with a production of 8.88 lakh tonnes. Madhya Pradesh is on first ranked with respect to acerage 39.81% (3.72 lakh ha) followed by UP 37.47% and Jharkhand 4.07% respectively. While in terms of production UP is on first ranked 41.22% (3.86 lakh tonnes) followed by Madhya Pradesh (33.27%) and Jharkhand (5.16%). The highest yield was recorded by the state of Rajsthan (1858 kg/ha) followed by Punjab (1333 kg/ha) and W.B. (1169 kg/ha). The National yield average was (950 kg/ha). The lowest yield was observed in the state of Chhatisgarh (431 kg/ha) followed by Maharshtra (387 kg/ha) and C.G. (431 kg/ha).
- The area, production and yield significantly increased during XIIth plan from previous plan.

(Table-8.6): Plan-Wise StatesøScenario - Major Stateøs

{Area-lakh ha, Production-Lakh Tones, Yield-kg/ha}

State		X th Plan	% to AI	XI th Plan	% to AI	XII th Plan	% to AI
	A	0.22	2.91	0.22	3.03	0.27	2.92
Assam	P	0.13	1.93	0.13	2.14	0.23	2.57
	Y	616		615		835	
	A	0.24	3.16	0.22	3.00	0.18	1.96
Bihar	P	0.21	3.08	0.22	3.47	0.18	2.04
	Y	906		1005		989	
	A	0.16	2.17	0.16	2.20	0.16	1.71
Chhattisgarh	P	0.06	0.82	0.06	0.89	0.07	0.78
	Y	352		352		431	
	A	0.01	0.19	0.01	0.18	0.00	0.03
Haryana	P	0.02	0.24	0.02	0.23	0.00	0.03
	Y	1214		1154		1000	
	A	0.08	1.07	0.31	4.28	0.38	4.07
Jharkhand	P	0.07	0.96	0.31	5.04	0.46	5.16
	Y	838		1019		1205	
Madhya	A	2.08	27.93	2.34	32.46	3.72	39.81
Pradesh	P	0.96	13.92	0.96	15.41	2.95	33.27
Tracesii	Y	463		412		795	
	A	0.17	2.24	0.22	3.06	0.11	1.19
Maharashtra	P	0.07	0.96	0.08	1.32	0.04	0.48
	Y	395		371		387	
	A	0.04	0.51	0.03	0.37	0.02	0.26
Punjab	P	0.04	0.61	0.03	0.50	0.03	0.36
	Y	1105		1192		1333	
	A	0.12	1.62	0.05	0.68	0.11	1.13
Rajasthan	P	0.27	3.84	0.07	1.04	0.20	2.22
	Y	2190		1327		1858	
	A	4.00	53.75	3.22	44.72	3.50	37.47
Uttar Pradesh	P	4.79	69.25	3.88	62.19	3.66	41.22
	Y	1197		1205		1046	
	A	0.04	0.56	0.06	0.84	0.06	0.66
Uttrakhand	P	0.04	0.57	0.03	0.47	0.06	0.68
	Y	952		500		968	
	A	0.13	1.75	0.10	1.44	0.14	1.46
West Bengal	P	0.11	1.60	0.10	1.58	0.16	1.79
_	Y	840		952		1169	
	A	7.45		7.20		9.34	
All India	P	6.92		6.24		8.88	
	Y	929		867		950	

Source: DES, DAC&FW,ND (*Twelfth plan - Avg. 2012-13 to 2016-17)




Fig.-8.3: Fieldpea Area: State@s Contribution (XIIth Plan)




Fig.-8.4:Fieldpea Production: State@s Contribution (XIIth Plan)

8.6.4 District Scenario (2016-17) – Potential Districts

Analysis of the intra-state status of Pea crop, is presented in table 8.7. Inter district analysis revealed that district Jalaun with 15.72% of production has the highest share followed by Lalitpur (12.11%) and Jhansi (10.40%) of U.P. District-wise area, production and yield of top ten district of India in respect of production are presented below which contributed 52.36% and 80.18% of area and production of the country. The yield index of potential districts revealed that the yield of these districts has been above the National average yield (950 kg/ha)in all districts except Mahoba.

(Table-8.7): Top Potential Districts: Fieldpea

{Area-Lakh ha, Production-Lakh Tonnes, Yield-kg/ha}

Sr.	Name of	State		Area		Prod.	Yield	
		State						
No.	District		Area	% to India	Prod.	% to India	Yield	ΥI
I	Jalaun	UP	0.79	8.50	1.40	15.72	1759	185
II	Lalitpur	MP	0.70	7.46	1.08	12.11	1545	163
III	Jhansi	UP	0.66	7.06	0.92	10.40	1400	147
IV	Panna	MP	0.52	5.57	0.87	9.76	1666	175
V	Jabalpur	MP	0.52	5.57	0.81	9.15	1562	164
VI	Datia	MP	0.35	3.75	0.58	6.51	1652	174
VII	Narsinghpur	MP	0.39	4.18	0.48	5.39	1227	129
VIII	Chattarpur	MP	0.25	2.68	0.34	3.79	1347	142
IX	Mahoba	UP	0.42	4.52	0.34	3.78	796	84
X	Dindori	MP	0.28	3.00	0.32	3.57	1132	119
	Total above		4.88	52.26	7.12	80.18	1459	154
	All India		9.34		8.88		950	

Source: State Department of Agriculture




Fig.-8.5: Fieldpea: Potential District

(Table-8.8): State-wise yield gap and additional return

(Yield: kg/ha; Return: Rs./ha.)

State	Yield (kg/ha)		Gap over FP		Gap over SAY		Yield 2016-17	Additional return by bridging yield gap (Rs/ha)		
	IP	FP	SAY	Actual	%	Actual	%		FP	SAY
Assam	1205	960	782	245	26	423	54	782	8575	14805
Bihar	1805	1256	993	549	44	812	82	1053	19215	28420
Chhattisgarh	764	586	419	178	30	345	82	381	6230	12075
J&K	1164	942	831	222	24	333	40	-	7770	11655
Jharkhand	1284	964	1188	320	33	96	8	766	11200	3360
Manipur	784	582	936	202	35	-152	-16	940	7070	-5320
Tripura	1420	1390	850	30	2	570	67	-	1050	19950
UP	1628	1278	1110	350	27	518	47	997	12250	18130
Average	1225	933	904	292	31	321	36	827	10220	11235

Source-Annual Report- 2016-17, GoI, DPD, Bhopal (Ave. 2013-14 to 2015-16)

State Average Yield - E&S (Ave. 2011-12 to 2015-16); IP: Improved Practise FP: Farmers Practise SAY: State Average Yield

8.7 Fieldpea Agronomy

8.7.1 Botanical Description

It is an annual herbaceous, semi erect to erect, succulent plant with a tendency to climb when support is available, grow up to a height of 30-200 cm. Plants bear tap root system with nodules on the surface. Leaves are typically compound, with each leaf has one to three pairs of leaflets and terminal branched tendrils. Inflorescence is axillary raceme. Peas are generally self fertilized but cross pollination may also occur up to some extent. Fruit is a typical pod varying 5-9 cm in length containing 5-10 seeds inside them. Seed germination is hypogeal when cotyledons remain below the ground surface during emergence.

- Two types of peas are generally cultivated all over the world as:
 - **Garden pea:** (*Pisum sativum var. hortense*) also called as table pea, young green seeds are used mostly as vegetables and also for canning purposes. Seeds are bold and wrinkled. Flowers are generally white.
 - Field Pea: (Pisum sativum var. arvense) this group contain ripe, matured seeds and mostly used for dal, some times they are also grown for forage and green manuring purposes. The plants are hardy and grown mostly as rainfed without any irrigation. They are also able to withstand frost. Seeds are round and white, grayish green to grayish yellow. Flowers are coloured.

8.7.2 Production Technology

• Climate: Being a winter season crop it requires a cool growing season with moderate temperature throughout the life. High temperature is more injurious to pea crop than frost. Frost can damage the plants during flowering stage. High humidity associated with cloudy weather results into spread of fungal diseases like damping-off and powdery mildew. Optimum monthly temperature suitable for growth is 13-18°C.

- Soil and Field preparation: A well-drained loamy soils free from excessive soluble salts with neutral pH range of 6.5 to 7.5 is suitable for successful cultivation of the crop. Prepare a level field for even distribution of irrigation water, free from stubbles and crop residues of previous crops by one deep ploughing through disc or mould board plough followed by 2-3 harrowing and planking after each operation. To ensure good drainage and aeration in the field, powdery seed beds must be avoided.
- Sowing time: IInd fortnight of October in north Indian condition and Istweek of October in Central India is the optimum sowing time for rainfed conditions.
- Seed Rate & Spacing: Tall varieties-70-80 kg./ha & 30-45 X 10 cm;Dwarf varieties-100 kg./ha & 22.5 X 10 cm.; 25-30 cm (row to row) and 8-10 cm (plant to plant) for dwarf genotypes like Aparna 30-40 cm (row to row) and 10-12 cm (plant to plant) for tall varieties like Rachna.
- Varieties: Selection of variety as per the adaptability to the region, recommendation, time of sowing, purpose of cultivation and use of inputs etc from Table 8.9.
- Cropping System: In general, peas are sown after harvest of kharif crops. The most common rotations are maize ópea; paddyópeaówheató(being popular in Northern India); cotton ó pea; jowarópea; and bajraópea.
- **Intercropping**: It can be sown as intercrop with autumn sugarcane as two rows of pea at 30 cm row spacing in the centre of two sugarcane rows at 90 cm apart.
- Water management: Fieldpea is mostly grown as rainfed/un-irrigated on residual soil moisture and can sustain drought conditions up to some extent. One or two irrigations at 45 DAs and if needed, at pod filling stage, may be the best recommended irrigation schedule.
- Plant nutrient management: Apply 2.5-5 tones biogas slurry/compost per ha, apply 60 kg P₂O₅ per ha as basal dosein furrow bands for higher P useóefficiency for which single super phosphate (contains 12 % S) to di-ammonoum phosphate should be prefered. On light tectured soils of northern region, application of 0.5 kg molybdenum (1 kg sodium molybdate) per ha has additional effect on yield of fieldpea. Foliar spray of B @ 1-1.5 kg B per ha or soil application of 4 kg borax per ha is recommended on boran deficient soils. Apply 20 kg K₂O per ha. Along with NP is beneficial in K deficient areas. Apply 20 kg sulphur per ha. In acid soils, rhizobium innoculated seed should be treated with 1.5 kg of finaly powdered lime (CaCO₃, 300 mesh).

For correcting Zn deficiency, foliar spray of 0.5 kg ZnSO₄ with 0.25 kg lime or soil application of ZnSO₄ @ 25 kg per ha to one crop on Zn deficient soils is helpful to both the crop of pulse based cropping system.

• Weed management: One weeding 30-45 DAS, depending upon the field conditions. Application of solution MCPB or 2,4 D-B @ 1.2 kg a.i./ha in 500-600 liters of water after 6 weeks of sowing, as post emergence, is effective in sandy loam soils. Application of

Pendimethalin (STOMP) 30 EC @ 1 kg a.i./ha as pre-emergence application can also be used to control the weeds up to 50 days. Application of Metribuzin 250 g a.i./ha at 0-3 DAS or 15-20 DAS.

- Harvesting, threshing and storage: Field peas should be harvested when they are fully ripe and threshed after sufficient drying in the sun. The clean seed should be sun dried for 3-4 days to reduce their moisture content up to 9-10% to be safely stored in appropriate bins. To avoid further development of bruchids and other storage pests, it is recommended to fumigate the storage material before onset of monsoon and again after the monsoon with ALP @ 1-2 tablets per tonne. The small quantity of the produce can also be protected by mixing inert material (soft stone, lime, ash, etc) or by smearing edible/non-edible vegetable oils or by mixing plant products like neem leaf powder at the rate of 1-2% w/w basis.
- Yield: 20-25qtls of grain and straw per ha (irrigated) and 10-15qtls grains per ha (rain fed).

8.7.3 Plant Protection Measures

A. Disease

i) Wilt

Symptoms: The symptoms may be seen in seedling stage. The symptoms are premature yellowing and withering of young leaves during seedling stage and advance stage. Disease caused maximum loss if crop is early sown.

Control Measures

i) Seed Treatment with Thirum (2 gm.) + Carbendazim (1 gm.) /kg of seed; ii) Adopt crop rotation; iii) Avoid early sowing in badly infested areas.



ii) Powdery Mildew

Symptoms: The symptoms first appears on the leaves then on other green parts of the plant. They are characterized by patchy growth on both the surfaces of the leaf and also on the tendrils, pods and stem. In case of severe infestation the plant dies prematurely.

Control Measures

i) Adopt resistant var. like Pant Pea-5, Malviya-15, JP-885, HUP-2 etc.; ii) Spraying with Karathane @ 1 ml/litre or wettable sulphur @ 3 gm/litre or Dinocap @ 1 ml/litre of water and repeat after 10-



15 days, if necessary; iii) Avoid late planting; iv) After harvest collect the plants left in the field and burn them.

iii) Rust

Symptoms: It is caused by fungus. The stem of the plant becomes malformed and the affected plant dies out. All the greenparts of plant are affected. Yellow spots having aecia in round or elongated clusters. Then the uredopustules develop which are powdery and light brown in appearance.

Control Measures

After harvest, the affected plants trash should be burnt; ii) Spray the crop with Mancozeb 75 WP @ 2 g/ liter of water.



B.Insect-Pest

i) Pea Stem fly

Nature of Damage: The maggot of the insect damages the internal tissue, consequently the entire plant dies. The damage is more acute when crop is sown early.

Control Measures

i)Mix 30 kg/ ha Carbofuran (Furadon) 3 % granules or 10 kg /ha Phorate (Thimet) 10 % granules in the soils before sowing the crop; ii) Avoid early planting.



ii) Leaf Miner

Nature of Damage: Larvae of the insect makes tunnel in the leaf causing severe damage. The damage is more during the month of Dec.to Mar.

Control Measures

i) 1 liter of Oxydemeton methyl (Metasystox) 25 EC in 1000 liter of water per hectare when the attack begins and repeat at 15 days intervals.



iii) Pea Aphids

Nature of Damage: They suck the sap of the cells, owing to which the leaves turn pale and yellow. In case of severe infestation the plant growth is checked. Ultimately plant growth get stunted.

Control Measures

i) Spray 1.25 liter of Dimethoate 30 EC or oxydemeton methy (Metasystox) 25 EC in 1000 liter of water per hectare. Reperat the spray after 10-12 days.



iv) Spiny Pod Borer

Symptoms: It is a polyphagous insect. Caterpillar makes hole in pods feed upon developing seed. Late varieties are prone to more damage than earlier one.

Control Measures

i)Picking of green pods should be done 15 days after spraying; ii) Spray of 1.25 liter of cypermethrin in 1000 liter of water per hectare is safe and effective.



8.8 Recommendation to achieved higher production

- Deep summer ploughing once in 3 years.
- Seed treatment should be done before sowing.
- Application of fertilizer should be based on soil test value.
- Weed control should be done at right time.
- Powdery mildew resistant varieties : VL matar -42, VL matar -47, IPF 4-9, Pant P 14, Paras, Prakash (IPFD-1-10), Aman, Gomati (TRCP-8), HFP- 529, HFP-715.
- Late planting should be avoided for preventing powdery mildew.
- Adopt integrated approach for plant protection.

(Table-8.9): Recommended varieties of fieldpeas/characteristics

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
JP-885	JNKVV	1992	CZ (MP, Maharashtra & Gujarat)	21.0	120-140	Resistant to PM.
KFP-103 (Shikha)	CSAUAT	1993	NWPZ (Punjab, Haryana, Delhi, West UP & North Rajasthan)	15-20	130-140	Resistant to PM.
DMR-7 (Alankar)	IARI	1996	NWPZ (Punjab, Haryana, Delhi, West UP & North Rajasthan)	20-25	115-135	Resistant to PM.
Uttra (HFP-8909)	CCSHAU	1996	NWPZ (Punjab, Haryana, Delhi, West UP & North Rajasthan)	20-25	120-140	Resistant to PM., dwarf
Sapna (KPMR- 1441)		1997	Uttar Pradesh.	20-25	120-130	Resistant to PM. Dwarf
Jayanti HFP-8712	CCSHAU	1998	Haryana	20-25	120-140	Res,.To PM., Bold Seeded
Swati (KFPD-24)	CSAUAT	1999	U.P.	25-30	110-125	Resistant to PM. & tolerant to rustDwarf, escapes leaf minor
Malviya Matar-15 (HUDP-15)	BHU	1999	NEPZ (East UP, Bihar, West Bengal). NHZ	25-30	110-130	Resistant to PM., rust and leaf miner
DDR-23 (Pusa Prabhat)	IARI	2000	NEPZ (East UP, Bihar, W.B).	15.0	95-115	Extra early, Resistant to PM
Ambika	IGKV	2000	CZ (MP, Maharashtra & Gujarat)	15-20	100-125	Resistant to PM, Tall Plants
DDR-27 (Pusa Panna)	IARI	2001	NWPZ (Punjab, Haryana, Delhi, West UP & North Rajasthan)	18.0	100-115	Very early, Resistant to PM
Indra (KPMR-400)	CSAUAT	2001	CZ (MP, Maharashtra & Gujarat)	20.0	105-115	Dwarf type, Resistant to PM
Shubhra (IM-9101)	IGKV	2001	Chhattisgarh	15-20	90-95	Resistant to PM
Jay(KPMR-522)	CSAUAT	2001	NWPZ (Punjab, Haryana, Delhi, West UP & North Rajasthan)	23.0	120-140	Dwarf type, Resistant to PM
Adarsh (IPF 99-25)	IIPR	2004	CZ (MP, Maharashtra & Gujarat)	23	110-115	Resistant to Powdery Mildew

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
Vikas (IPFD 99-13)	IIPR	2005	H.P., Maharashtra, C.G., Gujarat & Bundel khand region of U.P.	23	102	Resistant to PM and tolerant to rust
Prakash (IPFD-1-10)	IIPR	2006	M.P., C.G., Maharashtra, Gujarat, Bundel khand region of UP, J&K, H.P. and Uttrakhand	21	94-121	Resistant to PM and tolerant to rust
Paras	IGAU, Raipur	2006	Chhattisgarh	18-24	92-119	Resistant to powdery mildew
Pant P-14	GBPUAT	2006	Uttrakhand	15-22		Resistant to rust and powdery mildew
VL-Matar-42	VPKAS, Almora	2007	Eastern U.P., Bihar, Jharkhand, East Bengal, Assam	20	108-155	Resistant to PM, Moderate resistant to rust
Hariyal (HFP- 9907B)	CCSHAU	2007	Punjab, Haryana, Rajasthan, Delhi, Western U.P.	17-20	128	Resistant to PM & tolerant to rust
Pant Pea -25	GBPUAT	2007	Uttrakhand	18-22	125-128	Resistant to PM &Mod. Resistant to rust
HFP -9426	CCSHAU Hisar	2008	Irrigated areas of Haryana	20	135	Res. To PM and tolerant to root rot. Mod. Resistant to nematodes.
Pant Pea -42	GBPUAT	2008	Western UP, Northern Rajasthan, Punjab, Haryana and plains of Uttrakhand	22	113-149	Resistant to powdery mildew and mod. Resistant to pod borer and stem fly
Swarna Tripti	ICAR,RS, Plandu, Ranchi	2008	Jharkhand, Bihar, & WB.	25	65-70	Resistant to rust and powdery mildew. Tolerant to pod borer
Vivek Matar -10 (VP101)	VPKAS, almora	2008	Uttar Pradesh & Uttrakhand	72-98 (pods)	120-130	Mod. Resistant to PM, white rot, wilt & leaf blight. Less incidence of pod borer
Pant p 13	Central	2008	Western UP,Rajasthan	24-26	110-115	Resistant to powdery mildew

Variety	Source	Release/	Area of adoption	Ave. yield	Days to	Special characteristics
		Notification Year	(Zone/State)	(q/ha)	maturity	
GOMATI (TRCP-8)	ICAR NHE	2010	Uttrakhand Hills, Jammu & Kashmir	22-24	87-97	Suitable for late sown
	Regional		and North Eastern states			condition resistant to PM.
	centre,					Tolerant to pod borer and
	Lembuherra					stem fly
Aman (IPF 5-19)	IIPR	2010	Punjab, Haryana. Plains of Uttrakhand	22	124-137	Res. To PM and tolerant to
			west UP, Delhi and Parts of Rajasthan			rust. Mod. Resistant to pod
						borer and stem fly
IPF 4-9	IIPR, Kanpur	2011	Suitable to irrigated areas	17	129	Resistant to powdery mildew
						and mod. Resistant to pod
						borer and stem fly
VL Matar 47	VPKAS,	2011	Uttrakhand	14.0	142-162	Resistant to wilt, Rust and
(VL47)	Almora					powdery mildew
Dantiwada Field pea	S.D.Agri.	2011	Uttar Pradesh, Bihar, Jharkhand, and	17.0	98-123	Resistant to powdery mildew
1 (SKNP 04-09)	university,		West Bengal.			_ •
		od – Modowotoky DM-				

Res.= Resistant, Tol.= Tolerant, Mod.= Moderately, PM= Powdery Mildew.

MOTHBEAN (MOTH)







Chapter-9

Mothbean

Botanical Name: Vigna acontifolia

Origin : India Synonym : Moth

Chromosome no.: 22

9.1 Economic Importance

- Mothbean (Vigna acontifolia) is a native crop of hot and dry habitats of northern and western parts of India. In severe soil moisture deficit situations, encountered with exceeding evaporative demands, this crop is rated as most economic and useful annual grain legume. This is probably due to genetic buffering embeded in this arid legume to quickly adjust and adapt to the fast fluctuating situations starved due to soil moisture depletion and nutritional deficiency. These very adjusting abilities have rendered this crop as an indispensable component of cropping system prevailing in arid regions.
- Thus, boosting the productivity of this very drought hardy crop in major growing state like Rajasthan, might help in breaking the ceiling of pulse production stagnated in India for last six decades.
- It is most commonly recognized as the potent source of several confectionary items like Papad, Bhujia, namkeen, wada etc. used as daily snaks by the people along with its main use as Đalø
- This crop is used as a source of food, feed, fodder, green manuring and green pasture. Green pods are delicious source of vegetables. Being a pulse, it is a cheap source of vegetable protein for balancing nutritional deficiency. Mostly common on less productive soils on which financially less equipped people having been depending for their livelihood. Mothbean is known for higher proportion of albumin and glutamin fractions of protein along with a good source of lysine and leucine amino acids.

9.2 Agronomic Significance

• Mothbean with deep fast penetrating root system concomitant with drought avoidance capabilities can thrive and survive upto 40-50 days in open fields exhibiting fast depletion of soil moisture and right from seedling emergence, atmospheric temperature heighting to more than 40°C. These adoptive features embodied in mothbean against harsher, harder and unhospital growing situations for unspecified intervals have led this crop to be recognized as arid legume. It also endowed with broad canopy; wing and semi training growth habit also prove useful in keeping the soil moist and lowering soil temperature besides help in reducing the possibilities of soil erosion. Thus, it is a biological means of soil and moisture conservation as temporary in situ shetlter belts.

• These multi adoptive and adjusting natures have scaled mothbean as the only alternative annual crop of sand dunes, requiring no inputs and physical care. This crop is an essential component of sub segments of cropping systems prevalent and common in arid zone like agri-hortic, silvi-pasture, agro forestry, mix cropping, inter cropping and sole cropping, as well. It is, therefore, part of all systems including texturally common poor lands representing the holding of common people, characterized with poor, physical and financial resources. It grows well under uniform rainfall upto 750 mm per annum.

9.3 Nutritive Value

Protein	24-25%	Carbohydrate	61-62%
Fat	1.4-1.5%	Calorific value	330 Kcal/100 g
Dietary Fiber	16%		

9.4 Production Trends

9.4.1 States' Scenario: Plan Analysis (Xth –XIIth)

- *Tenth Plan (2002-2007):* The area coverage and production were 12.42 lakh hectares and 2.97 lakh tonnes respectively, during the tenth plans. Rajasthan ranked first both in area (92.43%) and production (90.24%). Maharashtra stand second in area (3.62%) followed by Gujarat (3.54%), while in production Gujarat stands second with 5.39% followed by Maharashtra 4.04% respectively. The yield was recorded above the National average in the state of Gujarat i.e. (364 kg/ha) followed by Maharashtra (267 kg/ha) remaining state below the National average yield.
- *Eleventh Plan (2007-2012)*: During eleventh plan, the area and production of moth were 14.06 lakh hectares and 4.27 lakh tonnes respectively. Rajasthan occupied first position accounting 94.66% area and 92.82% production share followed by Gujarat (2.65% and 3.79%) and Maharashtra with 1.97% area and 2.25% production share in the country. The yield was observed below the National average in Rajasthan (297 kg/ha) which is major producing state.
- *Twelfth Plan (2012-2016)*: A total of 11.83 lakh hectares and 3.89 lakh tonnes of Moth production was recorded in the country during the twelfth plan period. Area and production of mothbean highest in Rajasthan contributing (86.81% and 82.26%) followed by Madhya Pradesh (10.99% and 14.65%) respectively. However, yield of Madhya Pradesh (438 kg/ha) was higher than the National average yield of (315kg/ha). Need to adopt improved package of practices and varietal break through.

The overall area and production declined during XII plan period from previous plan. Emphasis needed to adopt the improved technology recommendations and varieties to increase the moth production in the country.

(Table-9.1): Plan-Wise StatesøScenario óMajor Stateøs

{Area- lakh ha, Production-Lakh Tones, Yield-kg/ha}

State		X th Plan	% to AI	XI th Plan	% to AI	XII th Plan	% to AI
Gujarat	A	0.44	3.54	0.374	2.659	0.21	1.78
	P	0.16	5.39	0.162	3.798	0.10	2.57
	Y	364		433		467	
Haryana	A	0.05	0.4	0.045	0.319	0.01	0.08
	P	0.01	0.34	0.014	0.323	0.003	0.08
	Y	200		308		286	
Himachal	A			0.009	0.062	0.02	0.17
Pradesh	P			0.012	0.274	0.02	0.51
	Y			1351		1250	
Jammu &	A			0.042	0.296	0.04	0.34
Kashmir	P			0.02	0.465	0.02	0.51
	Y			477		373	
Madhya	A					1.30	10.99
Pradesh	P					0.57	14.65
	Y					438	
Maharashtra	A	0.45	3.62	0.278	1.973		0.00
	P	0.12	4.04	0.096	2.25		0.00
	Y	267		346			
Rajasthan	A	11.48	92.43	13.314	94.661	10.27	86.81
	P	2.68	90.24	3.96	92.819	3.20	82.26
	Y	233		297		311	
All India	A	12.42		14.07		11.83	
	P	2.97		4.27		3.89	
	Y	239		303		315	

Source: DES, DAC&FW,ND (*Twelfth plan - Avg. 2012-13 to 2016-17)




Fig.-9.1: Mothbean Area: State@s Contribution (XIIth Plan)




Fig.-9.2: Mothbean Production: State@s Contribution (XIIth Plan)

9.4.2 District Scenario (2012-13)-Potential Districts

Inter district analysis revealed that the all the potential districts having area and production of the country about 5% and 9% respectively and yield above the National average yield (281 kg/ha) in all the potential districts.

(Table–9.2): Top Potential Districts

{Area-Lakh ha, Production-Lakh Tonnes, Yield-kg/ha}

Sr.	Name of District	State	Aı	rea	Pro	od.	Yie	ld
No.			Area	% to	Prod.	% to	Yield	YI
				India		India		
I	Kutch	Gujarat	0.26	2.92	0.11	4.32	415	148
II	Banas Kantha	Gujarat	0.06	0.67	0.04	1.48	617	220
III	Ahmedabad	Gujarat	0.05	0.52	0.03	1.16	630	225
IV	Patan	Gujarat	0.03	0.37	0.02	0.64	485	173
V	Surendranagar	Gujarat	0.02	0.25	0.01	0.52	591	211
VI	Mehsana	Gujarat	0.01	0.14	0.01	0.32	667	238
VII	Bhavnagar	Gujarat	0.01	0.11	0.01	0.24	600	214
VIII	Rajkot	Gujarat	0.004	0.05	0.003	0.12	750	267
IX	Gandhinagar	Gujarat	0.003	0.03	0.002	0.08	667	238
X	Kheda	Gujarat	0.002	0.02	0.002	0.08	1000	356
	Total above		0.45	5.07	0.22	8.96	496	177
	All India		8.91		2.50		281	

Source: State Department of Agriculture



Fig.-9.3: Mothbean: Potential District

9.5 Mothbean Agronomy

9.5.1 Major Constraint in Production

Besides low productivity, crop is known for plant types of primitive nature, conferring its evolution for survival but not for productive gains. Therefore, treated as neglected crop having marginal and secondary choice.

A. Abiotic Stresses of Mothbean

- i) Mechanical injury: in desert Rajasthan due to hot (>40°c temp.) desicating wind causing removal of epidermis wilting and death.
- ii) Jhola (Hot streaming): When plants are 30-40 days old, 43°c or more temp in concomitant with high wind velocity, in September causes physiological disruption of growth, may lead to plant death.

B. Remedy to over come constraint

Alteration in plant type (Research efforts) which should be high yielding and physiological efficient *i.e.* early partitioning, early maturing and semi erect to erect growth habits along with high Harvest Index, resistance to YMV and Bacterial leaf spot for yield proliferation. Insect pests, particularly Jassids, whiteflies, grubs, and storage pests also deserve special management strategies so that yield losses could be brought at the minimum.

9.5.2 Production Technology

- Climate: It can tolerate high temperature without any adverse effect on flowering and fruit development. Optimum temperature requirement for growth and development is 25-37°c. Bulk of the cultivation is, confined to dry-lands of arid zone with 250-500 mm rainfall requirement with arrangement of proper drainage.
- Field preparation: Preparation of soil aims at to store maximum soil moisture and to reduce subsequent requirement of tillage operations especially when sowing time is limitted. In a good rainfall year, one ploughing with mouldboard plough and a cross harrowing serve the purpose in arid conditions of western Rajasthan. Other alternative is Sweep Cultivation with a ferti seed drill (developed at CAZRI) that can also be used for inter cultivation in wide spaced crop.
- Tillage: Apply emergency tillage for stopping/reducing drafting of surface soil by increasing degree of surface run off that will reduce the surface wind velocity. (Emergency tillage-Making of rough strips on the filed at right angle to the wind direction to temporarily halt the surface movement). Practices for better soil moisture conservation like Dust mulch by sweep cultivator, making staggering trenches with Pitter dicker (CAZRI) and Water harvesting contour bunding soil amendments, soil cultivator & mulching should be followed.
- Sowing Time: With the onset of monsoon. Generally start with first soaking rain to second rain after onset of monsoon. Optimum sowing time IInd to IIIrd week of July. Delay in sowing may result in poor growth, poor germination, increased seedling mortality and incidence of pest and diseases and more conspicuously moisture stress at the flowering, the most critical stage.

- Seed Rate & Spacing: 10-15 kg/ha (short statured, spreading to erect RMO-40 type) for grain and 4-5 kg for mixed crop. For fodder purpose 20-25 kg/ha seed required.; 30-45 cm x 10-20 cm.
- **Seed treatment:** 2 g thiram + 1 g Carbendazim/kg of seed. After fungicide treatment seed inocultion with Rhizobium and PSB culture @ 5-7 g /kg of seed.
- Varieties: Other than the following specific, varieties may be selected from Table 9.3.
 - a) Normal maturity group: (>90 days) Moth Guj. 1 (MG-1), Jadra (IPCMO 943), Jwala (IPCMO-926), IPCMO 880 (26% Protein).
 - b) **Medium maturity group:** (70-90 days) with uniform rainfall throughout season (i) IPCMO 912 (ii) CZM 1 (both 75-80 days duration).
 - c) Early maturity group 60-65 days, higher yield, escape terminal drought especially suitable for lat season, drought areas, resistant to YMV.(i) RMO-40 (62-65 days) (ii) RMO 257 (65 days) (iii) FMM 96 extra early (58-60 days), 5-7 Q/ha short statured + 18-20 Q Fodder (25-30 cm) and non spreading with synchronus maturity (iv) Maru Vardan (RMO 225) (v) Maru Bahar (RMO 435) 15% high yield over RMO 257.

• Cropping system

- Generally grown as single (mono) crop in a year mixed or as a sole crop. However, in a year of good rainfall, it can be rotated with mustard.
- Mixed cropping with pearlmillet, cluster bean, cowpea, mung & sesame in risk prone areas during monsoon. Varieties recommended are RMO 40 & FMM 96 of mothbean and HHB 67 of Bajra.
- Inter cropping (2:1) 2/3 rows of mothbean in between two rows of pearl millet.
- Water Management: It is cultivated in dry land and rainfed condition but in long dry spell one irrigation should be given at pod formation stage.
- Plant nutrient management: Besides their N-fixing capacity they have greater power for absorbing less soluble form of 'P'. Roots have greater CEC hence capable of absorbing divalent catious like Ca++ and Mg++ but can not complete with cereals for monovalant K+. Recommendation is 20-25 tones FYM for improving physical condition and improving water holding capacity of soil along with 10 kg N + 40 kg P₂O₅/ha as basal at the time of sowing or last preparation.
- Weed management: One hand weeding at 30 DAS + pre plant incorporation of fluchloralin (Basalin) @ 0.5 to 1 kg *a.i.*/ha effectively controlled the weeds in mothbean. As pre emergence use Pendimethalin 30 % EC @ 0.75 -1 kg *a.i.*/ha and one hand weeding at 25-30 days after sowing.
- Harvesting and storage: Crop is ready to harvest when pods get mature and turn brown. Plant show drying symptom or yellowing of leaves. Estimated Post harvest losses are 9-10% during threshing transportation, processing and storage. Sun drying, heat treatment, and storage at low temperature with low moisture percentage in seeds (8-9%), is recommended.
- Yield: Fodder 12-25 Q/ha, Grain 3-8 Qtls/ha.

9.5.3: Plant Protection Measures

S. No.	Common Nan	ne Active Period	Incidence	Control Measures
	Sucking Pest	•	•	
i.	Jassids	II week of August	Regular	- Early sowing
		to harvest		- Inter-croping with Pearl Millet (1:4).
ii.	White fly	II week of August	Regular	- Application of Phorate or aldicarb @
		to harvest		1.25 kg a.i. effective upto 4 week.
iii.	Thrips	II week of August	Regular	- Spray with monocrotophos @ 25
	_	to harvest		kga.i./ha or dimethoate @ 0.15 kg
iv.	Aphid & mite	II week of Aug. to I	Sporodic	a.i./ha.
	_	week of Sept.	minor pest	
	Soil/Foliage Pes	t		
v.	White grub	II week of August	Sporodic	Soil application of Phorate or aldicarb @
		to harvest	minor pest	1.25 a.i./ha before sowing.
vi	Termite	Entire cropping	Sporodic	Soil application of Phorate or aldicarb @
		Season	minor pest	1.25 a.i./ha before sowing.
vii.	Root Knot			Use Aldicarb @ 1 kg a.i./ha or
	Nematode			carbofuran @ 2 kg a.i./ha.
	Storage Pest			
viii.	Pulse beetle	During storage	Regular	- Carry Seed moisture level below 10%
	calosobruchus			before storing.
	chinensis			- Fumigation.
				- Mixing/Smearing with neam leaves /cake
				& edible oils.

(Table-9.3): Recommended varieties of mothbean/characteristics

Variety	Source	Year of Release/ Notifica.	Area of adoption Zone/State	Ave. yield (Q/ha)	Days to maturity	Remarks
Maru Bahar	RAU	2002	Rajasthan, Gujarat,	6-6.5	65-67	Early maturing
(RMO-435)			Maharashtra			
CAZRI Moth 2	CAZRI	2003	Rainfed areas	5-7	70-72	
CAZRI Moth 3	CAZRI (Jodhpur)	2004	Rainfed areas	6-5	60-88	Erect, Res. to YMV and dry root rot
RMO-423	ARS, Bikaner	2004	Rajasthan	5-6	67-70	Tolerant to disease insect & pests
RMO-435	Rajasthan	2002		5-6	62-67	Erect, medium nearly, good yield potential, leaves green coloured, mutant from RMO-40
RMO-257	RAO	2007	Rajasthan	6-7	63-65	Semi erect, Tolerant to YMV
TMV (Mb)1	TNAU	2007	T.N.	6	65-70	
RMO-2251 (I)	Rajasthan	2016	Rajasthan, Gujarat	6	60-67	Erect stem with 3-5
						branches, fodder remain
						green upto maturity, average
						incidence of YMV

YMV= Yellow Moasaic Virus

9.6 Recommendation To Achieved Higher Production

- Deep summer ploughing once in 3 years.
- Seed treatment should be done before sowing.
- Application of fertilizer should be based on soil test value.
- Weed control should be done at right time.
- Adopt integrated approach for plant protection.

HORSEGRAM (KULTHI)





Chapter-10

Horse Gram (Kulthi)

Botanical Name : Macrotyloma uniflorum (Lam) Verdc

Synonym: Kulthi

Origin : Peninsular India

Chromosome : 2n = 24

10.1Economic Importance

Horse gram is an important crop of south India. Its grain is used for human consumption as \pm daløas well as in preparation of so called \pm asamøand also as a concentrated feed for cattle. It may also be used as green manure. This crop is generally grown when the cultivator is unable to sow any other crop for want of timely rains and also grown in vacant space of citrus orchard. Horse gram is mainly cultivated in the states of Karnataka, Andhra Pradesh, Odisha, Tamil Nadu, M.P., Chhattisgarh and in foot hills of Uttrakhand and H.P., in India. It is also cultivated in other countries mainly Sri Lanka, Malaysia, West Indies etc.

10.2Nutritive Value

Protein	23-24%	Carbohydrate	59-60%
Fat	2.3-2.4%	Calorific value	321-322 Kcal/100 g
Dietary Fiber	15-16%		

10.3 Production Trends

10.3.1 States' Scenario: Plan Analysis (Xth -XIIth)

- *Tenth Plan (2002-2007):* During the tenth Plan, the total area coverage of Kulthi in the country was 7.16 lakh hectares with a total production of 2.56 lakh tonnes. Karnataka ranked the first both in area and production with 41.65% and 42.48% respectively. Odisha is second in area (11.09%), while, Tamilnadu in production (11.14%). Andhra Pradesh held third position both in area and production (9.74% and 11.25%). The highest yield was recorded in the state of Bihar (804 kg/ha) followed by W.B. (444 kg/ha) and A.P. (413 kg/ha).
- *Eleventh Plan (2007-2012)*: The total area and production during Ninth plan was 3.29 lakh hectares and 1.43 lakh tonnes respectively, Karnataka stands first in respect of area and production with 28.15% and 33.15% respectively. The second position in respect of area and production is occupied by Odisha (18.74% & 12.55%) followed by Chhatisgarh (14.48% & 9.94%). The highest yield was, however, recorded in the state of Bihar followed by W.B. (825 kg/ha) and A.P. (581 kg/ha).
- *Twelfth Plan (2012-2016):* In India, the total area under Horsegram and its production during this plan was 4.36 lakh hectares and 2.04 lakh tonnes respectively. In terms of area and production, Karnataka is on the first position on all India basis contributing 38.53% and 34.22% followed by Tamil Nadu (16.40% & 20.10%). Chhatisgarh ranked third in area (11.15%) and fourth for production (7.84%). Jharkhand is on the third position for production with 8.14% contribution. The highest yield was recorded in the state of Bihar (951 kg/ha) followed by Jharkhand (689 kg/ha) and Tamil Nadu (573 kg/ha).

The trend of area and production during the last three plan period showed significant increased.

(Table-10.1): Plan-Wise StatesøScenario - Major Stateøs

{Area- lakh ha, Production-Lakh Tones, Yield-kg/ha}

State		X th Plan	% to	XI th Plan	% to	XII th Plan	% to AI
			AI		AI		
Andhra Pradesh	A	0.70	9.74	0.09	2.61	0.32	7.39
	P	0.29	11.25	0.05	3.50	0.15	7.45
	Y	413		581		472	
Bihar	A	0.14	1.93	0.11	3.32	0.08	1.86
	P	0.11	4.34	0.10	7.15	0.08	3.77
	Y	804		938		951	
Chhattisgarh	A	0.55	7.68	0.48	14.48	0.49	11.15
	P	0.16	6.41	0.14	9.94	0.16	7.84
	Y	298		298		329	
Jharkhand	A	0.17	2.32	0.16	4.95	0.24	5.53
	P	0.07	2.66	0.09	6.41	0.17	8.14
	Y	410		563		689	
Karnataka	A	2.98	41.65	0.93	28.15	1.68	38.53
	P	1.09	42.48	0.47	33.15	0.70	34.22
	Y	365		512		415	
Madhya Pradesh	A	0.31	4.33	0.23	7.04	0.17	3.99
	P	0.09	3.40	0.07	4.88	0.07	3.43
	Y	281		301		402	
Maharashtra	A	0.65	9.12	0.32	9.65	0.05	1.17
	P	0.20	7.97	0.12	8.48	0.02	0.74
	Y	312		382		294	
Odisha	A	0.79	11.09	0.62	18.74	0.39	8.99
	P	0.20	7.97	0.18	12.55	0.14	6.86
	Y	257		291		357	
Tamil Nadu	A	0.75	10.43	0.20	5.97	0.72	16.40
	P	0.29	11.14	0.08	5.39	0.41	20.10
	Y	382		392		573	
West Bengal	A	0.04	0.50	0.12	3.77	0.03	0.62
	P	0.02	0.63	0.10	7.15	0.01	0.69
	Y	444		825		519	
All India	A	7.16		3.29		4.36	
	P	2.56		1.43		2.04	
	Y	357		433		467	

Source: DES, DAC&FW,ND (*Twelfth plan - Avg. 2012-13 to 2016-17)




Fig.-10.1: Horsegram Area: State@s Contribution (XIIth Plan)




Fig.-10.2: Horsegram Production: State@s Contribution (XIIth Plan)

10.3.2 District Scenario (2016-17) – Potential Districts

Inter district analysis revealed that the major potential district contribute area and production about (35.07% & 35.80%) respectively. Most of the potential district having yield below the National average yield (467 kg/ha) except Simdega district of Jharkhand.

(Table-10.2): Top Potential Districts: Horsegram

{Area-lakh ha, Production-Lakh Tones, Yield-kg/ha}

S. No.	Name of	State	A	rea	Prod	uction	Yie	eld
	District		Area	% to India	Prod.	% to India	Yield	YI
I	Bolangir	Odisha	0.29	6.66	0.13	6.27	440	94
II	Sundargarh	Odisha	0.22	5.12	0.10	4.98	455	97
III	Kalahandi	Odisha	0.22	5.06	0.10	4.85	449	96
IV	Keonjhar	Odisha	0.19	4.28	0.08	4.13	452	97
V	Simdega	Jharkhand	0.07	1.51	0.08	3.88	1200	257
VI	Angul	Odisha	0.13	2.95	0.06	2.80	445	95
VII	Nuapada	Odisha	0.12	2.75	0.05	2.64	449	96
VIII	Kandhamal	Odisha	0.11	2.60	0.05	2.30	415	89
IX	Ganjam	Odisha	0.09	2.06	0.04	1.98	448	96
X	Cuttack	Odisha	0.09	2.09	0.04	1.97	442	95
	Total above		1.53	35.07	0.73	35.80	478	102
	All India		4.36		2.04		467	

Source: State Department of Agriculture




Fig.-10.3: Horsegram: Potential District

10.4 Horsegram Agronomy

10.4.1 Botanical Description: It is an annual herb, slender, with slightly twinning branches, semi-erect, low growing habit 30-50 cm height. Leaves are trifoliate yellowish green to green in colour. Pods are short, 3-5 cm long, linear, with secured beak and 5-7 seeds. Seeds are flattened, 3-6 mm long, light red brown, black or mottled with hard seed coat.

10.4.2 Production Technology

- Climate: Horsegram is extremely drought-resistant crop. Moderately warm, dry climatic conditions are suitable for its optimum growth. It does not grow well on higher altitudes because of cool and wet climate. Horsegram can be cultivated up to an altitude of 1000 m above the sea level. The temperature range of 25-30°C and relative humidity between 50 and 80% is optimum for its growth. Heavy rains during the initial stages of crop growth affect nodule formation owing to poor aeration in the soil. A well-distributed rainfall of about 800 mm is sufficient for its successful cultivation, but it performs well even under low rainfall areas.
- Soil type and Field preparation: Generally grown on lateritic soil (poor in fertility) in south India. The crop can be grown on wide range of soils from light to heavy soils which are free from alkalinity. The crop needs minimum field preparations. Only 1-2 ploughings followed by planking provides desirable seed-bed.
- Sowing time: The main season for sowing horse gram is late August-November. As a fodder crop it is sown during June-August. In Tamil Nadu, it is sown in September-November. In Maharashtra, horse gram is sown as a kharif crop, mixed with bajra or sometimes Niger and also in the Rabi in rice fallows. In M.P. it is a Rabi crop. In northern parts it is grown as **kharif** crop. In West Bengal the sowing period is October-November.
- **Seed Rate & Spacing:** Generally sown as broadcast with 40 kg/ha seed rate for dual purpose i.e. grain and fodder. For line sowing 25-30 kg/ha is enough for grain crop. Row Spacing: 40-45 cm during kharif and 25-30 cm during rabi and about 5 cm plant to plant spacing.
- Seed treatment: Seeds must be treated with seed treating fungicide to reduce infection by fungal pathogens found in the soil. Horse gram seeds are treated with carbendazim (bavistin) 2g for every kg of seeds. Now-a-days bio fungicide like *Trichoderma viridi* is recommended for pulses at the rate of 4g per kg seed. After fungicide treatment seed should be inoculate with Rhizobium and PSB culture @ 5-7 g/kg of seed.
- Selection of varieties: Select a best variety as per the growing season and purpose of cultivation from Table ó 10.3.
- Cropping System: Crop is grown as pure crop as well as mixed crop with sorghum, pearl millet, pigeon pea, sesame or niger.
- Water Management: Grown as rainfed.
- Plant Nutrient Management: 20 kg nitrogen and 30 kg P₂O₅ per ha as basal application at the time of sowing 2-5 cm below and in the side of the seed with the help of ferti.-seed drill is enough for good management of crop.

- Weed Management: Due to luxuriant growth an early weeding/hoeing is enough for weed. Application of Pendimethalin @ 0.75-1 kg a.i./ha as pre emergence application. After that, one hand weeding at 20-25 days after sowing.
- Harvesting, threshing & storage: As usual with other kharif pulses of Vigna group, clean seed should be sun dried for 3-4 days to bring their moisture content at 9-10% to be safely stored in appropriate bins. To avoid further development of bruchids and other storage pests it is recommended to fumigate the storage material before onset of monsoon and again after the monsoon with ALP @ 1-2 tablets per tonne. The small quantity of the produce can also be protected by mixing inert material (soft stone, lime, ash, etc) or by smearing edible/non-edible vegetable oils or by mixing plant products like neem leaf powder at the rate of 1-2% w/w basis.
- Yield: By adopting improved package of practices one can harvest 6-10qtls of grain/ha depending upon the monsoon behavior.

10.4.3 Plant Protection Measures

Insect Pest/Disease/ Causal Organism	Nature of Damage/ Symptoms	Control Measures
i. Aphids	The adults and nymphs suck the juice from the leaves as a result turn brown and crumpled and the plants look sick.	Spray of Oxydemeton methyl 25 @ 1 ml/liter or
ii. Jassids	The adults and nymphs suck the juice from the leaves as a result leaves turn brown an leaf surface become uneven. In severe infection leaves dry up and fall and weaken the plants.	Dimethoate 30 EC @ 1.7 ml/liter water
iii. Pod borer	It is a polyphagous insect. Caterpillar makes hole in pods, sometime also feed seed.	Spray of NPV @ 250 LE/ha. or Quinolphos 25 EC @ 2 ml/liter water
iv. Yellow Mosaic Virus vector-white fly	The symptoms firstly appear on young leaves in the form of yellow, diffused, round spots scattered on the leaf lamina. The infected leaves turn necrotic. The diseased plants usually mature later and bear relatively few flowers and pods. The pods are stunted and mostly remained immature but whenever seeds are form they are small in size.	 i. Grown resistant varieties. ii. Destroy the infected plants. iii. Spray of Oxydemeton methyl 25 @ 2 ml/liter or Dimethoate 30 EC @ 1.7 ml/liter water and repeat after 15 days, if necessary.
v. Root rot	Roots rot and plants show yellowing of the lower-most leaves followed by wilting.	 i. Seed treatment with 2 g Captan or Carbindazim /2 kg of seed. ii. Avoid early sowing in infested areas

10.5 Recommendation To Achieved Higher Production

- Deep summer ploughing once in 3 years.
- Seed treatment should be done before sowing.
- Application of fertilizer should be based on soil test value.
- Weed control should be done at right time.
- Adopt integrated approach for plant protection.

(Table-10.3): Recommended Varieties/Characteristics

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
KS 2	RAU	1991	Rajasthan	6-7	80-85	Early maturing, seed brown
Palem 1	ANGRAU	1998	A.P	10-12	80-85	Early maturing, Semi- spreading
Palem 2	ANGRAU	1998	A.P	8-9	100-105	Med. maturing
Arja Kulthi 21 (AK-21)	MPUAT (Bhilwara)	1998	Rainfed areas of NW parts	8-9	70-105	Early maturing
Paiyur 2	TNAU	2001	SZ (Karnataka, AP, Odisha, TN).	8-9	100-106	For Sept- Oct sowing
PHG 9	UAS	2001	SZ (Karnataka, AP, Odisha, TN).	7-9	100-105	Semi spreading thick foliage
Pratap Kulthi - 1 (AK 42)	MPUAT	2005	Rajasthan, Gujarat, M.P. Haryana	10-12	83-87	Protein 30% lush green foliage with wax deposition
VL Gahat-8	VPKAS, Almora	2007	Uttrakhand	12	92-106	Resistant to anthracnose and stem root
VL Gahat-10	VPKAS, Almora	2007	Uttrakhand	10	110-115	Resistant to YMV & root rot and leaf spot.
GPM 6	AICRP, Bijapur	2008	Karnataka	8-9	120-130	Resistant to YMV, moderately resistant to Rhizoctonia root rot
VL Gahat 15	VPKAS, Almora	2009	Northern India	5-6	95-105	Resistant to Anthrocnose and leaf spot
VL Gahat 19	VPKAS, Almora	2010	North Zone	5	88-94	Multiple disease resistance to important disease
CRIDA 1-18 R	CRIDA, Hyderabad	2009	Karnataka, AP and TN	8	72-102	Tolerant to YMV, powdery mildew, leaf blight, and root rot
Cridalatha (RHG 4)	CRIDA	2010	South Zone	8.0	72-110	Tolerant to YMV, powdery mildew, leaf blight, and root rot & mites
Indira kulthi 1 (IKGH 01-01)	IGKV	2010	Chhattisgarh	7.0	92	Up lands under rainfed co ndition with sowing time of august 15 onwards
Gujarat Dantiwada Horsegarm- 1(GHG-5)	SDAU, SK Nagar	2012	Gujarat, Rajasthan, Uttrakhand, Jharkhand, UP & Maharashtra	5-6	89-100	Resistant to root rot, moderately resistant to PM, Collar rot, Cercopsora leaf spot and leaf blight.


Variety	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
Cridaharsha (CRHG-19)	2014	AP, Karnataka, Kerala and Tamil Nadu	100	900	Tolerant to pod shattering, YMV and Cercospora leaf blight
Pratap Kulthi-2 (AK 53)	2016	Rajasthan,CG, MH & Gujarat & UK	80	650	Extra early maturing, lowest tannin content
Cridavardha n (CRHG-22)	2016	South India	94-100	800-900	Semi compact and semi erect plant type and black seed colour, moderately resistant to anthracnose.
Phule Sakas (SHG-0628- 4)	2016	Maharashtra	110-120	1000- 1100	Resistant to YMv and escape drought due to earliness
Chhattisgarh kulthi-3 (BHG-03)	2017	Chhattisgarh	85-90	500-600	Resistant to collar rot, powdery mildew leaf spot disease.
Chhattisgarh Kulthi-2 (BHW-1)	2017	CG & Neighbouring states	104-112	754	Resistant to powdery mildew & significantly low YMV incidence.

SZ- South Zone (A.P., Karnataka, Tamil nadu, Odisha), NWPZ- North Western Plane Zone (Punjab, Haryana, Delhi, West UP & North Rajasthan)

LATHYRUS (KHESARI)







Chapter-11

Lathyrus (Khesari)

Botanical Name : Lathyrus sativus L.

Synonym : Grass pea, Chickling pea, Teora, Kasari (Bengali) & Kisara (Nepali)

Origin : South Europe and Western Asia

Chromosomes : 2n = 14

11.1Economic Importance

Lathyrus is considered as drought-tolerant hardy crop, and is grown in low-rainfall regions under rainfed conditions, during winter when lentil and chickpea are not expected to give good yields. The crop has unique tolerance ability against stress environmental conditions not only drought but also for water logging. In addition to use as dal and chapatti, it is usually grown as fodder crop. Lathyrus leaves about 36-48 kg/ha nitrogen economy for the succeeding cereal.

11.2 Nutritive Value

Protein	31.9%	Fat	0.9%
Carbohydrate	53.9%	Ash	3.2%

- The major Lathyrus cultivating states in India are Chhattisgarh, Bihar, Jharkhand, Maharashtra, Odisha, Assam, West Bengal and Eastern Uttar Pradesh. In the Rice-Based Cropping system, utilizing the available moisture, it is grown as a relay crop and itos a better option to earn income from rice fields. It is also taken as mixed crop and intercrop during *rabi* and sole crop under outerao conditions.
- Grasspea contains 34% protein and other essential micro-nutrients and may provide nutritional security to the low income people in the society. However the seeds contain Beta-ODAP (-N-oxalyl-L-, -diaminopropionic acid), a toxin known to cause neuro-lathyrism, if consumed as staple food for as long period of 4 to 5 months continuation. In view of this, a ban on the sale of its produce was imposed in some states but its cultivation was not under ban.
- Chhatisgarh (Raipur, Durg, Ranjandgaon, Kabirdham, Bilaspur, Dhamtari, Raigarh, Mahasamund, Janjgir-Champa and Jashpur) and its adjoining areas of Vidarbha region of Maharashtra and MP are the major areas of its cultivation and consumption. In Chhattisgarh, its major cultivation is mainly under õUteraö system, where the seeds of grass pea are broadcasted on the standing water in the paddy field about 10-15 days before harvest.

11.3 Background: Policy on Consumption/Sale of Lathyrus

Khesari Dal (*Lathyrus sativus*) has been a subject of controversy among the agricultural scientists, nutrition experts and the farming community in the country for many decades. Though, admittedly a high protein pulse, its sale was banned by the Government as early as in 1961, under the Prevention of food Adulteration Act, 1954, on the ground that its consumption was harmful to health. The controversy arose from the conclusions of certain studies conducted in the past that it contains a toxic element called BOAA (B-N-oxalyl-

aminoalanine), now ODAP, which causes a crippling affliction of the central nervous system called Lathyrism.

11.3.1 High power committee on lathyrus

- The States of Chhattisgarh, Jharkhand, West Bengal, M.P. and Odisha, did not have any ban on cultivation, sale and consumption of lathyrus during 2007-08. However, this was banned in Maharashtra state.
- On consistant request from the Maharashtra based NGO, ANI, (Nagpur) for lifting of ban on sale in their state, the Parliamentary Standing Committee on Agriculture constituted a High Power Committeeø under the Chairmanship of Secretary DARE ó cum-DG, ICAR, with the approval of Agriculture Minister. The other Members of the Committee were the Secretaries of Agriculture and Health, Government of India; State Agriculture Secretaries of Government of M.P., MS, Odisha and West Bengal; Directors from NIN, Hyderabad; IARI New Delhi; ITRC, Lucknow; Pulses Research Institute (Now IIPR, Kanpur); Directors of Research BCKVV, West Bengal; OUAT, Odisha; IGKVV, M.P.; DG Maharashtra Council of Agriculture Research, Pune; DDG, Crop Sciences (ICAR); Director ó DG ó ICMR, New Delhi including Dr. S.L. Kothari, the president of the Maharashtra based NGO, ANI, Nagpur.
- The High Powered Committee, in its recommendation revealed that continued consumption of khesari (*Lathyris sativus*) which contains BOAA, causes lathyrism. The Committee, however, did not give any õthreshold levelsö to precisely establish to determine in a specific quantitative terms and inter alia recommended further research/data generation to determine the threshold doses from where the real risk starts. The Parliamentary Standing Committeeø on Agriculture unanimously adopted the recommendation report of the High Powered Committee in its sitting dated 8th November, 2001.
- Accordingly, the Ministry of Agriculture on the recommendation of this Committee and also of the Ministry of Health & Family Welfare, had funded a study to NIN, Hyderabad on project proposal on Experimental Neuro lathyrism in goats/sheepø to determine the threshold doses of consumption of khesari dal for Rs.11,89,400/- for a period of two years (2002-03 to 2003-04).
- **A. Brain storming session on lathyrus** dated 20.12.2005 was convened under the Chairmanship of Mrs. Radha Singh, the then Secretary (A&C) specifically for two reasons:
- i) Academy of Nutrition Improvement, Nagpur, {Soyamilk Complex, Sitabuldi, Wardha Road, Nagpuró440 012 (MS)}, an NGO has been making complaints/putting the case before the Department of Agriculture & Cooperation for lifting of the ban on sale of lakh/lakhodi dal imposed in 1961 by the State Government of Maharashtra in pursuance to Ministry of Health & Family Welfare circular under PFA rule, 1955 on the ground that its consumption is associated with the disease õLathyrismö causing õCrippling paralysisö due to presence of Beta-n-oxalyl-aminoalanine (BOAA) content, now termed as ODAP.

- ii) Rule 44-A-Sale of khesari grain prohibited:- The Ministry of Health & Family Welfare says õNo person in any state shall with effect from such date as the State Government concerned may, by the Notification in official Gazette specify in this behalf, sell or offer or expose for sale, or have in his possession for the purpose of sale, under any description or for use as an ingredient in the preparation of any article of food intended for sale of khesari grain and its mixtureö. The concerned State Governments, based on the consumption behavior vis-a-vis incidence of lathyrism causing crippling analysis, in consultation with the State Health & Family Welfare Department promulgated the advice of the Ministry of H&FW under the provision of the PFA 1954.
- iii) "Project on Enhancing Grass pea production for safe human food, animal feed and sustainable rice-based production systems in India funded under NFSM Monitoring Report regarding".
- iv) The promotion of this crop and its cultivation has not been covered under the NFSM-Pulses, A3P and 60000 Pulse Village Programme up till 2015-16. However, the DAC-ICARDA collaborative project was funded during the 11th Plan (last two years 2010-11 to 2011-12) with an out lay of Rs. 362.03 lakh. The pilot states in the first Phase were U.P (Jhansi, Lalitpur, hamirpur, Mirzapur, Chandauli); Chhattisgarh (Raipur, Durg, Bilaspur), Bihar (Patna, Nalanda) and West Bengal (Coochbehar /Nadia). Initially for two years (2010-11 to 2011-12), the project continued during 2012-13.
- v) The other cooperating centres were Indian Grassland and Fodder Research Institute, Jhansi, IIPR, Kanpur, IGKV, Raipur, Society for Promotion of Agricultural Research & Knowledge (SPARK), Patna (Bihar), Uttar Banga Krishi Viswavidyalaya, Cooch Behar (West Bengal) Bidhan Chandra Krishi Vishwavidyalaya, Kalyani (West Bengal) and Pulses & Oilseeds Research Station Berhampore, Murshidabad (West Bengal).

vi) The Objectives of the project were:

- Enhancing fodder and straw yields through introduction of high-biomass and low toxin
 grass pea varieties to support nutritional feed & fodder where only paddy straw is
 available as cattle feed.
- Replacement of indigenous high toxin grass pea varieties available with farmers with low toxin & high biomass varieties through farmers participatory approach.
- Identification of new grass pea varieties through adaptive research, multi-locational testing by farmers participatory selection.
- Developing strong seed production and distribution system of quality dual purpose seeds of farmers- preferred varieties along with matching production technologies.
- Capacity building of farmers, extension personnel etc. for farmer-participatory adaptive
 research and technology transfer for adoption and expansion of improved production
 technologies, quality seed production through training, visits, workshops and seminars
 etc.
- Back-up research (farmers participatory) for further identification of grass pea varieties and refinement of production technologies.

B. Lathyrus under Development Programmes

- Assam, West Bengal has taken this cultivation of Lathyrus under NFSM during 2016-17 (Rabi). The state of C.G. has also taken the development programme on lathyrus under RKVY.
- Varieties like Nirmal, Prateek and Mahateora having less ODAP content may be proposed.

11.4 Production Trends

11.4.1 States' Scenario: Plan Analysis (Xth -XIIth)

- TenthPlan (2002-2007): The total area and production of Khesari was 6.35 lakh hectares and 3.76 lakh tonnes respectively. Out of these, Chhattisgarh ranked first both inarea and production (64.89% and 54.32%) followed by Bihar (18.47% and 25.69%). Madhya Pradesh is in third position for acreage with 7.09%, while W.B. ranked third in production (8.62%). Due to highest yield (960 kg/ha) among the lathyrus producing states. The major contributing state of Chhattisgarh recorded yield below (496 kg/ha) the National average yield (592 kg/ha).
- Eleventh Plan (2007-2012): The total area and production of lathyrus were 5.16 lakh hectares and 3.42 lakh tonnes. C.G. stands first in respect of area and production (65.64 % and 58.38%) followed by Bihar (16.83% and 24.53%). And M.P. (9.10% & 9 %). The highest yield was recorded in the state of Bihar (965 kg/ha) followed by West Bengal (778 kg/ha) and M.P. (654 kg/ha). However, major contributing state i.e. C.G (589 kg/ha) was observed below the National average yield (662 kg/ha).
- Twelfth Plan (2012-2016): The total area and production of Khesari were recorded at 4.58 lakh hectares and 3.84 lakh tonnes respectively. Chattisgarh ranked the first position both in area and production (76.06 % and 66.38%), followed by Bihar (14.04% and 17.21%) and West Bengal (9.28% and 13.93%) respectively.

(Table-11.1): Plan-Wise StatesøScenario

{Area- lakh ha, Production-Lakh Tones, Yield-kg/ha}

State		X th Plan	% to AI	XI th Plan	% to AI	XII th Plan	% to AI
Bihar	A	1.17	18.47	0.87	16.83	0.64	14.04
	P	0.97	25.69	0.84	24.53	0.66	17.21
	Y	824		965		1028	
	A	4.12	64.89	3.39	65.64	3.35	73.06
Chhattisgarh	P	2.04	54.32	1.99	58.38	2.55	66.38
	Y	496		589		762	
Madhya	Α	0.45	7.09	0.47	9.10	0.09	1.90
Pradesh	P	0.31	8.35	0.31	9.00	0.07	1.88
	Y	697		654		828	
Maharashtra	A	0.27	4.24	0.15	2.97	0.08	1.77
	P	0.11	3.03	0.06	1.69	0.02	0.52
	Y	423		376		247	
West Bangal	A	0.34	5.32	0.28	5.46	0.43	9.28
	P	0.32	8.62	0.22	6.41	0.54	13.93
	Y	960		778		1259	
All India	A	6.35		5.16		4.58	
	P	3.76		3.42		3.84	
	Y	592		662		837	

Source: DES, DAC&FW,ND (*Twelfth plan - Avg. 2012-13 to 2016-17)




Fig.-11.1: Lathyrus Area: State® Contribution (XIIth Plan)




Fig.-11.2: Lathyrus Production: State@s Contribution (XIIth Plan)

11.4.2 District Scenario (2016-17) – Potential Districts

Inter District analysis revealed that major top ten potential district contributing the area and production about 60.84% and 53.86% respectively. All the Potential districts comes under C.G. and one from West Bengal and Bihar.Most of the districts average yield below the National average yield (837 kg/ha) except Purba Medinipur (1437 kg/ha) and Patna (1368 kg/ha) districts of W.B and Bihar.

(Table-11.2): Top Potential Districts: Lathyrus

{Area-lakh ha, Production-Lakh Tones, Yield-kg/ha}

Sr.	Name of District	State	Area Prod.		od.	Yie	ld	
No.			Area	% to India	Prod.	% to India	Yield	YI
		~~	0.10		0.10			
I	Mungeli	CG	0.62	13.47	0.48	12.55	779	93
II	Bilaspur	CG	0.27	5.94	0.23	5.92	834	99
III	Balod	CG	0.44	9.58	0.23	5.90	515	62
IV	Purba Medinipur	WB	0.15	3.18	0.21	5.47	1437	172
V	Baloda bazar	CG	0.31	6.72	0.19	5.06	629	75
VI	Rajnandgaon	CG	0.30	6.55	0.16	4.26	544	65
VII	Dhamtari	CG	0.23	4.93	0.16	4.08	693	83
VIII	Patna	Bihar	0.11	2.32	0.15	3.80	1368	163
IX	Raipur	CG	0.23	4.95	0.14	3.72	628	75
X	Janjgir	CG	0.15	3.20	0.12	3.11	813	97
	Total above		2.79	60.84	2.06	53.86	740	88
	All India		4.58		3.83		837	

Source: State Department of Agriculture.




Fig.-11.3: Lathyrus: Potential District

11.5 Lathyrus Agronomy

11.5.1 Botanical Description

Plant of lathyrus is herbaceous annual with slender, glabrous, well branched, winged procumbent stems. Pods are flattened, oblong, up to 4 cm long; two winged dorsally, up to five seeded. Seeds are wedge shaped, angular, white or brown sometimes mottled. Germination is hypogeal.

11.5.2 Production Technology

- Climate: Being a winter season crop it prefers temperate climate with good adoption under climatic extremities.
- Soil and Field preparation: Thrive well in all types of soils except of very acidic nature. It prefers heavy soils belonging to low lying areas which are not suited to other crops. It grows abundantly in loamy and deep black soils and produce excellent crop. For cultivation of lathyrus under *utera* system (relay cropping), no tillage is required. However, for planting after harvest of rice, one deep ploughing followed by cross harrowing and planking is necessary.
- Sowing Time: Crop is sown on residual soil moisture after harvest of kharif during last October to early November as pure crop. In utera cropping last week of September or first week of October.
- Sowing Method: Utera/Paira cultivation- In utera cropping seeds of small seeded lathyrus is generally broadcasted in standing paddy crops (2-3 weeks before its harvest, after draining the excess water by the end of September or early October). However, planting time largely depend upon cessation of monsoon rains and maturity of rice crops. Seed must be inoculated with Rhizobium and PSB before broadcasting.
- Seed Rate & Spacing: 70-80 kg/ha for broadcasted sowing in utera system and 40-60 kg/ha in line sowing is required.; Under utera cropping sown as broadcasted in-between the rice rows. Whereas normal spacing 30 cm x 10 cm is recommended.
- Varieties: Refer table 11.3.
- Cropping System: It is grown as single crop of the year in areas where water gets accumulated during rainy season or as a relay crop after paddy often as utera/paira crop in standing paddy, due to its ability to withstand in high moisture conditions at sowing time and moisture stress during growth period.
- Water Management: The crop is grown as rain fed crop on residual moisture. However, under high moisture stresses one irrigation at 60-70 days after sowing may be remunerative in terms of production.
- Plant Nutrient Management: Under utera cropping the crop is grown on residual fertility of rice. However, it respond well to phosphorus up to 40-60 kg /ha except in the case if

- grown on highly phosphorus fertilized paddy field. For normal crop 100 kg DAP + 100 kg gypsum/ha is a optimum dose of fertilizer applied as basal dose 2-3 cm side and below the seed with the help of ferti-seed drill, is recommended.
- Weed Management: For normal sown crops one hand-weeding at 30-35 days after sowing (if soil condition permit). Weeds can also be managed effectively by spray of fluchloralin (Basalin) 35 EC @ 1 kg a.i./ha in 500-600 liters of water as pre-plant incorporation.
- Harvesting, threshing & storage: Harvest the crop with the help of sickle when colour of pods change to brown and grains are at dough stage having approximately 15% moisture in-side them. Harvested produce may be allowed to dry in sunlight for a week. Harvested produce after 3-4 days sun drying is roaped in the bundles and transferred to threshing floors. Threshing is done by beating with sticks or trampling under the feet of bullocks. The clean seed should be sun dried for 3-4 days to reduce their moisture content up to 9-10%. Now the produce should be safely stored in appropriate bins. The small quantity of the produce can also be protected by mixing inert material (soft stone, lime, ash, etc).
- Yield: A well managed crop can easily give 8-10qtls/ha yields under direct sowing and 3-4 qtls under utera cultivation.

11.5.3 Plant Protection Measures

Insect Pest/ Disease/	Nature of Damage/	Control Measures
CO	Symptoms	
i. Aphid	The adults and nymphs suck	Monocrotphos @ 0.04% or
	the juice from the leaves as	Metasystox.
	a result, leaves turn brown	
	and crumpled and the plant	
	look sick.	
ii. Rust	Pink to brown pustules	i. Grow early maturing variety.
(Uromyces fabae)	appeared on leaves and	ii. Seed Treatment with
	stems. In severe attack, the	Agrosan GN @ 2.5 g/kg seed.
	affected plants amy dry.	iii. Spray the crop with Maneb,
		Zined or Ferbam @ 2.5
		g/litre.
iii. Downy Mildew	Brownish cottony growth of	Spray with Agrosan GN
(Peronospora spp.)	fungus may be seen on the	(0.25%)
	lower surface of leaf. Inside	
	growth yellow to greenish	
	spots are also visible.	
iv. Powdery	Symptoms first appeared on	Wettable Sulphur @ 3 gm/ litre
Mildew	all the aerial part of plant.	or Dinocap @ 1 ml/litre of
(Erysiphe polygoni)	While powdery masses of	water.
	spores formed on leaves	
	which may collapse and	
	cover the whole leaf with	
	powdery growth.	

(Table–11.3): Recommended Varieties/Characteristics

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
Bio L-212	IARI	1997	NEPZ (East	15.0	108-116	Tolerant to stress,
(Ratan)			UP, Bihar,			Low ODAP, Bold
			West Bengal)			seed, Blue flower.
Prateek	IGKV,	2001	M.P.	6-9	110-115	Tol. to downy
	Raipur			(Utera)		mildew & mod.
				11-15		Resistant to
				(sole)		powdery mildew.
Maha	IGKV	2007	Chhattisgarh	15	94	Tol. to nematode
Teora						& thirps, mod.
						Resistant to PM

CZ- Central Zone (MP., Maharashtra, Chhattisgarh, Gujarat) NEPZ-North East plane Zone (East Uttar Pradesh, Bihar, Jharkhand, West Bengal). $ODAP = \beta$ -N-Oxalyl-L- α , β diaminopropionic acid

11.6 Recommendation to achieved higher production

- Deep summer ploughing once in 3 years.
- Seed treatment should be done before sowing.
- Application of fertilizer should be based on soil test value.
- In Lathyrus foliar spray of 2% urea or 20 ppm Salicylic acid at flowering and pod formation stage increases the yield.
- Weed control should be done at right time.
- Adopt integrated approach for plant protection.

RAJMASH (FRENCHBEAN)



Chapter-12

Rajmash

Botanical name : *Phaseolus vulgaris L*.

Synonym : Kidney bean, common bean, haricot bean, snap bean and French bean

Origin : Central America and south Mexico

Chromosome nos: 2n = 22

12.1 Economic Importance

Rajmash, an important pulse crop, with high yielding ability as compared to gram and pea, require focussed attention both at the development and policy front. It is grown in Maharahstra, H.P., U.P., J&K., and NE states covering 80-85 thousand ha area. However, its cultivation during rabi and summer is also gaining popularity in northern Indian plains. Traditionally Rajmash is grown during kharif in Hills of Himalayas, however; high yield is attainable in Rabi in plains due to better management.

12.2 Nutritive Value

Protein	26-28%	Calcium	260 mg/100g
Fat	0.3-0.5%	Phosphorus	410 mg/100g
Carbohydrate	62-63%	Iron	5.8 mg/100g
Fiber	17-18%	Calorific value	345-346 Kcal/100 g

12.3 Rajmash Agronomy

12.3.1 Botanical Description

Plants may be bushy or climbing type. Bushy cultivars are day neutral, early maturing, dwarf plants, 20-60 cm tall with lateral and terminal inflorescence and consequently determinate growth habit. Climbing cultivars are indeterminate, and may grow 2-3 m tall if they have support to climb by twining. The pods are slender, 10-20 cm long, straight or curved and terminated by a pointed beak. They contain 4-6 seeds which vary greatly in size and colour. Germination is epigeal

12.3.2 Production Technology

- Climate: In the hilly region it is grown during kharif and in lower hills/tarai region, sown as spring crop. In north-east plains and hilly tracts of Maharashtra, it is cultivated during rabi. It is highly sensitive to frost and water logging. The ideal temperature range for proper growth of this crop is 10-27°C. Above 30°C, the flower drop is a serious problem. Similarly, below 5°C the flowers and developing pods and branches are damaged.
- Soil & Field Preparation: The crop can be grown in light loamy sand to heavy clay soil under adequate moisture. Among various pulses, Rajmash is most sensitive to salt stress and sodicity. Therefore, soil must be free from excessive soluble salts and neutral in reaction. Rajmash having bold and hard seed coat needs a good seed bed accomplished by thorough primary tillage like ploughing, harrowing or discing and planking. A good seed bed have

friable but compact soil adequate moisture and free from weeds and plant debris of earlier crop. Acidic soils of the hills must be treated with lime before sowing.

- Sowing time & spacing: Kharif (Hills) last week June to first of July; Rabi (Plains) IInd fortnight of October and for spring (Lower hills) IInd fortnight of March and for bold seeded 100-125 kg/ha. Kharif (Hills) 45-50 cm x 8-10 cm; Rabi & Spring 40 cm x 10 cm (irrigated) 30 cm x 10 cm (Rainfed).
- Varieties: Selection of varieties as per the growing season and purpose of cultivation from Table ó 11.1.
- Cropping System: In hills, it is grown as intercrop with maize in 1:2 ratios. In-between two rows of maize sown at 90 cm apart, two rows of Rajmash are adjusted at 30 cm spacing with the plant population of 120000 of Rajmash and 40000 of maize. It is also grown mixed with maize and soybean. In plains it is grown as spring season crop after harvesting of potato and mustard. It is also found quite compatible for intercropping with early potato due to its high nitrogen requirement and wet moisture regime in 2:2 or 2:3 row ratios.
- Water Management: Rajmash is the most irrigation responsive pulse crop due to its shallow root system and high nutrient requirements. It requires 2 to 3 irrigations in NEPZ and 3 to 4 irrigation in CZ for achieving highest productivity. Irrigation at 25 days after sowing is most critical followed by irrigation at 75 days after sowing.
- Plant nutrient management: Unlike other Rabi pulses, Rajmash is very inefficient in biological nitrogen fixation owing to poor nodulation due to non availability of suitable and efficient Rhizobium strain for Indian plains. Hence, it requires relatively higher doses of fertilizer N. For enhanced productivity, application of 90-120 kg N ha⁻¹ has been found optimum. Half of the nitrogen should be applied as basal during sowing and rest half as top dressing after first irrigation.Rajmash responds well to phosphorus application like cereals. Its P requirement is distinctly higher than other pulse crops, significant response to P application has been obtained up to a level of 60-80 kg P₂O₅ per ha.
- Weed Management:One hand weeding/hoeing at 30-35 days after sowing or application of a pre-emergence herbicide like pendimethalin @ 1 to 1.5 kg a.i./ha in 500-600 liters of water immediately after sowing helps to keep the losses by weeds below ETL (Economic Threshold Level).
- Harvesting, threshing & storage: The crop mature in 125-130 days. Plants are cut with sickles after attaining full maturity judged by severe leaf fall, changing colour of pods and hardness of the grains.
 - Harvested materials, after 3-4 days sun drying, is collected in bundles to the threshing floors. Threshing is done by beating with sticks or trampling under the feet of bullocks. The clean seed should be sun dried for 3-4 days to bring their moisture content at 9-10%. To avoid further development by bruchids and other storage pests it is recommended to fumigate the storage material with ALP @ 1-2 tablets per tonne before onset of monsoon and again after

the monsoon. The small quantity of the produce can also be protected by mixing inert material (soft stone, lime, ash, etc) or by smearing edible/non-edible vegetable oils or by mixing plant products like neem leaf powder at the rate of 1-2% w/w basis.

• Yield: A well managed crop can easily give 20-25 qtls/ha yields under irrigated conditions of plain and 5-10 gtls/ha under rain fed conditions of hill with 40-50 gtls/ha of straw for cattleds

12.3.3 Plant Protection Measures

A. DISEASES

i) Anthracnose

Symptoms: Pale brown sunken spots may appear on the cotyledons of infected seedlings. Lesions on leaves are dark brown. They are restricted to the veins on lower leaf surface. On stems, lesions are elongated and sunken.

Control Measures: Seed treated with Carbendazim & Thirum (1:1); ii) Spray Mancozeb 0.25 % or Carbendazim 0.1 % of 2-3 foliar spray at 45, 60, 75 DAS; iii) Remove from the field and destroy crop debris after harvest; iv) Practise a 2 to 3 year rotation; v) Avoid overhead irrigation; vi)

Avoid movement of workers in the field when wet.



Stem Blight

Symptom: Small water-soaked spots are first symptoms observed on leaves & appear within 4 to 10 days of infection. Develop, centre spots becomes dry and brown. The lesion is surrounded by a narrow band of bright yellow tissue.

Control Measures: i) Foliar spary of Carbandazim @ 0.2 % is recommended; ii) Early or timely sowing; iii)Planting in well drained soil; iv) Avoid dense planting.



iii) Angular leaf spot

Symptoms: Fungus produces a grey mould on the lower surface of the spots. Infected pods have brown blotches. The spots may increase in size, join together, and cause yellowing and necrosis of the affected leaves.

Control Measures: i) Seed treated with Carbendazim @2-3 g/kg. of seeds; ii) 3 Foliar spray of Carbendazim @0.1% (1 gm/lit.) starting at the appearance(5-6 weeks after sowing) at 15 days interval; iii) Plough under bean debris after harvest; iv) Practice a 2-3 year crop rotation without legumes; v) Do not work in bean fields when the plants are wet.



B. Insect-Pest

i) Leaf Miner

Nature of damage: Severely mined leaves may turn yellow and drop. Severely attacked seedlings are stunted and may eventually die. It may be seen in vegetative stage.

Control Measure: Spray of Oxydemeton methyl (Metasystox) 1 ml/liter of water and repeat at 15 day interval if required; ii) Roughing of infected plants; iii) Handpick & destroy mined leaves; iv) Whenever necessary spray the crop with neem products; v) Neem water extracts and neem oil give good control of leaf miners; vi) Remove and destroy crop residues and all plant parts with symptoms of damage by bean flies.



ii) Stem fly

Nature of damage: Stem becomes to swell and split and reducing formation of lateral roots. Attacked plants produce adventitious roots in compensation. Young seedlings and plants under stress wilt and die.

Control Measures: i)Seed Treated with Chlorphyriphos @8ml/kg seed; ii) Soil application by Phorate 10 G @ 10 kg./hac; iii) Mulch (e.g. with straw and cut grasses) helps conserve moisture, promote adventitious root development and enhances tolerance to maggot damage; iv) Avoid planting beans near cowpea, soybean and many other leguminous crops, that may be the source of bean flies.



iii) Black Aphids

Nature of damage: Aphids feed by sucking plant sap. Heavily infested plants usually have wrinkled leaves, stunted growth and deformed pods. Plants, in particular young plants, may dry out and die under heavy aphid attack.

Control Measures: i) Bio treat.-Inundative release of Coccinella septempunctata @ 1000 adult/400 sqm; ii) Practice a 2-3 year crop rotation without legumes; iii) Spraying the systemic insecticide like Dimethoate or Oxydemeton methyl @1 ml/liter of water.



12.4 Recommendation to achieved higher production


- Deep summer ploughing once in 3 years.
- Seed treatment should be done before sowing.
- Application of fertilizer should be based on soil test value.
- Weed control should be done at right time.
- Adopt integrated approach for plant protection.

(Table-12.1): Recommended Varieties/Characteristics

Variety	Source	Release/ Notification Year	Area of adoption (Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
Varun (ACPR 94040)	IIPR	2002	Maharashtra	14-16	66-68	Tolerant to Anthracnose
IPR 96-4 (Amber)	IIPR	2002	NEPZ (East UP, Bihar, W.B.).	15-16	139	Res. to BCMV & Leaf Curl. Red
Ankur (RSJ-178)	Central	2005	Rajasthan	12	110-120	Moderately resistant to root rot, leaf crinkle and leaf spot dry root rot.
Gujarat Rajma-1	SDAU	2006	Gujarat	20	30-35	Moderate resistant to bean common mosaic virus
VL Rajma 125	VPKAS	2007	Uttrakhand	14-15	82-85	Resistant to root rot, Mod. Resistant to Anthracnose, angular leaf spot & rust
VL Bean 2	VPKAS, Almora	2008	Uttrakhand	14-15	82	Resistance to root rot, mod. Resistant to anthracnose, angular leaf spot and rust
Arka Anoop		2012	Karnataka	18-20	70-80	Suitable for eastern dry zone of karnata ka in both kharif an d rabi season

NEPZ-North East plane Zone (East Uttar Pradesh, Bihar, Jharkhand, West Bengal).

COWPEA (LOBIA)





Chapter-13

Cowpea

: Vigna anguiculata Botanical Name

: Lobia, Barbati, Black eyed pea Synonymous

Origin : Africa Chromosome : 2n = 22

13.1 Economic Importance

This crop is known as drought hardy nature, its wide and droopy leaves keeps soils and soil moisture conserved due to shading effect. Initial fast growth with fast penetrating root system and strong stomatal sensivity justify its initial establishment in soil moisture deficit conditions. It is also known as Cowpea, black-eyed pea or southern pea etc. and has multiple uses like food, feed, forage, fodder, green manuring and vegetable. Cowpea seed is a nutritious component in the human diet, and cheap livestock feed as well. Choice of cowpea as vegetable is due to being palatable, highly nutritious and relatively free of metabolites or other toxins. Fresh leaves and fast growing twigs are often picked up and eaten like spinach. Immature shaped pods are used in some way as snap beans often being mixed with other foods. Both the green and dried seeds are suitable for canning and boiling as well.

13.2 Nutritive Value

Protein	26-28%	Calcium	0.08 ó 0.11%
Fat	0.30.4%	Iron	0.005%
Dietary Fiber	18.2%	Calorific value	345-346 Kcal/100g
Carbohydrate	63-64%		

Essential amino acids (lysine, leucine and phenylalanine)

13.3Agronomic Importance

An important component of farming system in resource constraints agriculture, this legume has great potential in India for successful cultivation in kharif and summer in northern India and throughout the year in peninsular India. It also leave 30-40 Kg N/ha in the soil for the succeeding crop.

13.4Crop Status

It is widely grown in tropics and subtropics of Asia, Africa, central and southern America and parts of southern Europe and USA. However, central and western Africa alone account for more than 60% of world acreage with marginal and sub marginal farmers in the semi-arid and sub-humid regions. According to an FAO estimate, Nigeria alone produces 2.1 mt of dry grain out of 3.3 mt of total worldwide, in 2000. During the same year, global area sown to cowpea was 9.8 mha (9.3 mha in West Africa) with average productivity of 337 Kg/ha whereas, productivity of Nigeria was comparatively higher (417 Kg/ha).

In India, cowpea is grown as sole, inter-crop, mix-crop and in agro-forestry combinations. Exact statistics on its area is not available but, is estimated to be cultivated in almost half of 1.3 m ha of area occupied by Asian region. Other Asian countries are Sri Lanka, Bangladesh, Myanmar, Indonesia, China, Korea, Pakistan and Nepal. In Indian context, it is a minor pulse cultivated mainly in arid and semi arid tracts of Rajasthan, Karnataka, Kerala, Tamilnadu, Maharashtra and Gujarat. In North India, it is grown in pockets of Punjab, Haryana, Delhi, and West UP alongwith considerable area in Rajasthan.

13.5 Cowpea Agronomy

13.5.1 Production Technology

- Climate: Cowpea is warm weather and semi arid crop, where temperature ranging from 20°C to 30°C. Minimum temperature for seed establishment is 20°C and above 32°C temperatures development of root is cease. For maximum production day temperature 27°C and night temperature 22°C required. It is sensitive to cold and below 150C temperature yield adversely affected. It can grow under shade of tree but can not tolerate cold or frost.
- Soil: Well drained loam or slightly heavy soil are best suited. In colder climate somewhat sandy soil preferred as crop mature earlier in them. It can grow successfully in acidic soil but not in saline/alkaline soil. In hard soil, one deep ploughing followed by two or three harrowing and planking are sufficient. In normal soil only two harrowing & planking is enough. For summer season crop give a irrigation immediately after harvesting of Rabi crop.

Field preparation and mulching:

In hard soil, one deep ploughing followed by two or three harrowing and planking are sufficient. In normal soil only two harrowing & planking is enough. However, field leveling is must to avoid water logging. However, reduced and zero tillage method can also be followed if effective weed control is assured through chemical herbicides.

Addition of grass mulch increase soil moisture in the root zone (0-15 cm soil depth) and significantly decreases maximum temp of soil along-with diurnal fluctuation. This provides a stable environment for seedling establishment and growth than the unmulched soil. A combination of minimum tillage and straw mulch as the least risky and hence, most appropriate soil management system for dry season Cowpea in rice fallows. However, growth and yield of cowpea grown after rice do not affect significantly by tillage or no tillage but the mulch application significantly increase growth and yield due to better ability or mulched plot in storing soil moisture during the growing seasons.

- Sowing time: Kharif- With onset of monsoon ranging from early June to end of July, Rabi- October-November (southern India), Summer - 2nd to 4th week of March (grain), February (Fodder), Hills: April-May, Green manuring- Mid June to 1st week of July.
- Sowing method: Broadcasting, in centre of furrow areas then modified into ridges after a month. Draw 30 cm wide and 15 cm deep drainage channel at 2 meter interval to drain excess rainwater after sowing. Sowing on rice bund on either side on the day of paddy transplanting during second season. Sowing by broadcasting immediately after paddy harvest in summer. Seed depth should be 3-5 cm.
- Seed rate & Spacing: For pure crop: 20-25 kg ha (grain), for fodder and Green Manure-30-35 kg/ha. During summer 30 kg/ha for grain and 4- kg/ha for fodder and green

manuring; Row to row-30 (Bushing) to 45 cm (spreading), Plant to Plant-10 (Bushing) to 15 cm (spreading).

- Varieties: Varieties is given in table -12.1 other than the specific as follows (a) Grain: C-152, Pusa Phalguni, Amba (V 16) (M), Ramba (V240) (M), Swarna (V-38) (M), GC-3, Pusa Sampada (V-585), Shreshtha (V-37) (M).
- Fodder: GFC 1, GFC 2, GFC 3,-Kharif season, GFC-4 Summer (25-35 tonnes/ha), Bundel Lobia-1, UPC-287 and UPC-5286, Russian Giant, K-395, IGFRI-5450 (Kohinoor), C-88 (20-35 tonnes/ha inPunjab), UPC 5287, UPC-4200 (NE India).

• Cropping system

Grain/vegetable	Fodder
Cowpea-Wheat-Mung/Cheena	Sorghum+cowpea-berseem-maize+cowpea
Cowpea-Potato-urd/bean	Maize-berseem/oat- maize+cowpea
Maize/Rice-Wheat-Cowpea	Sudan grass- berseem/oat- maize+cowpea
Maize-Toria-Wheat-Cowpea	Cowpea-berseem-maize+cowpea
Rice-Rice-Cowpea	
Rice-Cowpea	
Rice-Mustard-Cowpea	

• Inter cropping:

Growing one or two rows of cowpea in widely spaced crops and in-corporating the biomass after picking pods can increase soil fertility and yield of companion crop. The improvement in this system can further be made by pairing the rows of main crops and taking one or two rows of cowpea in between two paired rows of either of pigeonpea, maize and sorghum. Here, we can get 5-7 qtl/ha grain yield of cowpea without any adverse effect on main crop yield.

It can also be grown as floor crop in coconut garden and intercrop in tapioca in Kerala and as sole crop in single or double crop rice fallows in rabi or summer season respectively.

- Water Management: For rainy season crop drainage is more essential than irrigation.
 Crop can tolerate flooding upto 2 days at flowering and pod setting thereafter, a marked decrease in yield and its attribute. Early sown rainy season crop may require one or two irrigation in pre monsoon/delayed onset of monsoon.
 - For summer crop, irrigation is most critical among all inputs followed by weeding and fertilizer. Generally, crop required 5-6 irrigation depending on soil, prevailing weather conditions etc, at an interval of 10-15 days. Increasing moisture regime from dry to medium wet, result in significant yield improvement. The response to irrigation is in order of flowering> pod filling>vegetative.
- Abiotic stress management: Salinity, water logging, toxicity or deficiency of minerals are common abiotic stress. The crop is more sensitive to drought at onset of flowering and during reproductive phase. Maintenance of adequate *Kø in soil improves plant water relations, photosynthesis and yield and overcoming soil moisture stress alongwith improving carbon partitioning in cowpea.

Seed treatment with thiourea (seed soaking in 500 ppm soln) followed by two foliar spray at vegetative and flowering phase is another option to avoid moisture stress through

- enhancement of photosynthesis efficiency and nitrogen metabolism there by giving higher yield.
- Crop improvement and breeding programme are needed for varieties with dwarf and erect growth habit, extra-earliness (65-90 days) with synchronous maturity, development of multi-purpose varieties, breeding for insect-pest and disease resistance, tolerance to drought, high nutritional quality.
- Plant nutrient management: Apply FYM/compost- 5-10 t/ha as basal with last ploughing. Both these bulky organic manure can be substituted by humic substances granule. 15-20 kg N/ ha as starter dose in poor soils (organic carbon<0.5%), 50-60 kg/ha P₂O₅ and 10-20 kg. K₂O/ha to promote growth and to mitigate the impact of water stress in plants when subjected to sub optimal soil stress. In acidic soil, lime pelleting of seed is beneficial alongwith Rhizobium inoculation. Add finely powdered (300 mesh) calcium carbonate to moist freshly Rhizobium treated seeds and mix for 1-3 minutes until each seed is uniformly pelleted. Lime requirement varied from 0.05 Kg to 1 Kg/ 10 Kg seed depending on seed size.
- Weed management: Weed can reduce crop yield upto 50-62%. Integrated approach includes agronomic (improved) practices like sowing at proper time, proper cropping geometry, optimum plant density, intercropping, intercultivation, irrigation and the need based supplement, use of chemical herbicides. One hand weeding at 20-30 DAS-followed by one more weeding after 20-25 DAS if required.
- Chemically, weed can be controlled by pre-planting spray of Basalin @ 1 kg a.i./ha as pre emergence in 800-900 litres of water. Application of pendimethaline @ 0.75 kg.a.i./ha combined with one hand weeding at 35 DAS resulted in two fold increase in marginal benefit cost ratio and highest weed control efficiency.
- Harvesting & threshing: The crop matures in 125-130 days. Plants are cut with sickles after attaining full maturity judged by severe leaf fall, changing colour of pods and hardness of the grains. Harvested materials after 3-4 days sun drying, is collected in bundles to the threshing floors. Threshing is done by beating with sticks or trampling under the feet of bullocks. The clean seed should be sun dried for 3-4 days to bring their moisture content at 9-10%.
- Yield By adopting improved management practices yields up to 12-15 Q/ha could be realised.

13.5.2 Plant Protection Measures

A. Diseases

i) Bacterial Blight

Symptoms: The germinating seedling turn brown-red and die. Irregular to round spots brown in colure with chlorotic halos, appear on leaves, and later spread to stem. Stem may break, pods are also infected leading to shrivelled seeds.

Control Measures: i) Grow resistant varieties; ii) Use healthy and disease free seeds; iii) In case of severe infection, crop may be sprayed with 0.2 % (2g/liter) copper oxychloride (Blitox).



ii) Cowpea Mosaic

Symptoms: It is caused by a virus transmitted by aphids. The affected leaves become pale yellow and exhibit mosaic, vein banding symptoms. The affected leaves become reduced in size and show puckering. Pods are also reduced and become twisted. **Control Measures:**i) Use healthy seed from healthy crop; ii) For controlling aphids spray Oxydemeton methyl 25 EC (Metasystox) @ 1 ml/liter or Imidacloprid 17. 8 SL @ 0.2 ml/liter of water and repeat the spray after 10 days of first spray.



iii)Powdery mildew

Symptoms: Powdery mildew are visible on all the aerial parts of the affected plants. Symptoms first start from leaves and then spread to stem, branches and pods. This white growth consists of the fungus and its spores. Affected leaves become twisted and smaller in size.

Control Measures:i) After harvest, collect the plants left in the field and burn them; ii) The disease can be controlled by spray of wettable sulphur @ 3g/liter or carbendazim @1 g/liter of water.



C. Insect-Pest

i) Cowpea pod borer

with the top shoot. Caterpillar bore into the pods and feed on the seeds, if flower and pods are not available larvae feed on foliage. **Control Measures:**i) Collect and destroy the eggs and young larvae; ii) The young caterpillar can be killed by dusting 2% methyl parathion @ 25-30 kg per hectare or spray of quinalphos @2 ml/liter of water; iii) Fix 3 feet stick in the field @10/ha bird parches to attract predatory birds.

Nature of Damage: The caterpillar rolls the leaves and web these



ii) Hairy caterpillar

Nature of damage: It is major insect of cowpea. It is cut juvenile plants and eat away all the green matter of the leaves.

Control Measures: Collect and burn the eggs and burn the eggs and larva of insect; ii) The young caterpillar can be control by spray of Chloropyriphos or Quinolphos @ 2ml/liter of water.

iii) Aphids and Jassids

Nature of Damage: The adult and nymphs of these pests suck the juice from the leaves and the damage is more severe when the plants are young. As a result of sucking of sap, the leaves turn brown and crumbled and the plant look sick.

Control Measures: Spray of Oxydemeton Methyl 25 EC (Metasystox) @ 1 ml/ liter or Dimethoate 30 EC @ 1.7 ml/ liter of water.



iv) Bean fly/Stem fly

Nature of damage: Bean fly causes the characteristics swelling of stem at ground level where the maggots burrow onto the stem. The maggots puppets at the base of the plant and the stem grows it often cracks. The petiole often shows dark streaks where the maggots have move through and damage tissue.

Control Measures: Keeping the field clean from legume debris; ii) Application of Phorate (Thimet) 10 G @ 10 kg per hectare in furrows at the time of sowing is effective for avoiding infestation.



13.6 Recommendation to achieve higher production

- Deep summer ploughing once in 3 years.
- Seed treatment should be done before sowing.
- Application of fertilizer should be based on soil test value.
- Weed control should be done at right time.
- Adopt integrated approach for plant protection.

(Table - 13.1): Recommended varieties of cowpea/characteristics

Variety	Source	Release/	Area of adoption	Ave. yield	Days to	Special characteristics
		Noti. Year	(Zone/State)	(q/ha)	maturity	
Gujarat	GAU	1990	CZ (MP, MH& Gujarat)	12-14	65-85	Seed bold, amber colour
Cowpea-3						
V-240	IARI	1993	All Zones	14.0	80	Tall, Indeterminate, seed
						red
Vamban - 1	TNAU	1997	Tamil Nadu	9.5	65	Erect, dwarf, seed white
Gujarat	GAU	1999	Gujarat	8-5	80-90	Seed bold, amber colour
Cowpea-4						
KBC-2	UAS	2001	Karnataka	9.5	95-105	Semi-determinate, seed
						light brown
RC-101	TNAU	2001	Rajasthan	8.5	85-90	Early, Determinate, seed
						white
CO-6	TNAU	2001	Tamil Nadu	14.0	85-90	Early, bold seeded
V 578 (Pusa	IARI	2004	Delhi	12		Early, Resistant to
sampada)						yellow mosaic virus
CL-367	PAU	2006	Punjab	12	95-100	Tolerant to YMV
RCP-27 (FTC-	RAU	2006	Rajasthan	6-13	69-79	Resistant to YMV
27)						
UPC 622	GBPUAT	2007	Uttrakhand Assam,	4-5	145-150	Tolerant to drought
			U.P., M.P., J & k, H.P.,			resistant to YMV,
			Punjab, Raj., Har.,			Anthracnose, root/collar
			WB., Odisha, Bihar,			rot and bacterial leaf
			and Jharkhand			blight,
Khalleshwari	IGKV,	2007	Chhattisgarh	6-7		RRF in rabi with
	Raipur					restricted irrigations
						and rainfed upland in
						kharif season
Kashi	IIVR,	2008	Punjab, UP, Bihar,	150-175	50-55	Reistant to golden
Kanchan	Varanasi		Jharkhand, Orisha,	(Pods)		mosaic virus, <i>Pseudo-</i>
(VRCP 4)			CG, MP, AP	,,		cercospora cruenta
(1101)			,,			diseases,
	1	l				arocases,

Variety	Source	Release/Noti. Year	Area of adoption(Zone/State)	Ave. yield (q/ha)	Days to maturity	Special characteristics
Swarna Harita (IC285143)	ICAR Res. Station,	2008	Assam, U.P., M.P., Kerela, A.P., Punjab, Raj., WB., Odisha, CG., Jharkhand, TN.	60- 150(Pods)	75-90	Resistant to rust and mosaic viral disease & tolerant to pod borer.
UPC 628	GBPUAT	2010	Punjab, UP, Bihar, Jharkhand, Orisha, CG, MP, WB,MS	3.5-4.0	145-150	Iriigated Summer, and rainfed condition,Medium late variety
IT ó 38956-1	UAS,GKVK, Bangalore	2009	Karnataka	10-12	80-85	Rainfed areas of eastern dry region
Hisar Cowpea 46 (HC 98- 46)	CCSHAU	2009	Haryana	10	65-70	Resistant to YMV
Pant Lobia -1	GBPUAT	2010	Uttrakhand, UP	20	130-135	Moderately resi. to Aphids, Thrips, Bruchids & other field pests. Suitable for spring ,summer and Kharif season
UPC 628	GBPUAT	2010	Uttrakhand, HP, J&K, Punjab, Harya., Raj.,UP, MP, CG, Bihar, Jharkhand, WB, Odisha, Assam, Gujrat & MS	350-400 (Pods)	145-150	Tolerant ot drought and other edephic /abiotic stresses, reis. To YMV, Anthacnose/leaf blight, Aphids, Semilooper, Flea Beetle/Defoliators, Pod borer/bugs & Root knot nematode, tolerant to storage Beevil
HIDRUDAYA	ORARS, Kerela	2010	Kerela	10-11	50-55	Tolerant to leaf rust, Aphids, Pod borer & American Serpentine leaf minor, summer season
C 519 (Himachal Lobia 11)	CSKHPKV, Palampur	2010	Himachal Pradesh	15-16	80-85	Resistant to Cercospora leaf spot, YMV, Low hills, Sub-tropical zone under rainfed condition in kharif
PKB 4	UASGKVK, Banglore	2012	Karnataka	11-13	80-85	Resistant to Bacterial leaf blight, Rust & Pod borer, suitable for early kharif season
PKB 6	UASGKVK, Banglore	2012	Karnataka	10-12	80-85	Resistant to Bacterial leaf blight, Rust & Pod borer, suitable for late kharif and summer season

Variety	Year of Release	Area of adaptation	Maturity	Yield (kg/ha)	Specific Traits
DCS 47-1	2014	Southern region	80	1400	Suited for cultivation under light and loamy soil, sowing time during late kharif, resistant o kharif.
Pant Lobia-4	2015	All cowpea growing regions	60-65	1400- 1800	It has tolerance to major bacterial and viral diseases like yellow mosaic, photo insensitive and drought tolerant, adaptable to zaid season.
Pant lobia-3	2016	All cowpea growing regions	65-70	1400- 1800	50-55 cm tall. Resistant to YMV and bacterial blight, bush type, seeds are kidney to oval shape and brown in colour. It has 27% protein.
Phule Vithai (Phule CP-05040)	2016	Maharashtra	70-80	1200	Moderately resistant to color rot and leaf spot.
Pant Lobia-5	2017	Uttarakhand	65-70	1600-200	It is tolerant to aphid thrips bruchids and resistant to CYMV, suitable to summer and kharif season, protein content is 23-25 %.
Phule Rakhumai PCP 0306-1	2017	Southern zone	70-80	1000- 11002	Early maturing, suitable for optimum sown for rainfed and irrigated condition, moderately resistant to Cercospora leaf Spot.
TPTC 29 (Tirupati cowpea- 1)	2017	Andhra Pradesh, Karnataka, Tamilnadu, odisha	80-90	1000- 1100	Moderately resistant to dry root rot and YMV, Suitable, suitable for kharif, rabi and summer season.
DC 15	2017	Karnataka, Andhra Pradesh, Tamilnadu and Kerela	75-80	1000-	Tolerant to aphids, wide adaptability from deep black to red loamy soil, tolerant to pod borer, moderate resistant to dry root rot and YMV.

CZ- Central Zone (MP, Maharashtra, Chhattisgarh, Gujarat), SZ- South Zone (A.P., Karnataka, TN, Odisha)


MINOR PULSES

A.BROAD BEAN

B. RICEBEAN









A. BROAD-BEAN

Botanical Name : *Vicia faba* L.

Synonym : Bakla, Fababean

Origin : Mediterranean Region of Southern Europe and Western Asia

Chromosome : 2n = 24

- 1. Introduction: Broad-bean has high yield potential. In many countries this species is the main food legume. This crop is presently being grown sporadically in our country as a minor vegetable. However, dry seeds are also used as ¿Dalø Its seed is edible and nutritive. There appears to be every possibility of popularizing broad-bean as a new pulse crop in India. Broad-bean has shown response to inputs and better management practices and hence can be fitted into intensive cropping systems.
- **2. Botonical Description**: Strong, erect annual herb with the plant height up to 1.5 meter. Roots like other legume. Inflorescence ó axil with 6 flowers of 3-7 cm long, mostly white in colour. Self pollination is a rule but cross pollination by insect may also occur. Pods are strong and semi-cylendrical up to 30 cm in length. Seeds are greenish or brownish white to black. Test weight vary from 10-40 g/100 seed weight.
- 3. Production Technology
- 3.1 Climate: Spring season with mild summer is best.
- **3.2** Soil: Like other legumes (Rajmash) lime addition in acidic soil gives best results.
- **3.3 Field Preparation**: Like Rajmash (1 deep ploughing + 2 harrowing followed by planking).
- **3.4 Seed & Sowing** *Spring*: April (first fortnight)/Rabi: October (IInd Fortnight ó 1st week of November) with crop spacing as row to row 30 ó 35 cm and plant to plant 10 cm with sowing depth of 7.5 ó 10 cm.
- 3.5 Seed rate: 70-100 kg/ha.
- **3.6** Cropping system: Maize-Broadbean, Pearlmillet/Maize-Potato-Broadbea.
- 3.7 PlantNutrient management: 20 kg N + 40-50 kg P₂O₅/ha.
- **3.8 Weed management**: Two howing at 30 and 60 DAS. Alternatively, Fluchloralin or Pendimethalin (Pre-emergence) @ 1 kg a.i./ha can be used for effective weed management.
- **3.9 Diseases**: Root rot, Aschochyta blight, Botrytis grey mold, Cercosporal Leaf spot & Rust. (Control measures like gram).
- **3.10Insect**: Aphid, Leaf minor, Leaf Loeevil, Stem borer (control measures like lentil)
- **3.11** Harvesting, threshing: Similer to lentil.
- **3.12**Yield: 10-40 Q/ha

B. RICE-BEAN

Botanical Name : *Vigna umbellate* (Thunb.) Ohwi & Ohashi}

Origin : Himalayin region of North east

Synonym : Japanese Rice bean, bomboo bean, climbing and mountain bean,

Chromosome : 2n = 22

- 1. Importance: One of the important minor food legumes can be grown under a wide range of soil and climatic conditions in the hilly areas of Himachal Pradesh, Uttrakhand and north-east hill regions, generally as dual purpose. Its grains are also cooked in place of rice that why it is known as rice bean. It is also grown as green manure crop. Its grain production potentiality is considerably high and can serve as a good pulse crop. It has the potential to yield as high as 15-25q/ha. All the pods on a plant mature almost simultaneously and can be harvested in a single operation. There are many types and varieties of rice-bean differing in maturity, plant type and seed characteristics. Some varieties are completely free from fungal and viral diseases during kharif.
- 2. Nutritive Value: Rice-bean has a protein content of 14 to24% and is free from antinutritional factors. Thus, rice-bean offers itself for cultivation during the monsoon season, when green gram and other pulses suffer greatly from diseases.
- 3. Botanical Description: It is an annual, deep rooted herb with plant height of 30-100 cm with fast spreading habit surrounding 100-120 cm. Leaves are oval and trifoliate with 6-9 cm long. Inflorescences are 4-7 cm long with 10-18 bright yellow flowers. Flowering in 100 days. Pod length vary 12-18 cm with 6-10 grain inside them. Grain colours vary from yellow, brown, black or straw with epigeal germination and white hilum.
- 4. Production Technology
- **4.1 Climate**: Tropical climate of kharif. It can be grown successfully in high rainfall areas with good drainage where other pulses are failed due to excessive growth and diseases and pest attack.
- **4.2 Soil:** Generally grown on slopy hilly land with poor fertility.
- **4.3 Land preparation**: One normal ploughing is enough as excess field preparation can accelerate the rate of soil erosion.
- **4.4 Seed & Sowing:** IInd fortnight of August for grain, however, can be grown up to September for fodder with spacing: 45-60 cm row to row, 5-10 cm plant to plant.
- **4.5 Seedrate**: 40-50 kg/ha for grain & 60-75 kg/ha for fodder.
- 4.6 Varieties: Pant rice bean 1, Pant rice bean 2, K-1, Bidhan rice bean-2 (KRB-4)
- **4.7 Cropping system**: Grown as mixed with Jute, Maize and Finger millet on Hills. Also grown commonly in kitchen gardens for meeting vegetables pulse and forage need of house hold.
- **4.8 Plant nutrient management**: Grown on residual soil fertility.
- **4.9 Water management**: Grown as rainfed in high rainfall areas hence instead of irrigation, drainage is important.
- **4.10 Weed management**: One hoeing 30 DAS is enough.
- **4.11 Plant Protection**: No need to do any spray as very rare infestation of pest & diseases is observed.
- **4.12 Harvesting**: August sown crop ready to harvest in Feb. last (Duration 120-130 days).
- **4.13 Threshing:** Like moong.
- **4.14 Yield:** 10 qtl/ha.

Chapter-14

Market Scenario: 2016-17

14.1 Production Scenario- 2016-17: An Analysis

- **Total Pulses** During, 2016-17 the production of pulses in India has been 229.54 Lakh tonnes (*IV*th*Adv*. *Est.*) which is ever highest production. This is 30% higher over the Normal and 40% higher than the last year.
- Arhar-During, 2016-17 the production of Pigeonpea has been 47.78 Lakh tonnes $(IV^{th}Adv.$ *Est.)* which is ever highest production. This is 68% higher over the Normal and 87% higher than the last year.
 - Maharashtra, with about 30% of National production remains at Ist rank in the country. More than 90% of the production of this crop is contributed by 8 states *viz.*, MS, Karnataka, MP, UP, Gujarat, Jharkhand, Telangana and AP.
- Urdbean-During 2016-17, the production of Blackgram has been at 28.05 lakh tonnes (kh-21.70 + rabi- 6.35 lakh tons), the ever highest production. This is 50% higher over the Normal and 44% higher than the last year.
 - Madhya Pradesh, with >26 % of National Production during kharif and >18% of total production in a crop year (kh. + rabi) ranks at I^{st} position at all India level. The rabi season, highest urd production is from AP which is >50 % of national production.
 - More than 90% of urdbean production comes from MP, AP, UP, TN, MS, Rajasthan, Jharkhand, Gujarat, WB and Karnataka.
- Mungbean-During 2016-17, Greengram also recorded the ever highest production at 21.63 lakh tons (kh-16.15 + rabi- 5.48 lakh tons), which is 44% higher over the Normal and 36% higher than the last year.
 - Rajasthan, with 48% of total all India production during kharif and >31 % collectively in a crop year, ranks Ist in the country. During, rabi, TN with 19 % of national production stands at Ist position.
 - More than 90% of Mungbean production comes from 10 states, namely Rajasthan, Maharashtra, TN, AP, Bihar, MP, Odisha, Gujarat, Telangana and Karnataka
- **Gram-**During 2016-17, Gram production at 93.26 lakh tons, which is 9% higher over the Normal and 32% higher than the last year. The ever highest production of gram was 95.30 lakh tonnes during 2013-14.
 - Highest production of Gram is from MP with 41% of contribution to the National Production followed by Rajasthan (14%), Maharashtra (13%), Karnataka (>8%) and AP (>6%).
- Lentil-Lentil and kulthi are considered under *other pulses category* by the DES, which provides the statistics of these pulses only at the final production estimates stage. During 2015-16, the production of lentil was at 9.76 lakh tons which was 7% less than the Normal

production. UP with 33% of national production ranks at Ist followed by MP (32.75%), Bihar (16.96%) and WB (6%).During 2016-17, as per the WWWR coverage and the Normal yield, the tentative production of lentil is likely to be 12.74 lakh tonnes.

(Table-14.1): National production of pigeonpea, urd, mung & lentil

(Production: Lakh tonnes)

Crop/Year	Normal	2015-16	2016-17	Chang	ge Over
1 st July to 30 th	(Avg. 2011-12 to 2015-16)	Final Est.	4 th Adv. Est.	Normal	2015-16
June	2010 10)				
Pigeonpea	28.44	25.61	47.78	19.34 (68%)	22.17 (87%)
Urd	18.72	19.45	28.05	9.33 (50%)	8.60 (44%)
Mung	15.05	15.93	21.63	6.58 (44%)	5.70 (36%)
Gram	80.90	70.60	93.26	12.36 (15%)	22.66 (32%)
Lentil	10.44	9.76	12.74*	2.30 (22%)	2.98 (31%)
Total Pulses	176.37	163.48	229.54	53.17(30%)	66.06 (40%)

Source: DES (DAC&FW); * Estimated by DPD on the basis of area coverage reported in WWWR & Avg. Yield

(Table- I): State-wise area production of pigeonpea

{Area: lakh ha, Production -lakh tonnes}

State	Nor	mal	201	3-14	201:	5-16		6-17*	Prod. % change	
										er
	Area	Prod.	Area	Prod.	Area	Prod.	Area	Prod.	Normal	2015-16
Maharashtra	12.00	8.31	11.41	10.34	10.39	4.66	15.33	13.89	20.11	63.96
Karnataka	7.27	4.05	8.24	5.88	6.48	2.63	12.14	8.66	52.80	125.20
M.P.	5.26	4.31	4.64	3.32	5.79	6.25	6.90	7.82	42.10	20.02
Gujarat	2.25	2.46	2.10	2.09	2.27	2.37	3.34	3.69	60.98	65.69
Uttar Pradesh	2.97	2.57	3.01	2.71	2.65	1.83	3.38	3.36	50.87	100.33
Telangana	2.61	1.19	2.64	1.40	2.48	1.04	3.87	2.15	151.83	198.78
Jharkhand	1.80	1.78	1.97	2.05	1.94	1.74	1.94	2.02	63.75	66.72
A.P.	1.88	0.93	1.85	1.04	2.20	1.29	3.48	1.31	151.46	78.72
Odisha	1.40	1.23	1.39	1.24	1.38	1.23	1.36	1.15	76.01	76.01
Chhattisgarh	0.55	0.30	0.51	0.31	0.64	0.30	0.69	0.46	511.11	511.11
Tamil Nadu	0.53	0.50	0.60	0.58	0.60	0.58	0.59	0.44	176.00	130.80
Bihar	0.22	0.37	0.22	0.37	0.22	0.32	0.22	0.34	248.36	332.03
Total Above	38.74	27.99	38.58	31.33	37.04	24.24	53.24	45.29	5.78	7.71
Others	0.50	0.45	0.46	0.41	0.42	0.34	0.63	0.70	345.68	605.54
All-India	39.24	28.44	39.04	31.74	37.46	24.58	53.87	45.99	5.69	7.61

Source: Normal: DES, (Ave. of 2011-12 to -2015-16), *IIIrd Advance Estimates of Production 2016-17

(Table-II): State-wise area production of urdbean

{Area: lakh ha, Production -lakh tonnes}

State	Nor	mal	2013	3-14	201	5-16	2010	6-17*	Prod. %	change
	Area	Prod.	Area	Prod.	Area	Prod.	Area	Prod.	ove	er
								•	Normal	2015-
										16
Madhya Pradesh	7.25	3.45	6.02	2.26	9.35	5.17	11.68	7.71	123.64	49.17
Andhra Pradesh	3.78	3.17	2.65	2.30	4.56	4.11	5.20	3.84	21.15	-6.57
Uttar Pradesh	5.70	3.09	5.42	2.48	6.17	2.44	6.42	3.48	12.65	42.87
Tamil Nadu	3.30	2.40	3.65	3.11	3.95	2.64	4.49	3.72	54.85	40.80
Maharashtra	3.24	1.64	3.34	2.06	2.86	0.61	4.45	2.50	52.25	309.84
Rajasthan	2.34	1.11	1.96	0.71	2.99	1.15	3.90	2.50	125.48	118.17
Jharkhand	0.94	0.78	0.94	0.88	0.95	0.72	1.52	1.40	77.90	93.72
Gujarat	0.84	0.54	0.91	0.55	0.64	0.38	1.99	1.21	124.91	218.42
West Bengal	0.74	0.49	1.14	0.63	0.74	0.55	0.76	0.55	13.70	1.47
Karnataka	0.93	0.37	1.02	0.50	0.91	0.25	0.88	0.43	16.85	72.00
Chhattisgarh	1.02	0.31	1.08	0.32	1.01	0.30	0.99	0.32	2.26	4.28
Total above	30.09	17.35	28.13	15.80	34.13	18.31	42.27	27.65	59.43	51.03
Other	2.55	1.37	2.49	1.19	2.11	1.14	2.66	1.61	17.24	40.90
All India	32.64	18.72	30.62	16.99	36.24	19.45	44.93	29.26	56.34	50.43

Source: Normal: DES, (Ave. of 2011-12 to -2015-16), *IIIrd Advance Estimates of Production 2016-17.

(Table- III): State-wise area production of mungbean

{Area: lakh ha, Production -lakh tonnes}

State	No	rmal	201	2013-14		5-16	201	6-17*	Prod. %	
									chang	ge over
	Area	Prod.	Area	Prod.	Area	Prod.	Area	Prod.		
									Normal	2015-16
Rajasthan	10.70	4.68	10.20	3.91	13.64	5.97	15.83	7.52	60.76	25.98
Maharashtra	3.97	1.66	4.31	2.08	3.66	0.69	5.15	2.68	61.86	288.84
Tamil Nadu	1.89	1.15	1.95	1.51	2.39	1.25	1.63	1.16	1.06	-6.95
Andhra	1.56	1.12	1.34	1.17	2.12	1.37	1.62	0.83	-25.84	-39.42
Pradesh										
Bihar	1.61	0.98	1.55	1.05	1.69	0.94	1.64	1.00	1.11	5.45
Madhya	2.10	0.93	3.16	1.46	2.95	1.31	2.94	1.39	49.37	5.81
Pradesh										
Odisha	2.66	0.86	2.52	0.89	2.90	0.85	2.66	0.85	-0.37	0.06
Gujarat	1.62	0.83	1.83	1.06	1.29	0.67	1.80	0.86	3.01	28.36
Telangana	1.24	0.69	1.26	0.53	1.11	0.56	1.48	0.88	27.76	57.14
Karnataka	2.80	0.61	3.20	0.81	3.48	0.44	4.14	1.15	89.83	161.96
Uttar Pradesh	0.86	0.47	0.79	0.39	1.11	0.51	1.09	0.59	26.61	15.69
Total above	31.00	13.97	32.12	14.87	36.34	14.57	39.98	18.91	35.37	29.83
Other	1.66	1.07	1.71	1.19	1.94	1.36	3.07	1.78	66.27	30.92
All India	32.67	15.04	33.83	16.05	38.28	15.93	43.05	20.70	37.57	29.93

Source: Normal: DES, (Ave. of 2011-12 to -2015-16), *IIIrd Advance Estimates of Production 2016-17.

(Table-IV): State-wise area production of gram

{Area: lakh ha, Production -lakh tonnes}

State	Nor	mal	201	3-14	201	5-16	2010	6-17*	Prod. %	6 change
	Area	Prod.	Area	Prod.	Area	Prod.	Area	Prod.	0.0	er
									Normal	2015-16
M.P.	30.60	32.11	31.60	32.99	30.17	32.65	32.22	35.49	10.53	8.70
Maharashtra	13.71	11.36	18.20	16.22	14.41	7.31	18.95	16.48	45.07	125.44
Rajasthan	15.30	12.98	19.24	16.40	9.42	8.03	14.86	13.82	6.47	72.10
Karnataka	9.23	6.22	9.46	7.16	13.72	8.97	10.25	3.90	-37.30	-56.52
Andhra Pradesh	4.68	5.21	4.72	5.97	4.71	5.00	3.92	4.34	-16.70	-13.20
Chhattisgarh	2.64	2.54	2.77	2.13	3.03	2.19	2.93	3.01	18.50	37.44
Uttar Pradesh	5.77	5.47	5.77	4.75	2.68	2.16	5.62	6.32	15.54	192.59
Jharkhand	1.30	1.48	1.56	1.82	1.64	1.73	1.86	2.19	47.97	26.59
Gujarat	1.99	2.30	2.47	3.09	1.15	1.53	1.70	2.10	-8.70	37.25
Telangana	0.96	1.42	1.14	2.46	0.70	0.49	1.01	1.24	-12.68	153.06
Bihar	0.59	0.70	0.61	0.70	0.61	0.60	0.60	0.59	-15.71	-1.67
West Bengal	0.24	0.28	0.25	0.29	0.31	0.37	0.30	0.33	17.86	-10.81
Total of above	87.01	82.07	97.79	93.98	82.55	71.03	94.22	89.81	9.43	26.44
All-India	88.37	83.23	99.27	95.26	83.49	71.69	95.39	90.75	9.04	26.59

Source: Normal: DES, (Ave. of 2011-12 to -2015-16), *IIIrd Advance Estimates of Production 2016-17.

(Table-V): State-wise area production of lentil

{Area: lakh ha, Production -lakh tonnes}

State	Nor	mal	201	3-14	201	5-16	2010	6-17*	Pro	d. %
_									chang	ge over
	Area	Prod.	Area	Prod.	Area	Prod.	Area	Prod.	Normal	2015-
										16
UP	5.08	3.8	4.49	3.10	3.35	2.38	6.63	5.01	31.84	110.50
MP	5.85	2.99	5.30	3.38	5.46	3.92	5.86	3.48	16.39	-11.31
Bihar	1.83	1.92	1.54	1.96	1.51	1.40	2.13	2.28	18.75	62.35
WB	0.62	0.56	0.65	0.63	0.95	0.94	1.13	1.04	85.71	10.64
Jharkhand	0.39	0.36	0.44	0.37	0.15	0.12				
Rajasthan	0.36	0.35	0.34	0.29	0.74	0.64				
Assam	0.27	0.17	0.30	0.22	0.28	0.20	0.33	0.22	29.41	11.99
Total above	14.40	10.15	13.07	9.95	12.45	9.60	16.08	12.03	18.52	25.31
Others	0.39	0.23	0.34	0.22	0.31	0.16	0.57	0.25	8.70	57.13
All India	14.79	10.38	13.41	10.17	12.76	9.76	16.65	12.28	18.30	25.83

Source: Normal: DES, (Ave. of 2011-12 to -2015-16), *IIIrd Advance Estimates of Production 2016-17

14.2 Availability Status: Total Pulses & Crop-Wise (2013-14 To 2016-17)

Crop-wise availability of Pigeonpea, Chickpea, Lentil, Mungbean and Urdbean based on domestic production, import and export is summarized under *Table 14.2*. During 2016-17, it is evident from table that the domestic availability of pulses has increased by 57% in total pulses, 89% in tur, 32% in chickpea, and 35% in urd and mung over the previous crop year 2015-16.

(Table-14.2): Import, Export and Availability

(Unit: Lakh Tons)

Year	Domestic Production	Import	Export	Total Availability
Pigeonpea				
2013-14	31.74	4.63	0.04	36.33
2014-15	28.07	5.74	0.09	33.72
2015-16	24.58	4.66	0.001	29.24
2016-17	48.73	7.03	0.12	55.54
Chickpea				
2013-14	95.26	2.76	3.33	94.69
2014-15	73.32	4.19	1.90	75.61
2015-16	70.58	10.31	2.17	78.72
2016-17	93.78	10.81	0.88	103.71
Lentil				
2013-14	10.18	7.09	0.01	17.26
2014-15	10.35	8.16	0.08	18.43
2015-16	9.76	12.60	0.12	22.24
2016-17	12.24	8.29	0.16	20.37
Moongbean /Uı	rdbean			
2013-14	33.04	6.24	0.02	39.26
2014-15	34.63	6.23	0.04	40.82
2015-16	35.38	5.82	0.06	41.14
2016-17	49.97	5.75	0.11	55.61
Total Pulses				
2013-14	192.53	36.44	3.46	225.51
2014-15	171.52	45.85	2.22	215.15
2015-16	163.48	57.98	2.56	218.90
2016-17	231.31	112.82	1.37	342.46

Source: GOI, MoA &FW, Min. of Commerce& Industry; Domestic production Final Est.and Import & Export 2016-17 (16th Feb., 2018).

14.3 Market Prices/Rates and Arrivals

Crop-wise market prices and arrivals of Pigeonpea, Urdbean, Mungbean, Gram and Lentil, worked out relative change over the period in terms of *arrivals* annually and *prices/rates*, monthly during 2015-16 and 2016-17 and, are summarized.

It is evident from table that the arrivals of pulses has increased by 72% in tur, 58% in urd and 29% in mung and 6% in lentil over the previous crop year 2015-16. It is also observed that the prices/rates are decreased over the period during the month of July by 53% in tur 48% in urd, 23% in mung and 34% in lentil.

(Table-I): State-wise market rates and arrivals of pigeonpea

State	Market	Rate	%	Market	Arrivals	% Change
	July, 2016	July,	Change	2016-17	2015-16	over
		2017	over			
Maharashtra	8644	3689	-57.32	799.71	535.40	49.37
Karnataka	9392	4174	-55.56	416.41	197.23	111.14
Madhya Pradesh	6938	3924	-43.44	321.31	174.53	84.10
Gujarat	7835	3262	-58.37	103.99	38.99	166.73
Telangana	8357	3722	-55.46	102.38	5.79	1668.56
Uttar Pradesh	8849	3999	-54.81	80.29	87.41	-8.14
Rajasthan	7588	4063	-46.45	24.21	14.74	64.23
West Bengal	12199	8342	-31.62	15.71	18.73	-16.16
Chattisgarh	7802	3944	-49.45	9.87	8.77	12.52
Assam	12381	6792	-45.14	7.82	10.96	-28.60
Andhra Pradesh	7799	3810	-51.15	6.26	1.02	512.32
Jharkhand	12901	6072	-52.93	4.68	3.86	21.18
Uttrakhand				2.69	1.11	142.34
Kerala	12919	7601	-41.16	1.42	0.60	134.93
Manipur	16361			1.29	1.37	-6.05
Punjab		2560		0.92	4.41	-79.12
NCT of Delhi	6267	2487	-60.32	0.90	0.81	11.91
Haryana				0.17	0.13	34.11
Tamil Nadu				0.08	1.16	-93.56
Orissa				0.00	0.20	-98.01
Average/Total	9749	4563	-53.20	1900.13	1107.23	71.61

Source: GOI, MoA &FW, DMI, Agmarknet

(Table-II): State-wise market rates and arrivals of urdbean

(Rate in Rs./Qtls; Arrivals-000 Tonnes)

State	Marke	t Rate	% Change	Market	Arrivals	% Change
	July,	July,	over	2016-17	2015-16	over
	2016	2017				
Madhya Pradesh	8798	4109	-53.30	349.82	284.01	23.17
Maharashtra	14190	3912	-72.43	123.54	36.16	241.65
Rajasthan	9943	3686	-62.93	113.68	79.98	42.15
Uttar Pradesh	10312	4977	-51.74	93.66	58.23	60.85
Gujarat	10500	3954	-62.34	82.00	18.16	351.53
Assam				77.39	29.90	158.81
Tamil Nadu	9206	5422	-41.10	43.38	22.08	96.51
Karnataka	12512	7660	-38.78	36.38	23.61	54.05
West Bengal	12841	7040	-45.18	23.11	35.71	-35.30
Andhra Pradesh	9246	5127	-44.55	5.00	5.20	-3.73
Kerala	15327	9196	-40.00	2.72	2.47	10.15
Pondicherry	7316	4088	-44.12	1.98	1.95	1.23
Telangana	7046	4050	-42.52	1.91	0.96	99.69
Chattisgarh	9935	4413	-55.58	1.74	2.37	-26.38
Orissa	8024	7515	-6.34	1.54	4.56	-66.15
Manipur	15391		-100.00	1.35	1.40	-3.51
Uttrakhand	7800	5916	-24.15	0.67	0.75	-10.47
Haryana				0.21	0.27	-21.48
Jharkhand		6272		0.12	0.00	5700.00
Punjab	9362		-100.00	0.02	0.05	-68.75
Average/Total	10456	5459	-47.79	960.22	607.81	57.98

Source: GOI, MoA &FW, DMI, Agmarknet

(Table-III): State-wise market rates and arrivals of mungbean

(Rate in Rs./Qtls; Arrivals-000 Tonnes)

G	Market Rate %					ivais-000 Tonnes)
State	Marke		- %		Arrivals	% Change
	July,	July,	Change	2016-17	2015-16	over
	2016	2017	over			
Rajasthan	5457	4347	-20.34	320.41	268.93	19.14
Madhya	5058	5142	1.66	244.12	165.53	47.47
Pradesh						
Karnataka	6263	5179	-17.31	127.37	71.27	78.70
Maharashtra	6242	4574	-26.72	39.34	22.22	77.02
Uttar Pradesh	5938	3872	-34.79	33.46	39.03	-14.28
Assam	8758	6774	-22.65	22.19	25.08	-11.55
Gujarat	5717	4592	-19.68	20.19	30.25	-33.24
Telangana	5015	3899	-22.25	18.35	9.93	84.69
West Bengal	10288	8386	-18.49	16.61	20.19	-17.72
Haryana	3800	5000	31.58	5.95	0.08	7245.68
Kerala	9011	7479	-17.00	2.57	2.69	-4.42
Manipur	13348			1.19	1.29	-7.30
Orissa	6242	5427	-13.06	1.11	3.32	-66.69
Andhra	5210	5564	6.79	0.90	1.69	-46.71
Pradesh						
Tamil Nadu	5070	4029	-20.53	0.86	1.56	-45.13
Pondicherry	5267	4214	-19.99	0.49	0.54	-10.85
Uttrakhand	5200	5644	8.54	0.44	0.91	-51.38
Jharkhand	9315	6341	-31.93	0.27	0.07	300.00
Punjab	5659		-100.00	0.10	0.08	27.85
Chattisgarh	10065	4200	-58.27	0.05	0.03	64.29
NCT of Delhi	4441	3214	-27.63	0.01	0.05	-79.63
A&N Island				0.01	0.01	-16.67
Mizoram				0.00	0.01	-87.50
Average/Total	6732	5151	-23.47	855.96	664.77	28.76

Source: GOI, MoA &FW, DMI, Agmarknet

(Table-IV): State-wise market rates and arrivals of lentil

(Rate in Rs./Qtls; Arrivals-000 Tonnes)

State	Marke	t Rate	%	Market	Arrivals	% Change
	July, 2016	July, 2017	Change	2016-17	2015-16	over
			over			
Madhya Pradesh	5688	3310	-41.80	279145.64	297636.97	-6.21
Uttar Pradesh	6674	4122	-38.24	122626.96	68523.86	78.96
Assam	8882	6164	-30.59	40626.30	36813.40	10.36
West Bengal	9620	7675	-20.22	28538.25	40004.41	-28.66
Rajasthan	6180	3329	-46.13	10122.80	9242.90	9.52
Maharashtra	7138	5336	-25.24	2295.00	2616.00	-12.27
Jharkhand	7877	5908	-25.00	2236.58	845.60	164.50
Chattisgarh	5159	3155	-38.84	2218.73	2495.60	-11.09
Manipur	11412			1350.19	1373.40	-1.69
Uttrakhand				796.20	811.80	-1.92
Gujarat				36.40	5.10	613.73
Haryana				8.50	6.60	28.79
Telangana				7.40		
Kerala	10433	7749	-25.73	3.70	1.21	205.79
Bihar					52.10	
Orissa					16.00	
Punjab					19.45	
Average/Total	7906	5194	-34.30	490012.65	460464.40	6.42

Source: GOI, MoA &FW, DMI, Agmarknet

(Table-V): State-wise market arrivals of gram

(Unit- Arrivals in Lakh Tonnes)

State	July 01, 2015 to	July 01,2016 To	% Change over 2015-16
	May 31, 2016	May 31,2017	2013-10
Andhra Pradesh	0.200	0.040	-79.86
Assam	0.136	0.131	-3.33
Chattisgarh	0.268	0.156	-41.84
Gujarat	0.339	0.520	53.30
Haryana	0.028	0.009	-68.79
Jharkhand	0.009	0.001	-94.26
Karnataka	0.872	0.847	-2.93
Kerala	0.042	0.023	-44.66
Madhya Pradesh	7.050	8.082	14.64
Maharashtra	3.173	3.635	14.57
Manipur	0.013	0.012	-5.58
NCT of Delhi	0.001	0.001	59.19
Orissa	0.000	0.000	-91.71
Punjab	0.030	0.000	-99.90
Rajasthan	1.117	2.381	113.23
Tamil Nadu	0.002	0.000	-91.91
Telangana	0.002	0.015	779.98
Tripura	0.000	0.003	-
Uttar Pradesh	1.838	1.957	6.46
Uttrakhand	0.016	0.006	-60.14
West Bengal	0.063	0.074	17.11
Total	15.199	17.893	17.72

Source: AGMARKNET

Note: During 2015-16 the AGMARKNET data for Gram was captured under two heads i.e. Bengal Gram and Big Gram and the same was merged and reported in a single commodity name: Bengalgram (Gram) w.e.f 11.03.2016.

(Table-VI): State-wise market arrivals and prices of gram

(Rate in Rs./Qtls; Arrivals-lakh Tonnes)

State		Mark	et Rates			Mark	et Arrivals	3
	2014	2015	2016	2017	2014	2015	2016	2017
MSP	3100	3175	3425	4000				
M.P.	2673	4241	5623	5370	0.967	0.379	1.558	1.942
Maharashtra	2519	4287	5587	5420	0.459	0.311	0.342	0.559
Rajasthan	2710	4201	5566	5229	1.190	0.322	0.252	0.671
Karnataka	2661	4674	5687	6367	0.132	0.068	0.065	0.091
Andhra Pradesh	2824	3712	5061	5954	0.005	0.012	0.000	0.007
Chhattisgarh	2661	4417	5510	5451	0.013	0.011	0.050	0.028
Uttar Pradesh	3350	4218	5735	5793	0.241	0.211	0.232	0.262
Jharkhand	3580	5528	6208		0.017	0.004	0.000	0.000
Gujarat	2698	4412	5719	5352	0.067	0.038	0.041	0.069
Telangana	2474	4229	4390	5756	0.000	0.000	0.000	0.001
Bihar					0.000		0.000	0.000
West Bengal			11250	9100	0.000		0.016	0.008
All-India	3312	4471	6135	6468	3.104	1.386	2.588	3.664

Source: AGMARKNET

- The all India market arrival of gram during May, 2016 was 2.59 lakh tons which was about 3.61% of the total gram production during 2015-16 whereas during May, 2017 it is 3.66 lakh tons which is about 4.04 % of total gram production during 2016-17 which is only 0.43 % higher than the last year.
- The market arrival during May, 2014 was 3.10 lakh tons which is only 3.26% of the total gram production during 2013-14, the ever highest gram production year with 95.25 lakh tons of production of Chana. It reveals that the arrivals percentage of gram during the current year (May 1st-31st, 2017) is higher by 0.78 % than the ever highest gram production year of 2013-14 during corresponding period (May 01st-31st, 2014).
- The trend of monthly wholesale prices of gram during last five years exhibit that the market rates are directly related with the production, the decreasing trend of prices observed during the 2014 and current year 2017 due to record production of gram during 2013-14 & 2016-17. The increasing price trend observed during 2015 and 2016 in view of less production during 2014-15 & 2015-16. During the current year wholesale price of gram during the month of January to May, 2017 show decreasing trend due to increasing trend of market arrivals with 17.72% higher than the last year in the same period.


14.4 Factors attributing to lower market prices (Below MSP)

- During 2016-17, as a result of significant increase in the area coverage and productivity of all major Pulses, total production of pulses is estimated at 22.95 million tonnes which is higher by 3.70 million tonnes (>19 %) than the earlier record production of 19.25 million tonnes achieved during 2013-14.
- The production of total pulses during the year under report is also higher by 4.77 million tonnes (>27%) than its Five yearsø average, as also higher by 6.60 million tonnes (>40%) over the last year i.e. 16.35 million tonnes.
- The availability status of the pulses during the current year, both under total pulses and individual crop category, has increased considerably. Once the availability is sufficient, the prices are bound to be comparatively low.
- The market arrival status of the pulses during the current year under individual crop category has increased significantly. Once the supply increased over its demand, prices/rates are declined.
- It is evident from the table depicting the increasing trend of availability of individual pulses as well as total pulses, the increased availability, may be a major factor for low ruling prices during the current month/year.
- Currency demonetization and implementation of GST may likely to impact the rotation of money under trading, resultantly poor holding of stocks in relation to the capacity of the traders/ processors.

- Imposing of 5% GST on *branded dals* and relaxing *non-branded dals* i.e., 0% GST, may also be attributed to comparatively poor procurement by the traders /processors/millers, fearing the assured profit-margin over the investment.
- Except, the big processors/traders, the other small scale entrepreneurs may be reluctant and unwilling to put the *brands*. During the course of interactions/ discussions with the millers, it is given to understand that the traders lobby anticipates the policy decision of the Government to open the export of this commodity.
- It is relevant to mention that the Lentil is exported to Myanmar (>35 %), USA (> 25 %), Kuwait (> 7 %) and Bhutan and Singapore (approx. 6 %). Similarly, tur is exported to Nepal (> 78 %), Canada (> 19 %) and Israel (approx 2%). Mung/Urd is also exported to USA (> 49 % of total export of this commodity) followed by Sri lanka and Canada (> 7 %).
- It should also be noted that the MSP procurement under PSF/PSS are for FAQ Grade, fetching MSP rates. Whereas, the lower prevailing rates of these pulses in the market are for ungraded/below FAQ grade of pulses.
- It is pertinent to record that for a stable remunerative price support under pulses, the state¢s own initiatives on procurement always pays dividends. In Karnataka, > 80000 qtls of tur dal have been procured during 2016-17 here the open market rates have been comparatively much better than the other states having no such provision.
- As per Agmarknet portal, the mandi arrivals of Bengal Gram (Chana) for all the states from 01.07.2016 to 31.05.2017 are 17.72% higher as compared to the corresponding period of last year (2015-16).
- The arrivals during the current year/month are higher than the corresponding year/month of the last year 2016. The probable reasons of higher arrival and prevailing market prices may be attributed to the followings:
- The higher MSP regime in pulses (Gram) is paying dividends to pulse growers in fetching remunerative prices of their produce. Currently price is above the MSP and is sustaining above MSP, because of procurement by the Government in major gram producing states.
- There is no carry forward stock (31st March, 2017) with the farmers, millers, stockist, traders and also the public sector agencies like NAFED and the produce being sold both at organized and non-organized trading routes.
- Holding of stocks by the farmers with access to information network, enhanced holding capacities anticipating further better prices correlating the previous year price regime.
- Farmersø enhanced access to get finance through sale proceeds of wheat crop and the KCCs and generally not constrained to distress sale.
- The price trends have been decreasing from March/April onwards during the current year as compared to the corresponding period of last year. It is an indicative that production is more.
- The status market arrivals during the current year is better than the last year due to excess production of gram during the current year.

the,	Generally, of the total production/ quantity of the gram commodity i.e., 42-43 percent of the, Total availability (Total production + Import - Export-Change in stock over year-NSSO based consumption) is retained as stock and the remaining 56-58 percent of the total available quantity remains under trading.								

SEED PRODUCTION



Chapter-15

Seed Production

15.1 Importance

Seed is the key input in pulse crop cultivation and vital in speeding and sustaining the crop productivity. The quality of seed alone is known to 10-15% increase in the total production of any crop. In the absence of quality seed, the inputs like fertilizer, water, pesticides etc., do not pay the desirable return. Lack of quality seed continues to be one of the greatest hurdles in reducing the vast yield gap between improved practices (FLD), farmersøpractice and statesøaverage yield. Concerted efforts and proper planning along with realistic execution the seed production programme are required to produce the quality seed of improved varieties insufficient quantities to phase out the old seed of absolete non-descript varieties.

15.2 Class of Seed and their production

i) Nucleus Seed

- Basic seed of variety should be sown in optimum area approximately 200 m². Field should qualify the norms in terms of topography, moisture availability and fertility. Recommended spacing (plant to plant and row to row) should be maintained through dibbling or thick rowing, followed by thinning.
- Select 500-1000 plant which exactly confirms the varietal descriptors. Number of plants to be selected will depend upon the seed production ability of individual plant *i.e.* yield per plant, if yield per plant is higher less number of plants selected and tagged.
- The tagged plants should be harvested separately. Seed of individual plants should be carefully examined and if the seed/plant produce is not conforming to the discriptors of the variety, the seed lots produce of individual plants should be rejected.
- The seed collected should be dried, treated and stored.
- During next cropping season, the individual plant progenies should be grown in standard and homogenous field. Row to row spacing should be wide than the normal recommendation of the crop. The main objective of spaced planting is to ensure genetic purity, otherwise the higher productivity per unit area. Here the row length may vary from one to three meters, depending upon the quantity of produce of individual plant.
- Due care should be taken for all the agronomic practices of the crop to ensure high seed to seed ratio.
- Individual plant progenies should be regularly visited by breeder, right from germination to harvesting.
- If any individual plant is not true to type and /or sister progeny are showing disease incidence, plant should be completely removed from the field, besides entire off type/diseased progeny should be discarded completely.
- From the remaining progenies, 500-1000 plants should be tagged for next year planting of single plant progenies.
- Individual plant should be harvested separately, as during the previous season and necessary steps, as enumerated above, should be strictly followed for next year planting, as well.
- After harvesting these 500-1000 plants, the individual plant progenies should be harvested separately.

- The seed lot of individual progenies should be examined for size, shape colour etc. of the seed. Any progeny exhibiting mixture or deviating from the seed descriptors of the original variety or sister progeny should be discarded.
- Remaining progenies left after rejection both at preand post harvest stage should be bulked, this bulk produce of selected progenies (bulk produce of 400 progenies out of 500 plants) is known as Nucleus Seed.
- The nucleus seed is used for production of breeder seed. Special care must be given to this seed during storage.

ii) Breeder Seed

All stake holders who deal in seed viz. NSC, SDAs, SAUs, SSC, Seed Grower Societies and private sectors, place their breeder seed indent to Seed Division, Govt. of India, DAC& FW. The indent, in compiled form, is given to ICAR who organizes. Breeder Seed Production (BSP) of various varieties of different crops through ICAR Institutes, SAUs, and AICRPs other organizations like NSC, etc. The different breeder seed production proformae are enumerated below:

BSP-1: In view of indents received, Project Coordinator unit formulate BSP-1 after detailed discussion in concerned crop Annual Group Meet group meet. The BSP-I proforma issued by PI/PC accounts for crop, variety, name of breeder to whom BSP is allocated, DAC & FW indent allocation and indentors.

BSP-II: After receiving the BSP-I proforma from PC unit, the breeder of SAUs/ICAR institutes sow the *nucleus seed/basic seed* for breeder seed production.BSP-II proforma is submitted by concerned crop breeder to PC unit after compilation of sowing of breeder seed production plot.

BSP-III: The BSP-III proforma is submitted by the breeder to PC unit after completion of monitoring by monitoring team. The team comprises breeders; in charge National Seed Programme, NSC representative and officer from State Seed Certification Agency.

BSP-IV: This proforma is issued by breeders after harvesting, threshing, cleaning and grading of breeder seed. It contains information on actual breeder seed of different varieties produced by the concerned crop breeder. On the basis of this proforma, seed division of DAC&FW arranges lifting of the breeder seeds by indentors.

BSP-V: After lifting of breeder seed by indentors, this proforma is submitted by breeders to PC unit and contains information on lifting of breeder seed by indentors.

iii) Certified Foundation Seed

This is the seed which is certified by a State Seed Certification agency notified under section 8 of Indian Seed Act 1966 or by any other foreign certification agency provided that the agency is recognized by Govt. of India through notification in official gazette. The certified seed consist of two classes:

- Certified Foundation Seed Stage I and II: CFS is the progeny of breeder seed or certified foundation seed it self. When seed is progeny of breeder seed, it is called foundation seed stage I, while it is called foundation seed stage II when it is the progeny of certified foundation seed stage I it is important to note that only certified foundation seed stage I can be multiplied to generate certified foundation seed stage II.
- Certified foundation seed stage II cannot be used to produce foundation seed; it can only be used to produce certified seed. The minimum seed standard for both foundation seed stage I and foundation seed Stage II are similar unless otherwise prescribed.

 Production of foundation seed stage II is undertaken only when it is clearly stated by Seed Certification Agency that the breeder seed of a particular variety is in short supply and Stage II foundation seed has to be produce to meet the seed demand. Bags of foundation seed carry white coloured tags.

iv) Certified Seed

This seed is progeny of foundation seed and it is produced under conformity of specific genetic identity and purity standard as prescribed for the crop being certified.

The certified seed can be progeny of certified seed provided this multiplication does not exceed three generation beyond foundation seed stage-I. Certified seed produced from foundation seed is called certified seed stage I while that produced by multiplication of certified seed itself is called certified seed stage II. Certified seed stage II can not further be used for multiplication.

The tag of certified seed is of blue colour (Shade ISI No. 104 *azure blue*) besides carrying all relevant information about the certified seed inside the bag.

15.3 Seed Replacement: The proactive efforts and various seed related programmes after 2014-15 has resulted to increase in the SRR in most of the pulses. The targeted SRR of 33% for self-pollinated crops was not only achieved, but also increased with a number of HYVs in the seed-chain. The details SRR is given below (*Table-15.1*).

(Table-15.1): Crop-wise & Year-wise SRR

(SRR-Per cent)

Crop	2014-15	2015-16	2016-17
Urd	30	34	38
Moong	24	31	34
Arhar	41	45	48
Pea	34	30	30
Gram	25	28	32
Lentil	31	27	35

Source: Seed Division, Min. of Agri. & FW (DAC&FW)



15.4 Seed Requirement

To achieve the targeted 36%, 44%, 52% & 60% Seed Replacement Rate, the requirement of breeder, foundation and certified seed by the end of 2030 is as under:

(Table-15.2): Seed Requirement

(Quantity in Qtl.)

Crop	Normal		Certifi	ed seed			Foundat	ion seed		Breeder seed			
	Area	2018-19 (36%)	2022-23 (44%)	2026-27 (52%)	2030-31 (60%)	2018-19	2022-23	2026-27	2030-31	2018-19	2022- 23	2026- 27	2030- 31
Arhar	41.91	301.72	368.76	435.81	502.86	7.54	9.22	10.90	12.57	0.15	0.18	0.22	0.25
Urdbean	27.00	194.37	237.56	280.76	323.95	6.48	7.92	9.36	10.80	0.32	0.40	0.47	0.54
Mungbean	24.93	179.48	219.37	259.25	299.13	5.98	7.31	8.64	9.97	0.30	0.37	0.43	0.50
Other Kharif	18.10	130.33	159.30	188.26	217.22	4.34	5.31	6.28	7.24	0.22	0.27	0.31	0.36
Tot. Kharif	111.93	805.90	984.99	1164.08	1343.17	24.35	29.76	35.17	40.58	0.99	1.21	1.43	1.65
Gram	89.45	2576.19	3148.68	3721.16	4293.65	171.75	209.91	248.08	286.24	17.17	20.99	24.81	28.62
Lentil	13.94	125.47	153.35	181.23	209.12	4.18	5.11	6.04	6.97	0.21	0.26	0.30	0.35
Urdbean	8.13	58.56	71.58	84.59	97.61	1.95	2.39	2.82	3.25	0.10	0.12	0.14	0.16
Mungbean	9.62	69.29	84.68	100.08	115.48	2.31	2.82	3.34	3.85	0.12	0.14	0.17	0.19
Other Rabi	19.69	283.59	346.61	409.64	472.66	14.18	17.33	20.48	23.63	0.71	0.87	1.02	1.18
Total Rabi	140.84	3113.10	3804.90	4496.70	5188.50	194.37	237.56	280.76	323.95	18.31	22.37	26.44	30.51
Total Pulses	252.77	3919.01	4789.90	5660.79	6531.68	218.72	267.32	315.93	364.53	19.30	23.59	27.87	32.16

15.5 Breeder seed production under new varieties

Breeder Seed

During 2014-15 Breeder seed production and indent was 10,910 qtls and 9702 qtls respectively. During 2016-17 both the production and indent increased by 40% and 37% (15242 qtls/13236 qtls) over the base year 2014-15. New varieties of pulses put to seed chain. The crop-wise production and indent of breeder seed is given as under (*Table-15.3*).

(Table- 15.3): Breeder seed production and indent

(Quantity-qtls)

Crop	2014	-15	2015	-16	2016	-17	2017	'-18
	Indent	Prod.	Indent	Prod.	Indent	Prod.	Indent	Prod.
Pigeonpea	390	653	226	734	308	653	351	735
Chickpea	6742	7464	7184	7630	10119	11174		
Mungbean	932	857	702	686	811	890	970	912
Urdbean	424	485	424	296	454	513	457	364
Lentil	259	312	449	474	467	535		
Fieldpea	464	637	1215	759	611	777		
Horsegram	8	4	10	2	17	21	25	41
Mothbean	95	36	59	23	62	66	25	42
Rajmash	4	88	2	7	3	14		
Lathyrus	4	88	9	14	115	166		
Clusterbean	341	272	269	257	227	289	231	306
Cowpea	39	14	168	26	42	144	19	31
Total	9702	10910	10717	10908	13236	15242	2078	2431

Source: ICAR-IIPR, Kanpur, U.P.




Fig.-15.2: Breeder Seed Production vs Indent: Total Pulses




Fig.-15.3: Breeder Seed Production: Kharif Pulses




Fig.-15.4: Breeder Seed Production: Rabi Pulses

• The seed production subsidy was one of the major initiatives during 2016-17. The quality pulse seed produced by various public sector/central agencies has been about 2.65 lakh qtls during 2016-17 and >6.00 lakhqtls in 2017-18. The new/promising varieties were made available to the farmers. The agency-wise quality seed is given below (Table-15.4).

(Table-15.4): Certified seed production programme under NFSM-Pulses

(Quantity-qtls)

Agency		2016-17		2017-18			
	Kharif	Rabi	Total	Kharif	Rabi	Summer	Total
NSC	35413	196720	232133	65000	269000	47000	381000
HIL	2681	23016	25697	22000	69000	3000	94000
KRIBHCO		7140	7140	1650	6060		7710
NAFED				48400	58900		107300
IFFDC				3400	16600		20000
Total	38095	226876	264971	140450	419560	50000	610010

Source: NFSM Cell, Min. of Agri. & FW (DAC&FW)




Fig.- 15.5: Availability of Certified Seeds of Pulses Enhanced

- As a result during 2017-18, enhanced availability of > 35.00 lakh qtls certified seeds of improved varieties which is >6.75 lakh qtls (23%) higher the availability over the base year 2014-15. Certified seed availability in gram was >2.00 lakh qtls, over it@s requirement during 2017-18, followed by urd and mung.
- During 2018-19 the targeted certified seed production is 31.22 lakh quintals.

(Table-15.5): All India: Crop-wise requirement and availability of certified seed

(Quantity- lakh qtls.)

Crop	2014-15 (Base Year)		2015-16		2016-17		2017-18		Change Over
	R	A	R	A	R	A	R	A	(+/-) 2017-18
Gram	16.11	15.72	18.14	14.86	17.65	16.01	17.16	19.27	2.11
Lentil	1.79	1.38	1.30	1.06	1.47	1.15	1.37	1.36	-0.01
Peas	1.96	1.57	2.12	1.83	2.67	2.91	2.39	2.36	-0.03
Urd	2.68	3.31	2.62	2.71	2.67	2.9	2.74	3.74	1.00
Moong	2.79	3.31	2.87	3.23	2.68	3.27	2.41	3.14	0.73
Arhar	2.64	2.78	2.51	2.72	2.71	2.97	3.31	3.81	0.50
Others	0.92	0.81	0.92	0.84	1.81	2.07	1.88	1.96	0.08
Pulses Total	28.88	28.87	30.49	27.24	31.66	31.28	31.26	35.64	4.38

Source: Seed Division, Min. of Agri. & FW (DAC&FW), R- Requirement, A – Availability



Fig.-15.6: Availability of Certified Seed

15.6 New Initiatives under NFSM-Pulses

õThe Committee for Monitoring Actions/ Strategy for Increasing Pulses Production". To enhance production of pulses in the country, availability of quality seeds SRR of latest/ promising varieties VRR and adoption of recommended technologies (TOT) has been viewed a major bottleneck. The committee under the Chairmanship of Dr. Ashok Dalwai, Additional Secretary, Govt. of India, following strategic interventions to address the seed sector during 2016-17 have neen initiated;

15.6.1 Enhancing Breeder Seed Production

As a major initiative to address SRR/VRR constraints with budgetary allocation of Rs. 20.39 crore EBSP programme has been started in projectile mode for a period of 03 years (2016-17 to 2018-19) in 08 states with 12 centers at Rajasthan (ARS, Kota/RARI-Durgapur), Bihar (BAU, Sabour), Maharashtra (ARS-Badnapur/MPKVV-Rahuri), Madhya Pradesh (JNKVV, Jabalapur/RVSKVV, Gwalior & IIPR-RS

Phanda), Odisha (ARS, Berhampur), Uttar Pradesh (ICAR, IIPR, Kanpur), Andhra Pradesh (ARS-Lam), Karnataka (UAS, Dharwad).

Crop-wise/Centre-wise targets and achievement for quality seed production and infrastructure created at each location is given below (*Table 15.6 & 15.7*).

(Table-15.6): All India-crop-wise additional breeder seed production targ. & achiev.

(Qty.: qtl)

Crops	2016-17		2017-18		2018-19	Total		
	Target	Ach.	Target	Ach.	Target	Target	Ach.	% Ach.
Tur	425	978	157	548	168	750	1526	203
Urd	317	720	183	304	212	712	1024	144
Mung	490	790	195	374	244	929	1164	125
Gram	2140	2917	235	3241	277	2652	6158	232
Lentil	165	250	140	205	132	437	455	104
Pea	180	350	68	190	73	321	540	168
Total Pulses	3717	6005	978	4862	1106	5801	10867	187

Source: ICAR-IIPR, Kanpur




Fig.15.7: Production of Breeder seed under ABSP

• Infrastructure created under EBSP: Breeder seed infrastructure strengthening comprises of several need based items at different centres such as works (threshing floors), Seed Processing Plants (SPP), Farm implements (tractors, sprinkler systems, power sprayers, Rotavator, BBF Planter, Ridge Planter, Storage godown, irrigation channel, fencing, bore-well, power thresher, seed-cum ferti-drill, pick-up van, irrigation pipes, hydraulic trolley, gravity separator, weighing machines, combine harvester).

(Table-15.7): Infrastructure: Strengthening of seed production farms

Allocation (Rs. Crore)

States	Institutes		Infrastructural Activities					
		Alloc.	CW	SPU	FI	Total		
Madhya	ICAR-IIPR, RS-	2.75	02	01	02	05		
Pradesh	Phanda, Bhopal							
	JNKVV, Jabalpur	1.90	06	-	02	08		
	RVSKVV, Gwalior	0.92	02	-	17	19		
Bihar	Agril. University,	1.95	03	01	02	06		
	Sabour (Bihar)							
Rajasthan	ARS, Kota Agri.	1.80	05	02	08	15		
	University, Kota							
	RARI, Durgapur,	1.47	02	02	04	08		
	Shri Karan Narendra							
	Agri. University,							
	Jobner							
Maharashtra	Agril. Research	1.45	05	01	01	07		
	Station, Badnapur,							
	VNM Krishi							
	vidyapeeth Parbhani							
	MPKVV, Rahuri	1.58	04	-	05	09		
Uttar Pradesh	/ 1	1.65	04	01	-	05		
Andhra	Agril. Research	1.75	03	02	-	04		
Pradesh	Station, LAM,							
	ANGRAU							
Karnataka	UAS, Dharwad	1.77	-	-	03	03		
Odisha	Agril. Research 1.40		03	01	01	05		
	Station, Berhampur,							
	OUAT,							
	KrishiBhubneshwar							
Total- 08	12	20.39	39	11	45	94		

SPU-Seed Processing Unit; FI-Farm Implements; CW-Civil Work; Alloc.-Allocation

15.6.2 Creation of Seed-Hubs

i) Creation of Seed-Hub

- To enhance the quality and quantity of pulses seed in the country, a project on creation of seed-hubs (2016-17 to 2018-19) has been initiated under (NFSM) with the mandated objectives and targeted seed production of latest varieties (150 locations in 24 states, at ICAR Institutes 07, ICAR-AICRPs centers -46, KVKs-97) across 24 states (Andhra Pradesh, Assam, Bihar, Chhatisgarh, Gujarat, Haryana, Himachal Pradesh, Jharkhand, J&K, Karnataka, Kerela, Manipur, Maharashtra, Madhya Pradesh, Nagaland, Odisha, Punjab, Rajasthan, Tamil Nadu, Telangana, Tripura, Uttar Pradesh, Uttarakhand and West Bengal) in the country. Each seed hub has to produce 1000 qtls of seed per year.
- With Budgetary allocation of Rs. 225.31 crores, each Seed-Hub has a financial assistance of Rs. 1.50 crore (infrastructure- Rs. 50 lakh for Storage of seeds/processing + Rs. 100 lakh revolving fund towards production, procurement, processing, of seeds during 2016-17 and 2017-18). Almost 90% of infrastructure has been completed and production has already started from 2016-17.

• Crop-wise/Centre-wise targets for quality seed production and state-wise permanent structure/Seed Processing Units (SPU) created are given below (Table-15.8)

(Table-15.8): All India-Crop-wise seed production target and achiev. under seed-hub

(Qty.: in qtl)

Crops	2010	2016-17		7-18	2018-19		Total	
	Target	Ach.	Target	Ach.	Target	Target	Ach.	% Ach.
Pigeonpea	15750	6865	25540	17927	31100	72390	24792	34
Urd	8675	4553	15850	15520	18600	43125	20073	47
Moong	16225	5857	28900	21050	34100	79225	26907	34
Chickpea	20510	11921	28700	39058	34150	83360	50979	61
Lentil	7325	4642	11550	9821	13700	32575	14463	44
Fieldpea	9550	1754	12530	4826	14500	36580	6580	18
Lathyrus	150	55	250	532	350	750	587	78
Rajmash	350		450	851	550	1350	851	63
Cowpea	900	222	2150	1316	2050	5100	1538	30
Mothbean	150	46	550	475	700	1400	521	37
Horsegram	400	64	600	425	650	1650	489	30
Total Pulses	79985	35979	127070	111801	150450	357505	147780	41

Source: ICAR-IIPR, Kanpur




Fig.-15.8: Production of quality seed under Seed-hub

15.6.3 Seed Village Programme

To address critical input, the seed village programme has been operationalised to improve the quality and stock of farm saved seeds of pulses enhancing crop production/productivity. To upgrade the quality and varieties of farm-saved seeds which is about 80-85% of the total seed used, of cluster of 50 farmers @ 1 acre is provided with 60% financial assistance towards foundation/certified seed of pulses for production of certified/quality seeds. The farmers are also imparted training on seed production technology.

During 2017-18 the seed production programme was conducted in >9 lakh ha area by distribution of 7 lakh qtls of seed. In all about 28 lakh farmers were benefitted in >1 lakh villages of the country. The state-wise detail is as under (Table-15.9).

(Table- 15.9): Seed distribution under Seed Village Programme (2017-18)

S. No.	State Agency name	Area (in ha)	F/C seed distribution (qtls.)	Qty. of Seeds Produced (qtl.)	No. of Seed Villages Organized	Total (M+F)
1	Tamilnadu		102771	228240	3258	450919
2	KOF,Karnatka	551.2	618	0	269	1378
3	ASC-Khrif	44400	16299	774600	1414	212100
4	ASC-Rabi	51450	7980	71100	1715	256290
5	ASC-Summer	19185	6171.6	369718	897	90526
6	ASSCA	6440	2576	80500	161	7793
7	Uttar Pradesh (K)	27350	8997	668189	15971	68376
8	Uttar Pradesh (R)	46564.8	46127	904794	25095	116412
9	IISS, Mau Uttar Pradesh	2699	2645	47678	317	6887
10	Chhattisgarh	173093	49805	1069901	4045	158841
11	Telangana	37650	26562	791495	3765	94125
12	Madhya Pradesh	196838	137612	1308655	7945	415506
13	Bihar (K-17)	18383	5430.1	425355	638	41294
14	Bihar(R-2017-18)	16325	15459	397720	719	52502
15	Uttarakhand K-17	4183	1274	19380	2850	15625
16	Uttarakhand R-17	9057	8467	132898	4256	23957
17	MSSC (R)	6772	88968	1514631	16697	286277
18	MSSC(K)		1768	23576	1433	5894
19	Andhra Pradesh	19370	12201	539040	1937	48425
20	RSSC-K	33167	8065	283548	1415	141456
21	RSSC-R	46581	42291	1319117	1165	116452
22	RSSC-zaid/ summer	2915	583	16300	72	7286
23	Himachal Pradesh	32664	32664	580721	1868	137647
24	Jammu & Kashmir K-17	28208	12491	727253	804	70913
25	Jammu & Kashmir R-18	45960	41112	0	938	124645
26	NSC		0	22022	189	3344
27	IARI, Karnal	52.8	6.6	2376	8	115
28	Odisha	9604	1921	104750	137	9604
29	Meghalya	32815	5913	237980	275	39625
30	Nagaland K-17			3246	27	1350
	Total	912278	686777	12664783	100280	3005564

Source: Seed Division, DAC&FW, N. Delhi

(Table-15.10): State-wise Physical and Financial Progress under Seed Village Programme

(Amount Rs. in lakhs; Qty. in qtl.)

S. No	Name of		2015-16				201	6-17		2017-18			
	State	Amount Released	Seed Village	Seed produced	farmers Covered/ Benefited	Amount Released	Seed Villag e	Seed produ ced	farmers Covered/ Benefited	Amount Released	Seed Village	Seed produc ed	farmers Covered/ Benefited
1	AP	650	6475	1135919	161850	37	1610	59031 5	40250	352	1937	539040	48425
2	Assam	1060	2771	1944810	415655	1000				1313	4186	1275918	566709
3	Arunachal Pradesh		36	10900	1800					52			
4	Bihar	600	885	648505	86321	635	376	238916	36210	324	1357	823075	93796
5	CG	245	1243	625093	89986	281	1150	684799	57538	518	4045	1069901	158841
6	Gujarat		58	9025	1223					24			
7	HP	248	1754	638868	172682	375	1498	505268	130480	386	1868	580721	137647
8	Haryana	1	18	1830	191	1	11	3186	154	1	8	2376	115
9	J&K	391	1178	415467	178531	691	1379	898826	208703	899	1742	727253	195558
10	Karnataka	20	380	91730	6372	25	163	51950	2338	83	269		1378
11	Kerala	4	9	18000	404	27							
12	MP	587	2441	1287398	239858	1395	1202	627440	75482	2056	7945	1308655	415506
13	MH		1320	110322	30260	600	729	177460	64318	2431	18130	1538207	292171
14	Meghalaya	98	305	69426	20755	11				142	275	237980	39625
15	Mizoram					25							
16	Manipur	4				55							
17	Nagaland	118					62	3969	2850	47	27	3246	1350
18	Orissa		213	298177	21295		105				137	104750	9604
19	Punjab	318	630	543570	63076				10300				
20	Rajasthan	122	312	296570	19266	101	1621	94536 1	99898	1511	2653	1625985	265194
21	Sikkim												
22	TN	970	1918	159079	447501	702	2484		707979	1330	3258	228240	450919
23	Tripura												
24	UP	900	2632	426050	62390	124	5149	390280	91511	1436	41383	1620661	191675
25	Uttrakhand	303	2888	75707	49475	88	4468	158819	32435	229	7106	152278	39582
26	Pondi.	0	1500	25.52.			4	12222	29	00.6	25.5	5 04.40 -	04425
27	Telangana	300	1783	375724	44575	550	2187	423385	54675	936	3765	791495	94125
28	NSC					980					189	22022	3344
29	West Bengal												
	Total	6939	29249	9182170	2113466	7701	24198	699974	1615150	14070	100280	12651803	3005564

Source: Seed Division, Ministry of Agri. & FW, (DAC&FW), N. Delhi




Fig.-15.9: Seed Village Programme - Phy. & Fin. Progress (Table-15.11): Crop-wise Seeds distributed and produced under SVP during 2017-18

(Quantity- 000 Qtls)

Crop	Production	Distribution	% Distributed
Lentil	19.94	1.37	6.89
Gram	925.40	83.33	9.01
Peas	6.78	1.07	15.74
Arhar	76.61	1.96	2.56
Moong	245.91	9.09	3.70
Urd	65.14	17.40	26.72
Moth	58.92	1.47	2.50
Rajmash	3.26	0.37	11.31
Lathyrus	0.84	0.06	7.61
Lobia	7.51	0.46	6.10
Total	1410.32	116.59	8.27




Fig.-15.10: Crop-wise Seed Distributed - 2017-18



Chapter-16

Production and Sustainability Constraints Identified

Based on the review of the planned agricultural development programmes under NFSM-pulses/RKVY/BGREI etc., across the states, Mid-term evaluation, Impact evaluation of NFSM and studies /NLMT reports of Directorate of Pulses Development on reasons for low production, coverage and productivity in pulses, the policy initiatives/interventions were taken under four major constraints/categories viz., *i) Production ii) Inputs iii) Marketing and iv) Technology dissemination.*

16.1 Constraints related to production

Production potential exhibited under different crops in analysis of yield gaps among major states, districts and within district; under the FLDs were considered as production constraints. Here complete package technology i.e. integration of all components viz. timely sowing, high yielding varieties, fertilizer management based on soil testing (including foliar nutrition), rhizobium inoculation, weed management, IPM etc., programme were felt necessary to be pursued vigorously for adaptation (*Table-16.1 and 16.2*).

(Table-16.1): Technological yield gap exhibiting the production related constraints-FLDs.

(Yield: kg/ha)

Crop	Yi	ield (kg/ha	a)	Gap ove	r FP	Gap ove	r SAY
_	IP	FP	SAY	Actual	%	Actual	%
Pigeonpea	1394	1078	863	316	29	530	61
Chickpea	1502	1244	907	257	21	594	66
Rice fallow Chickpea	1275	960	976	315	33	299	31
Mungbean(Kh)	781	608	435	173	28	345	79
Mungbean (R)	1398	1228	704	170	14	694	99
Mungbean (RF)	960	723	532	237	33	428	80
Mungbean Summer/Spring	931	559	674	372	66	257	38
Urdbean (Kha.)	813	622	368	191	31	445	121
Urdbean (R)	1203	986	774	217	22	429	55
Urdbean (RF)	1185	1002	774	183	18	411	53
Lentil	1289	966	777	323	33	512	66
Field pea	1225	933	904	292	31	321	36
Average (All Pulses)	1163	909	724	254	30	439	65

Source-Annual Report -2017-18, GoI, DPD, Bhopal (Ave. 2013-14 to 2015-16)

State Average Yield (SAY-E&S (Ave. 2011-12 to 2015-16) IP: Improved Practice, FP: Farmers Practices.

(Table-16.2): Yield gap exhibiting the production related constraints among the states

(Yield-kg/ha)

Crop/Season	National Yield	Highest/ Lowest Yield	States > National Avg.	States < National Avg.
Total Pulses	835	HP (1338) /J&K (397)	AP, Bihar, Gujarat, HP, Jharkhand, kerela, MP, Punjab, Telengana, UP, Uttarakhand, WB	Assam, CG, Haryana, J&K, Karnataka, Maharashtra, Odisha, Rajasthan, Tamil nadu,
Total Kharif	654	Kerela (1499) / J&K (373	Assam, Bihar, Gujarat, Haryana, HP, Jharkhand, kerela, MP, Punjab, Tamil nadu, Telengana, UP, Uttarakhand, WB	AP, CG, J&K, Karnataka, Maharashtra, Odisha, Rajasthan
Total Rabi	994	HP (2198)/Odisha (516)	AP, Gujarat, Jharkhand, HP, Rajasthan, MP, Telangana, UP, WB	Assam, Bihar, CG, Haryana,J&K, Karnataka, Kerela, Maharashtra, Odisha, Punjab, Tamil nadu, Uttarakhand
Tur	937	MP (1297)/ HP (360)	Bihar, Gujarat, Haryana, Jharkhand, Kerela, MP, Rajasthan, TN, UP, Uttarakhand, WB	AP, Assam, CG, HP, Karnataka, Maharashtra, Odisha, Punjab
Mungbean (K)	429	WB(900)/ Karnataka (237)	AP, Assam, Bihar, MP, Gujarat, Karnataka, UP, Punjab, TN, Telangana, WB	CG, Odisha
Urdbean (K)	630	Jharkhand (918)/ CG (320)	AP, Bihar, Gujarat, Jharkhand, MP, TN, Telangana, Uttarakhand, WB	CG, Haryana, HP, Karnataka, Maharashtra, Odisha, Punjab, Rajasthan, UP
Gram	1055	Telangana (1456) / Karnataka (600)	AP, Gujarat, HP, Jharkhand, MP, Punjab, Telangana, Rajasthan, UP,WB	Assam, CG, Haryana, Karnataka, Maharashtra, odisha, Tamilnadu, Uttarakhand
Urd (Rabi)	764	Madhya Pradesh (1400)/Chhatti sgarh (264)	AP, MP, Telangana, WB	Assam, CG, Gujarat, Karnataka, odisha, TN, UP
Mungbean (Rabi)	618	MP (1203)/ Chhattisgarh (234)	AP, Assam, Bihar, MP, Punjab, Telangana, UP, WB	Bihar, CG, Gujarat, Karnataka,Odisha, TN
Lentil	1008	Rajasthan (1408)/ CG (325)	MP, Rajasthan	Assam, Bihar, CG, HP, Jharkhand, Odisha, Punjab, Uttarakhand, WB

Source-Annual Report -2017-18, GoI, DPD, Bhopal

Based on the production related constraints, state-wise interventions were identified and States Annual Action Plan (AAPs) were approved accordingly (*Table-16.3*).

(Table-16.3): Identified production related constraints and their interventions

States	Production /	Crop	Loss	Measures adopted
	constraints	Стор	percentage	/ interventions
			l r · · · · · · · · · · · · · · · · · ·	made
UP, MP, PB,	Mid-season cold waves	Gram,	10-40%	Tolerant
Haryana	and terminal heat during	Lentil		Varieties, MIS
	Rabi	Pigeonpea		Intercropping
MP, MS, Guj,	Inundation of water in	Pigeonpea,	10-50%	Planting under BBF
AP, TN	black cotton soils during	Urd,Mung		Planting under
	heavy rains sub-optimal			Furrow Irrigated
	nutrient uptake			Raised Bed (FIRB)
				Inter-cropping
				RCT provisions
All states	Micronutrient	All Pulse	-	INM provided @
	deficiency (Zn, Fe, B,	crops		Rs.500/ha
	and Mo) - unbalanced			
	use/seldom soil test;			
100 100 00 1	Quality issues			0.7.7.0
MP, MS, Guj,	Sulphur deficiency;	All Pulse	-	@ Rs.750 per/ha
AP ,Karnataka,	inadequate availability	Crops		provision
UP	of Gypsum or pyrites	D'	10.500/	TD) (
UP, MP,	Podfly and maruca	Pigeonpea	10-50%	IPM provision
Bihar,				made
JH., Punjab, Haryana				
MP, UP,	Fusarium wilt	Chickpea	20-25%	IPM seed treatment
Bihar,	Fusanum wiit	Tur &	10-15%	irwi seed tieatilielit
Jharkhand		Lentil	10-13 /0	
Jilai Kilaila	YMV & Powdery	Urdbean&	10-50%	Rest. Varieties
All States	mildew	Mungbean	10 30 70	method of planting
including MP	milite W	Widingocan		- IPM
UP, Bihar,	Stray cattle/ Blue bull	All Pulse		Solar fencing under
MP,	meanace	Crops		RKVY Local
Jharkhand, RJ,		•		initiative
CG, Haryana				
	Region specific	All Pulse		Minikit provided,
	technologies-Pigeonpea	Crops		awareness and
All states	on bunds transplanting/			popularize these
An states	intercropping etc.			technologies
				through NFSM-
				Pulses

Source-Annual Report -2017-18, GoI, DPD, Bhopal

• Cluster Frontline Demonstration: Pulses

Government has initiated National Level Cluster Frontline Demonstrations on pulses, through Krishi Vigyan Kendra under 11 Agriculture Technology Application Research Institute (ATARIs) through 549 KVKs to demonstrate the production potential of new varieties and the related technologies; increasing production through area expansion and productivity enhancement in a sustainable manner in the identified districts of the country; restoring soil fertility and productivity at the individual farm level; and enhancing farm level economy (*i.e.* farm profits) to restore confidence amongst the farmers.

Cluster FLD on Pulses (minimum 10 ha each) by ATARI (Rs. 26.11 Crore for 2017-18) is operational in 31366 ha area @ Rs. 7500/ha (Rs. 750 for monitoring + literature + field day) across the country.

The transfer of technology through CFLDs have increased yield levels shown upto 42% and 54% over local check and normal yield. Crop-wise details of CFLDs conducted, yield gaps and varieties demonstrated during 2016-17 is given below (Table-16.4 to 16.5).

(Table-16.4): All India CFLDs targets and achievement (2015-16 to 2018-19)

(Area: ha)

Year	Target	% Area Increase from 2015-16	Achievement
2015-16	15382		13528
2016-17	31000 (By 534 KVKs)	102	29008
2017-18	31366 (By 549 KVKs)	103	30366
2018-19*	34750 (By 578 KVKs)	126	-

Source: ICAR-ATARI



Fig.-16.1: All India CFLDs Targets and Achievement (2015-16 to 2018-19)

(Table-16.5): All India: CFLDs yield gaps exhibited: Total Pulses (2016-17)

(Yield: qtls/ha)

Crop	Area (ha)		Yield		Actual Yi	eld Gap	% Yield	l Gap
	_	Demo.	Local	NY	Local	NY	Local	NY
			Check		Check		Check	
Pigeonpea	4824.80	14.50	9.92	7.25	4.58	7.25	46	100
Black gram	4469	8.67	6.39	6.32	2.28	2.35	36	37
Green gram	7491.20	8.30	5.90	5.00	2.40	3.3	41	66
Horse gram	228.00	8.15	6.05	4.83	2.10	3.32	35	69
Chick Pea	6054.46	15.07	11.02	9.32	4.05	5.75	37	62
Field Pea	1819.09	15.89	11.11	9.40	4.78	6.49	43	69
Lentil	3689.70	10.66	7.58	7.38	3.08	3.28	41	44
Rajmash	203.65	13.67	8.07	8.07	5.60	5.60	69	69
Moth bean	112.00	6.23	4.82	4.82	1.41	1.41	29	29
Lathyrus	116.00	7.16	5.53	7.76	1.63	-0.6	29	-8
Total	29007.90	10.83	7.64	7.02	3.19	3.82	42	54

NY: Normal Yield: (Ave. 2011-12 to 2015-16) Yield of Local Check considered as NY.

(Table-16.6): All India: CFLDs yield gaps: Kharif Pulses (2016-17)

(Yield:qtls/ha)

							(Tieia:qii	is/na)
Crop	Area (ha)		Yield		Actual Yie	ld Gap	% Yield Gap	over
	_	Demo.	Local	NY	Local	NY	Local Check	NY
			check		Check			
Pigeon pea	4741.8	12.22	8.51	7.25	3.71	4.97	44	69
Black gram	2734.6	8.39	5.85	5.19	2.54	3.2	43	62
Green gram	2103.8	8.66	6.34	4.14	2.32	4.52	37	109
Horse gram	228	8.15	6.05	4.57	2.1	3.58	35	78
Rajmash	19.2	13.31	6.2	6.2	7.11	7.11	115	115
Moth bean	112	6.23	4.82	4.82	1.41	1.41	29	29
Total	9939.40	9.49	6.30	5.36	3.20	4.13	51	77

NY: Normal Yield: (Ave. 2011-12 to 2015-16) Yield of Local Check considered as NY of Rajmash & Mothbean

(Table-16.7): All India: CFLDs yield gaps: Rabi Pulses (2016-17)

(Yield:qtls/ha)

Crop	Area (ha)		Yield (q/ha)	Actual Yield Gape over % Yield Gap over				
	-	Demo.	Local Check	NY	Local Check	NY	Local Check	NY
Chick Pea	6054.46	15.07	11.02	9.32	4.05	5.75	37	62
Lentil	3689.7	10.66	7.58	7.38	3.08	3.28	41	44
Field Pea	1819.09	15.89	11.11	9.40	4.78	6.49	43	69
Green gram	1449.4	7.99	5.24	5.77	2.75	2.22	52	38
Black gram	1020.2	9.33	6.88	7.45	2.45	1.88	36	25
Pigeon pea	83	16.82	11.33	11.33	5.49	5.49	48	48
Rajmash	180	19.60	11.88	11.88	7.72	7.72	65	65
Lathyrus	116	7.16	5.53	7.76	1.63	-0.6	29	-8
Total	14411.85	12.82	8.82	8.79	3.99	4.03	45	46

NY: Normal Yield: (Ave. 2011-12 to 2015-16) Yield of Local Check considered as NY of Rajmash&Mothbean.

(Table-16.8): All India: Varieties Demonstrated under CFLDs

Crop	Variety
Pigeonpea	TJT-501, AL-201, Malviya Arhar-13, Asha, UPAS-120, NA-2, BSM3-736, GRG-811
Blackgram	Mash 114, UG 218, Shekhar, PU-31, AKU-15, IPU-94-1, Shekar, PU-19
Greengram	MH421, Pant Moong 5, IPM 2-3, Tripura Moong-1, MGG-347, Uttara, GM-4, KKM-3,
	SML-668, SML-832, HUM-16, Samrat
Horsegram	BirsaKulthi, Pavur-2, Indrakulthi
Rajmash	WazejRajmash, Tripura Rajmash Sel-1
Mothbean	RMO-257, CZM-2
Lentil	HUL 57, DPL 62, L 4594.
Field pea	Prakash, Rachna, Anupam, Shalimar pea
Chickpea	JG 16, HC1, GNG 1581, HC 1, HC 5, JAKI 9218, Vijay, Vishal, Digvijay, NBEG-3,
	BDNG-797,GG-2, GJG-3, GG-5, JGG-1, GG-5
Blackgram	LBG 752, LBG-787, GBG-1, PU-31
Greengram	LGG460, WGG-42, TM96-2, MGG-295,GG- 2, Co-4,CO-4, IPM-99-125

(Table-16.9): Cluster FLD on Pulses 2017-18

(KVK-Nos.; FLDs-Area in ha; Budget-Rs. in Lakhs)

Implementing Agency/Zone	NFSM-Pulses		
[KVK	FLDs	Budget
ATARI-I (Ludhina)	52	1522	125.46
ATARI-II (Jodhpur)	49	3520	296.31
ATARI-III (Kanpur)	68	4220	354.80
ATARI-IV (Patna)	62	4350	360.36
ATARI-V (Kolkata)	49	3080	255.81
ATARI-VI (Guwahati)	32	1510	127.26
ATARI-VII (Barapani)	16	880	78.81
ATARI-VIII (Pune)	67	3810	315.36
ATARI-IX (Jabalpur)	63	4440	365.91
ATARI-X (Hyderabad)	58	2750	222.66
ATARI-XI (Banglore)	33	1284	108.51
Total (2017-18)	549	31366	2611.25
Total (2016-17)			

16.2 Constraints related to inputs

- Quality and timely availability of critical inputs seeds, varieties, bio-fertilizers, micronutrient and critical irrigation were identified across the states and felt necessary to be addressed as one of the major strategies under this category.
- Non-availability of location specific/recommended high yielding varieties quality certified seeds at all levels as the production and distribution is usually for the very old and known varieties which are generally poor performers.
- Poor availability of quality/certified seed/poor varietal development/limited varietal choise during last 10 years & poor varietal diversification of pulses in India. Crop-wise gap of availability of quality certified seeds, varietal development & varietal choice last one decade & Poor varietal diversification are given table 16.10 to 16.13.

- Non-availability of quality inputs at village level (sometimes even at block levels); inflow of spurious and sub-standard seeds, rhizobium culture/PSB, micro-nutrients, biointensive/bio-pesticides.
- Non-popularization/lack of demonstration and availability of implements like light seed drills, zero-till machine/rotavator/and ridge-maker (custom-hiring or community runbasis) in big areas of Bundelkhand region of U.P., and M.P.
- Pulses respond favorably to 1-2 critical irrigations for good yields, however, lack of power supply/low-voltage, non-opening of canal and less priority to the crop-group in addressing the water carrying/micro-irrigation related problems.
- Lack of domestic milling support and Post Harvest Technology (PHT)/value addition support.

(Table-16.10): Requirement and availability of certified seeds during 2016-17

Quantity: Thousand Tonnes

Crop	Requirement	Availability	Deficit/Surplus
Gram	181.43	148.55	-32.87
Moong	5.94	7.70	1.75
Urd	7.82	8.09	0.27
Arhar	0.13	0.62	0.49
Lentil	13.05	10.56	-2.49
Peas	21.17	18.28	-2.88
Cowpea	0.44	0.70	0.26
Horsegram	1.56	1.56	0.00
Indian Bean	0.13	0.13	0.00
Khesari	0.62	0.64	0.02
Rajma	0.62	0.56	-0.06
Total Pulses	232.91	197.39	-35.51

(Table-16.11): Poor Varietal Development (XIth and XIIth Plan)

(a) Varietal Release Profile-Notification during Last 10 Years (2006 to 2015)

Crop	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	Total
Arhar	2	6	4	3	-	1	2	3	2	1	24
Urd	2	-	4	4	4	3	3	1	3	1	25
Moong	3	2	2	4	4	3	3	1	2	-	24
Gram	7	11	5	3	7	1	6	5	2	2	49
Lentil	3	2	1	3	4	2	1	1	1	-	18
Peas	3	3	5	1	2	3	1	-	2	-	20

(Table-16.12): Limited statesøvarietal choice during last 10 years (2006-2015)

State/Crop	Arhar	Urd	Moong	Chickpea	Lentil	Peas
Maharashtra	7	3	2	10	1	2
Rajasthan	-	5	4	16	2	4
A.P.	6	8	3	1	-	-
Karnataka	5	5	4	3	-	-
Gujarat	4	1	-	7	-	-
Odisha	2	6	2	1	-	-
M.P.	4	<mark>2</mark>	-	18	<mark>1</mark>	2
U.P.	<mark>2</mark>	7	4	7	6	10
Bihar	<mark>2</mark>	2	1	1	<mark>1</mark>	3
Tamil Nadu	2	8	3	1	-	-

(Table-16.13): Poor Varietal Diversification (VRR)

State	Crop	Prevalent Varieties	Recommended Varieties	
N (- 11	D:	TIT 501 ICDI 07110 No.	(ICAR/SAUs)	
Madhya	Pigeonpea	TJT 501, ICPL 87119, Non-	TJT-501, ICPL 87119, ICPL	
Pradesh	* * 11	descript	88039, JA 4	
	Urdbean	T-9, HFP8909, IPU-94-1,	KU-96-3, PU 30, MASH 338	
		Non-descript		
	Moongbean	HUM-1, HUM-12, Non-	HUM 1 JM 721, TARM 1, HUM	
		descript	6	
	Chickpea	JG 11, JG 16, JG 130, JAKI	JG-130, JG-322, JG 63	
		9218		
	Lentil	JL 1, Mallika, DPL 62,	JL1, K-75, IPL 406	
		IPL 81		
	Peas	Arkel, Azad-1	KPMR-400, IM 9101 (Subhra),	
			Rachna	
Maharashtra	Pigeonpea	ICPL-87119, ICPL-8863,	ICPL-87119, ICPL-8863, BDN-	
		BSMR-736, Vipula	708, BDN-711	
	Urdbean	TAU-1	BDU-1, TPU-4, TAU-1	
	Moongbean	Kopargaon-1, Utkarsha	BPMR-145, BM-4, 2002-01,	
			Vaibav	
	Chickpea	Chaffa, Agnirekha	BDN-9-3, PKV-2, 5, 4-1, JAKI-	
	1	, ,	9218	
Maharashtra	Other Kharif	-	Seena, Maan (Kulthi)	
	Pulses		, , ,	
	Other Rabi	Ratna local(Khesari),	Ratna (Khesari), Pusa Komal	
	Pulses	Parvati (Cowpea)	(Cowpea)	
Rajasthan	Pigeonpea	ICPL-151, ICPL-87,	ICPL-151, ICPL-87, Gwalior-3,	
,		Gwalior-3	UPAS 120	
	Urdbean	T-9, Pant U 19	T-9, RBU 38, Pant U 19	
	Moongbean	K-851, RMG-62, RMG-268	K-851, RMG-62, RMG-268	
	Chickpea	Dahod Yellow, RSG 888	RSG 902, GNG 1581, Pratap Raj	
			Chana, RSG 991,	

State	Crop	Prevalent Varieties	Recommended Varieties (ICAR/SAUs)
Uttar Pradesh	Pigeonpea	Rajeev Lochan, PAU-881, VL Arhar-1, Pusa-992, Malviya Chamatkar (MAL-	NDA-2, Pusa-992, MAL-13, PAU-881, NDA-88-2, KA-32-1, K91-25
	Urdbean	13) Pant Urd-31 & 40, LAM-709, Azad Urd-3	NUL-7, Vallabh Urd-1, Azad urd- 1, Uttra, Shekhar-2, 3
	Moongbean	IPM-02-3, Pant Mung-6, TM-96-2, Meha	KM-2195, MH-421, HUM-16, Pant Mung-4, Pusa-9531, Pusa Vishal
Uttar Pradesh	Chickpea	RVG-101, 210, PKV Kabuli-4,RSG-991, Pusa- 1103, HK-94-124	GNG-1969, GNG-1958, WGG-3, HK-2, RSG-963, WCG-10, JGK- 1, RSG-88
	Lentil	Pant Lentil-7, PL-02, HUL-57, VL-507	KLB-303, KLB-320, PL-8, HUL-57, IPL-406, Pant Lentil-4, DPL-15
	Peas	Sapna, VL Matar-47, VP- 101	VP101, Pant P 13, IPF-5-19, SKNP 04-09
Andhra Pradesh	Pigeonpea	LRG-41, PRG-158 & ICPH-2740	LRG-41, PRG-158
	Urdbean	PU-31, LAM, TM-76-2	PU-31, LBG-752 & TM-76-2
	Moongbean	LGG-460, TM-96-2	LGG-460, TM-96-2
	Chickpea	JG-11, JAKI-9218 & PBH-4	
Karnataka	Pigeonpea	BRG-1, BRG-2	ICP 8863 (Maruthi), ICPL 87119 (Asha), ICPL 87 (Pragathi)
	Urdbean	TAU-1, T-9	Kargane-3, T-9, LBG-625
	Moongbean		PS-16, Pusa baisaki
	Chickpea	Annigeri-1, JG-11	Annigeri-1, JG-11, KAK-2, Vishal
	Kulthi	Hebbal Local	KBH-1, PHG-9
Gujarat	Pigeonpea	Gujarat Tur-100, Gujarat Vegetable Tur-1	BDN-2
	Urdbean	TPU-4, Gujarat Urd-1	T-9
	Moongbean	Gujarat Mung-3, CO-4	GM-4, K-851
	Chickpea	GG-1, Chaffa, Dahod Yellow, ICCC-4, Gujarat Junagarh Gram-3	Gujarat Gram-4
Telangana	Pigeonpea	Asha, ICPL 87119, ICPL 85063, PRG 158, WRG 65, MRG 1004, LRG 41 & LRG 158	Asha, ICPL 85063, LRG 41 & PRG 158
	Urdbean	PU 31, LBG 752, LBG 787	PU 31, LBG 752, LBG 787
	Moongbean	LGG 460, MGG 295	LGG 460, MGG 295, MGG 347 & 348, MGG 42
	Chickpea	JG 11, JAKI 9218	JG 11, JAKI 9218

16.3 Constraints related to marketing

- Distress sale, poor minimum support prices compared to cost of cultivation, exim policy issues, non-accessibility to market, post harvest losses etc., were identified as major marketing related constraints, especially in major pulse producing state. The policies to provide remunerative prices to the farmers including the procurement facilities were felt vital by the government.
- Wide price-gap between the whole and processed/milled product in the chain of farmer/producer-buyer-consumers, vulnerability to stored grains due to lack of scientific storage facilities at domestic level, lack of support to small scale processing, packaging, value addition and non-linking of pulses to procurement policy commensurate to staple food grains like wheat and paddy, are the other major market related constraints.

16.4 Constraints related to extension and their interventions

- Lack on guidance for proper certified seed production/variety identification, insectpest/diseases identification and management phases, importance and procedure of seed
 treatment/rhizobium inoculation, lack of information/knowledge on current advances in
 production, management technology, and also poor or no knowledge about organizing
 seed production and its protection for succeeding crop.
- Poor knowledge base on nutrient use efficiency (NUE), IPM, method of preparation of spray solutions and multiplicity of extension system on IPM, esp., pesticide dealers etc., identified as technology transfer or extension related constraints.
- Depleting public sector extension support, non-positioning of skilled/sound extension functionaries at the grass-root level (Block/villages) the technology dissemination/extension activities have adversely affected.
- The extension workers also lack advances in technological sector and there is a gap of HRD activities. Quality cluster demonstrations have been an observation across the board.
- Interface between State Department of Agriculture (SDA) and State Agricultural Universities (SAUs), ICAR (ATARI) and Department of Agriculture & Cooperation DAC and other allied state level/district level field functionaries also seems to be bleak and visible with the absolute communication gap in conduction/organization of FLDs, and cluster demonstrations, FFS, IPM, etc.

16.5 Suggestions

16.5.1 Input Related Interventions

Input related constraints are the major bottlenecks in increasing area and production of pulses in the country, following may, therefore, be suggested:

A tie-up arrangement amongst state + ICAR (breeder seed producers), Seeds and NFSM Divisions, Government of India, Department of Agriculture, Cooperation & Farmers Welfare need to be more strengthened for advance indenting of breeder seeds. For production of foundation and certified seed, besides making cent-per cent utilization of centrally sponsored schemes on pulses (NFSM). States need to enter in to MoU with the private seed producers, NGOs and FPOs/SHGs/FOs/FIGs etc.

- On going *seed hub programme* project under NFSM, operational since 2016-17, need serious implementation by KVKs and other associated agencies for their sustainability.
- To ensure the timeliness, availability of quality inputs at cost effective and approachable common panchait/village place, each potential district, its blocks should identify village-clusters, formulate Pulses Self-Help Groups (PSHG). Under the chairmanship of Rural Agriculture Extension Officer (RAEO) or ADO. A committee, comprising of representatives from PSHG, Cooperative society, local rural bank, pesticide dealer, block Electricity Board and panchayt representative may be constituted. The committee should prepare season-wise Strategic Pulses Production Plan (SPPP), delineating input requirement, much in advance. The SPPP should be fine-tuned by the ADO-further refined by the Deputy Director Agriculture for final appraisal/review/approval by Chief Executive Officer/District Magistrate, Chairman of DFSMEC/ATMA.
- Supply of electricity for critical irrigation at the critical period of crop growth, credit support and all such vital input aspects may be properly addressed in an institutionalized manner by the DFSMEC.

16.5.2 Production Related Interventions

• Based on the analysis of production and productivity on all India basis (crop-wise analysis), ten potential districts each for pigeonpea, chickpea, blackgram, greengram and lentil, categorised as the major contributors (5-40 per cent of total all India production in the specific pulse crop), may be adopted by the respective SDAs/SAUs. These districts may be saturated with the entire pulse related development and research programme on cent per cent implementation basis. At least 20 number of each FLDs, FFS,IPM, infrastructural development and minikits demonstration need to be taken in each block/panchayat on cluster demonstration basis: Crop-wise ten potential districts are indicated below:

(Table-16.14): Crop-Wise Potential Districts With 20-30% Prod. Shareó All India

Crop	Districts
Gram	Kurnool, Vidisha, Sagar, Raisen, Ashok nagar, Dewas, Rajgarh, Dhar,
	Chhatarpur, Panna
Arhar	Prakasam, Kurnool, Betul, Fatehpur, Hamirpur, Seoni, Sonbhadra,
	Mirzapur, Jabalpur, Morena
Moong	Jagatsingpur, East Godavari, Nayagarh, Kedrapara, Puri, Bolangir,
	Vizianagarm, Thiruvarur, Mahoba, Jhansi
Urd	Krishna, Lalitpur, Guntur, Jhansi, Mahoba, Srikakulam, Unnao,
	Damoh, Sagar, Jabalpur
Lentil	Bahraich, Sagar, Vidisha, Panna, Hamirpur, Balrampur, Jhansi,
	Damoh, Chitrakut, Shivasti
Field Pea	Jalaun, Lalitpur, Jhansi, Mahoba, Panna, Sagar, Chhatarpur,
	Narsingpur, Seoni, Allahabad
Total Pulses	
	Mahoba, Betul

To address the production related constraints amongst the pulse growers, usually with low socio-economic status (SES), poor resource base and least exposure to human resource development (HRD), followings may be suggested:

- i) Strong Development ó Research interface need to be in place to intensify research efforts to evolve still high yielding varieties and management recommendations suited to dry farming/moisture-stress conditions/utera under rice-fallow areas and for different agro-ecological situations (AESs).
- ii) There is need to evolve crop-management modules and low cost technology with best inter-cropping recommendations for various agro-climatic and agro eco-situations). These modules may be helpful to meet-out any contingent situation associated with such production constraints.
- iii) State Agriculture University/Agriculture Colleges/Zonal Research Station (ZRSs)/Krishi Vigyan Kendres (KVKs), etc. in consultation with the State Department of Agriculture now need to develop season-wise nutrient-use efficiency (NUE) plan for each districts on AES basis. Instead of simple recommendations of fertilizers based on the nutrient management practices, there is need to group and plan the practices as:
- Match between nutrient supply from soils and demand by crop on the basis of soil testing and optimization of split fertilizer application and soil and plant nutrition factors (soil moisture, pH, temperature, physical properties etc.).
- Improving nutrient application methods such as broad-casting, band placement, split application).
- Improving physical properties of fertilizers and use of inhibitors to reduce losses.
- Improving soil conditions, crop and water management practices, tillage, regulating soil moisture regimes, crop-rotations, weed control, residue management, break and catch crop etc.
- iv) To be more serious on the sustainability of cropping system and judicious use of natural resources in the rainfed regions, depleting ground water level and frequent drought, State Department of Agriculture may draw the successful experience/results from within the best districts.
- v) State may put a system and policy frame for pulses cultivation. This strategy would not only benefit the small and marginal pulse growers but would prove a boon to statesø proposed crop-diversification programmes involving horticulture etc.
- vi) Liberal credit policies and extending insurance cover under PMFBY with low premium offered by the Government of India also need to be aggresively addressed by the states.
- vii) State Department of Agriculture, in view of the state potential in a particular/group of pulse crop, may constitute a 'Pulse Board' (similar to Tur Board in the state of Karnataka) and procurement policy adopted by A.P. involving private sector, NGO etc, to seriously watch the interest of pulse producer.

viii) The Pulse Boardø could be a multi-disciplinary approach agency taking full care of marketing, domestic level processing, pricing, value addition, Import-Export, and consumption behaviour of states socio-economic-group of farmers.

16.5.3 Marketing Related Interventions

To motivate the pulse growers of different socio-economic-status (SES) in various agroeco-situations (AES) of the state, following interventions may be suggested.

- i) To minimize the price-gap in the chain of producer to consumers, it is important to assign active role and accountability to some institutional buyer like cooperatives, civil supplies, MARKFED etc. State Government may fix a procurement target of at least 20% of the total production in order to build an effective a purchase and price security environment.
- ii) The SDAs should strongly put-forth its procurement share during the all India rabi and kharif procurement meetings organized at the behest of National Agricultural Marketing Federations Ltd. (NAFED), Govt. of India, New Delhi.
- iii) The targetted pulses within the purview of Price support Scheme (PSS) are pigeonpea, gram, lentil, pea, mungbean and urdbean. In view of its major production share in the country, states need to strongly pursue its position to central nodal agency (DAC) for recommendation of more cash credit limit (CCL) to NAFED to be sanctioned by RBI through SBI (up to 75% of hypothecation of stock keeping a margin of 25% in accordance to banking norms).
- iv) State Marketing Federations can also initiate a similar PSS system in the larger interest of pulse growers by way of provisioning a revolving fund commensurate to proposed procurement.


16.5.4 Extension Related Interventions


Monitoring of pre-TMOP and post-TMOP projects (NPDP/ISOPOM) including ongoing NFSM-Pulses by the Directorate of Pulses Development, Bhopal conclude that pulse growers are usually resource poor, small and marginal group of farmers. The socio- economic status (SES) of this group inhibits them to have an immediate access to technology in put. It is, therefore, in the interest of this group in particular and the enhancement of pulses production and nutritional security of the country in general, under mentioned are suggested:

i) For strengthening technology dissemination and extension education, potential pulse producing districts/blocks should be identified. In each block, FPOs constituted during XIth and XIIth plan group of progressive farmers, FPOs, SHGs, Cooperatives, NGOs, KVKs, FIGs, Womenøs Group; Agri-business Companies and Input dealers etc should be organized, strengthened to function as local information kiosks or extension education points.

- ii) The district agriculture officer (DDA) should facilitate these private sectors in terms of local news papers, departmental scheme details, technical literature, credit and insurance consultancy, TV/internet facilities etc through on-going central sector or centrally sponsored, State Government run programmes, banks and input dealers in the field of fertilizers, seeds, pesticides, implements etc.
- iii) DDA/SDO/ADA to facilitate the group in organizing the meetings at common panchait place, developing of Kharif, Rabi and Zaid **crop-cultivation seasonal action plan** clearly indicating the input requirements. The district administration should also provide all administrative/technical input and help in interactions with all other stake-holders or service providers.
- iv) A certain percentage (10-15%) of total allocated developmental programmes (central sector/centrally sponsored/state-run) should be assigned to these identified groups (agents). Block demonstration, IPM demonstrations, production of certified seed etc components may also be given to these agencies for more accountability and ownership feelings.
- v) Under the varietal diversification programme commonly known as seed minikit distribution under the ongoing NFSM programme, at least 10% of the minikits, alongwith the technology package, be given to these FOs/SHGs/FIGs/NGOs. The SDA may also start their own seed minikit programme.
- vi) Each potential block is identified as processing centre and at least one small/domestic dal mill like IIPR dal chakki, CIAE Dal mill may be provided. The responsibility of running the mill is rest with the NGOs/Farmers Organization.
- vii) Methodologies and package of practices for improving fertilizer use efficiency (FUE) under various soil conditions and different crops, as brought out by Indian Council of Agriculture Research (ICAR) be documented in vernacular language by the state Directorate of Agriculture under the funds on publicity provided through NFSM-pulses and made available to these groups by the district agriculture officer/farmers.


POST HARVEST TECHNOLOGY AND MANAGEMENT





SMALL SCALE MILLING OF PULSES

MINI DHAL MILL (CFTRI)







MINI VERSATILE DHAL MILL

Chapter-17

Post-Harvest Technology and Management

Post-harvest protection of pulses assumes a greater importance in overall crop protection system as pulses are more susceptible to storage losses. Traditionally the produce is essentially stored for longer or shorter duration, either for consumption or as seed for sowing during the next cropping season.

17.1 Harvesting Precautions

To minimize quantitative and qualitative losses, besides harvesting the crop at 08 per cent of total pods maturity stage, under mentioned. *Advisory should be followed:*

- Harvesting prior to physiological and proper maturity usually result in lower yields, higher proportion of immature seeds, poor grain quality and resulting in susceptibility to infestation during storage. To fetch better prices and consumer acceptance, proper harvesting judgement is required.
- Avoid harvesting during adverse weather conditions i.e. rains and overcast weather, however, delay in harvesting may results in shattering pods and losses caused by birds, rats, insects etc. Rogue out the admixtures prior to harvesting.
- The harvested produce should be stacked in a dry, clean place in cubical way to facilitate circulation of the air around and keep the bundles for drying in the field after cutting on threshing floor.

17.2 Grading

Sorting of the homogenous lots of the produce according to the fixed grade standard in accordance with various quality factors is important. Grading of the produce before sale enables farmers to get better price and helps the consumers to get standard quality produce at fair price vis-a-vis facilitate the consumer to compare the prices of different qualities of a produce in the market. Grading assures the quality of the produce and also reduces the cost of the marketing and transportation. The quality parameters of pulses are wholesome, clean, odorless and less moisture content.

17.3 Storage/Packaging

The good packaging material must protect quality and quantity, prevent spoilage during transit and storage and should display about grade/quality, variety, date of packing, weight and price etc.It must also be convenient in handling operations, convenient to stack, cheap, clean and attractive.

Impotant packaging materials are (i) Jute bags, (ii) HDPE/pp bags, (iii) polythene impregnated Jute bags, (iv) poly pouches, and (v) cloth bags. About 10-15 per cent moisture is safe for storage of pulses. For small-scale storage, preferably air-tight

metallic bins, and for large scale storage of pulses, large silos are commercially available. The storage affects the cooking quality of whole and split pulses (dal).

17.4 Major Stored Grain Pests

The various factors responsible for deterioration of stored grains/seeds are broadly classified under two categories, biotic factors (insect, rodents, birds, fungi, mites and bacteria); Abiotic factors (moisture content/relative humidity, temperature) Pulse beetle (*Callosobrucus maculatus* (Lin), (Bruchid) in whole grain and *Tribolium castaneaum*, *Tribolium confusum* in milled product (besan), are the major stored insect.

• Prophylactic and Curative Measures

Selection of site, storage structure, cleaning and drying of structures/site/bags is important. The site/structure should be given *prophylactic treatment* by spray of Malathion 50% EC (1:100) one per cent solution @ 3 litres per 100 sqmt.

For *curative treatment*, Methyl Bromide and Aluminium phosphide are common fumigants. Aluminium phosphide @ 3g pallet per 5-10 qtls whole grain for 7 days is recommended. Control of rodents should be done through i) Multi-dose anti-coagulant (cumarin compounds) eg. *Rodaferin*, *Warferin* (proportion 1:19) ii) single dose anti-coagulant eg. Promadiotone (proportion 2:98) and iii) single dose acute poison eg. *Zinc phosphide* (proportion 2:98).

17.5 Storage Structures

Producers store pulses in bulk at farm godown or own house using various types of traditional and improved structures. Generally, these storage containers are used for short period. Different organisations/institutions have developed improved structures for pulses storage with various capacities like Hapur Kothi, Pusa bin, Nanda bin, PKV bin, etc. Different storage structures are also used for this purpose like bricks-built rural godown, mud stone godown etc. Producers also use flexible PVC sheets covering for temporary storage. Some producers also pack pulses in jute gunny bags or in gunny bags lined with polythene and stack in room.

Prevalent storage structure may be classified into two categories as domestic and commercial (Table 16.15).

(Table-17.1): Categories of prevalent storage structures

Domestic	Commercial	
Traditional structures	i.Warehouse CAP Storage (cover	and plinth
i.Mud-binds or Kachchi Kothi	storage) Soils.	
ii Metal drums	ii. Steel Silos	
iii Thekka		
iv Gunny bags		
Improved/scientific structures		
i Pusa Kothi		
ii Nanda bins		
iii. Hapur Kothi		
iv PAU bins		
v PKV bins		
vi Chittore stone bins		

17.5.1 Storage Infrastructure/Programmes/Facilities

I) Rural Godowns

- Considering the importance of rural storage in marketing of agricultural produce, DAC&FW, Directorate of Marketing and Inspection, initiated a Rural Godowns Scheme, in collaboration with NABARD and NCDC. Its objective is to construct scientific storage godowns with allied facilities in rural areas and to establish a network of rural godowns in the States and Union Territories.
- Eligibility: The project for construction of rural godowns can be taken up by individuals, farmers, group of farmers/growers, partnership/ proprietary firms, non-government organizations (NGO), self help groups (SHGs), Companies, Corporations, Cooperatives, Agricultural Produce Marketing Committees, Marketing Boards and Agro Processing Corporations in the entire country. However, assistance for renovation/expansion of rural godown is restricted to the godowns constructed by co-operatives only.
- Location: Under the scheme, the entrepreneur will be free to construct godown at any place and of any size as per his commercial judgment except for the restrictions that it would be outside the limits of Municipal Corporation area and be of a minimum capacity of 100 MT.

II) Mandi Godown

• Most of the States and Union Territories have enacted Agricultural Produce Market (Regulation) Act. The reduction of loss of produce was aimed in the scheme of regulated market. The regulated markets developed modern market yard with necessary infrastructural facilities. The APMCs have constructed godowns so that the agricultural produce brought into the market should be stored safely by market committees. The produce is weighed in the presence of producer/seller at the time of keeping the produce in the godown after grading for storing and receipt is issued indicating the quality and weight of produce to be stored. The receipt is issued by the licensed general commission

agents or brokers depending upon the case. The CWC, SWC and Co-operative societies have also constructed godown in the market yards.

• In most of the secondary and terminal regulated markets, central and state warehousing corporations also provide scientific storage facilities at prescribed storage charge and issue *warehousing receipt against pledge of produce*, which is a negotiable document for obtaining finance from the scheduled banks.

III) Central Warehousing Corporation (CWC)

• CWC was established during 1957. It is the largest public warehouse operator in the country. Apart from storage, CWC also offers services in the area of clearing and forwarding, handling and transportation, distribution, disinfestation, fumigation and other ancillary services like safety and security, insurance, standardization and documentation. The CWC has also introduced a scheme, called the FarmersøExtension Service at selected centres to educate farmers about the benefits of a scientific storage. The CWC is also operating custom bonded warehouses. These bonded warehouses are specially constructed at a seaport or airport and accept imported commodities for storage till the payment of customs duties by the importer of the commodities.

IV) State Warehousing Corporations (SWCS)

Different states have set up their own warehouses in the country. The area of operation
of the state warehousing corporations is district places of the state. The total share capital
of the state warehousing corporations is contributed equally by the Central Warehousing
Corporation and concerned State Government. The SWCs are under the dual control of
the State Government and the CWC.

V) Co-Operative

• Co-operative storage facilities are provided to the producer at cheaper rates, which reduces the storage cost. These Co-operatives also provide pledge loan against the produce and storage is more systematic and scientific than traditional storage. Financial assistance and subsidies are provided by government organisations/banks to build Co-operative storage. To meet the increasing need for storage capacity, the National Co-operative Development Corporation (NCDC) encourages construction of storage facilities by Co-operatives, particularly at rural and market level.

17.6 Marketing Channel

The production of a produce is complete only when it reaches the hands of consumers. Marketing channels are the routes through which agricultural products move from producers to consumers. A flow of pulse produce from farmers to consumer under organised and un-organised channel is exhibited under Table 17.2.

(Table-17.2): Processes of marketing of raw produce.

Private	Institutional		
i) Producer Dal Miller Consumer	i) Producer Procuring Agency Dal		
ii) Producer VillageTrader DalMiller Wholesaler	Miller Consumer		
Retailer Consumer	ii) Producer Procuring Agency		
iii) Producer Dal Miller Retailer Consumer	Dal Miller Wholesaler		
iv) Producer Wholesaler DalMiller Retailer	Retailer Consumer		
Consumer	iii) Producer Procuring Agency		
v) Producer Wholesaler DalMiller Wholesaler	DalMiller Retailer Consumer		
Retailer Consumer			
vi) Producer Wholesaler Retailer Consumer			
(For whole Green gram)			
vii) Producer Commission Agent Dal Miller			
Wholesaler Retailer Consumer			

17.7 Processing and Value Addition

- Promoted by the western habits, food consumption habit is under radical change in India. Export of value added products has retained the upward ladder. Food Processing industry is still at the category of small or cottage industries. The Industry has to be popularized due to the wide range of consumption of processed items of foodstuffs in the country. A close study on the issue imperatively amounted to a conclusion that the industry is not less important than the bigger industrial units on various consumer and non-consumer goods. Due emphasis has to be paid to the agro-industries based on the prevailing nature of perishable crops including pulses.
- Lack of Processing Technologies of applied nature has a far negative reaching implication vis-à-vis, value addition and by-product utilization of pulses. There is a great scope of canning fresh peas, but lack of facilities for preservation has not made much headway, particularly in the rural sector where all types of infrastructures are not available. The produces of the farmers are sold in situ at low prices hardly meeting the economical aspirations of the farmers themselves. Middlemen involved in the process of transporting the pulse grains exploit the rights of the farmers to sell them at reasonable prices. Lack of processing plants in the vicinity of the farmersø field have encouraged these the middlemen to interfere for converting the raw and fresh pulses into various processed food items and their by- products, where value additions are the prime target.
- In an effort to increase the value of foodstuffs in pulses, research & development on Post Harvest Technologies (PHT) would come to play a major role. The large loss of pulse grains during harvest operation and post harvest storage (25-30%) is a major concern. PHT is thus, an indispensable part of operation when food processing and value addition of pulse crops are concerned.

17.7.1 Domestic/Small Scale Pulse Milling in rural sector: Scope

- Setting up small scale pulse milling units in rural sector need to be expoited to boost-up the pulse sector. Although dal milling is an agro-based industry, the rural sector is rather deprived of this owing to following reasons common to the rural areas of the country: Non-availability of infrastructural facilities in rural sector; Inefficient methods of milling in rural sector (incompetent methods and machines for processing dal in rural sector); Non-availability of suitable cottage scale milling machines which are economical and can be easily adopted in rural sector with the existing methods of processing.
- Moreover, the capital investments, taxation policies, lack of skilled labour are coming in way of setting up a dal mill in rural sector. The producer, therefore, is almost forced to sell the pulses to the agent-cum-dal miller in large scale sector and in turn purchase dal from him, thereby giving him major share of profit.
- Keeping in view these difficulties some organization like CFTRI (Mysore), PKV (Akola, Maharashtra), IARI,ICAR (CIAE, Bhopal), have come up with several designs of small scale/cottage scale pulse dehusking machines, with capacities ranging from 40 kg ó 200 kg per hr. These low cost, low capital investment machines may help the producer to get value added product (dal), and useful by product óChunni-and husk for his cattle. These machines can be easily operated and maintained by a single family or by a village based small cooperative society either for their own use or as custom milling systems, thereby giving chance for more rural employment. This may have an impact on the overall village economy especially in the major pulse growing regions.

17.7.2 Benefits of mini/small scale mills

- Simple technology/mini machines easy to operate & maintain and repair by villagers; Low cost of processing and less power consumption; Low capital requirement, hence, can come within the limits of state financial corporations or KVIB of states; Can attract subsidy by State Governments and avoid taxation to some extent; Long distance transportation is not required, since raw material purchase and product sale are confined to local markets.
- The scope for setting up such small scale pulse dehusking machine is based on (i) the type and utility of the machine for the pulses grown and (ii) the status of pulse milling industry in that area. It is assumed that a small scale pulse dehusking unit like CFTRI mini dal mill processes about 5 quintals of pulses in a day on an average. If it works for 150 days in season (December to May), it can process 75 tonnes of pulses in one year. Assuming 50% of the produce is retained and processed to dal in rural sector, 6 such units can be set up in a district where the production is about 1000 tonnes (500 tonnes available for small scale). The number of such small scale units suggested to be set up in a district is based on the above assumption.

17.8 Processing Technology

- India is the largest producer and consumer of pulses in the world. Processing of dal is unique and indigenous to India. This is due to the fact that, substantial quantities of pulses are consumed in the country in the form of dal ó the dehusked split form. Though pulse milling is the third largest grain processing industry in the country, next only to rice and wheat, processing still remains largely traditional and employs empirical methods of processing which leads to inefficient processing and wastage of precious raw materials.
- Processing of pulses into dal or a variety of primary and secondary products adds more values to consumers. However, the operation is being coupled with losses and wastage estimated to be about 10-25%, depending on the technology adopted and machines used. Still, the processing of pulses is on the rise due to the consumersøneeds and the sound market price of processed pulse products. In India, more than three fourths (3/4th) of pulses produced are processed into dal. During the processes of milling only the losses (as powder and brokens) are estimated to be about 10-15%. Excessive scouring of pulse grains not only results in quantitative loss, but also qualitative loss since the peripheral layers contain substantial quantity of proteins. It is therefore, due to this that care must be taken to minimize the losses by using improved machineries and processing techniques.

17.8.1 Large Scale Processing

• As the traditional methods are laborious, time consuming and dependent on climatic conditions, attempts have been made to develop new technologies for efficient and economic milling of pulses. An improved method and machinery was developed by CFTRI in eighties which aims at minimizing the difficulties faced by traditional large scale pulse processors. The improved method gives a higher yield of dal in lesser time and at a lower cost of processing. The process is accomplished in two steps. In the first step, loosening of husk is achieved by an incipient toasting followed by tempering and the removal of husk and splitting is achieved by improved processing machines. The method consists of exposing the cleaned and size graded pulse, followed by tempering in bins to a critical moisture level. Removal of husk is done in an improved pearling machine in a single operation. The gota is split in an impact splitter after moisture treatment and aeration under controlled conditions. The method is independent of climatic conditions and can function throughout the year resulting in increased productivity. The technology has already been released to the industry.

17.8.2 Small Scale Processing

In order to revive the now-defunct traditional village level industry and to place the rural
dal processor on a competent and sound economic and technological footing, CFTRI
has recently developed an integrated smallscale pulse processing unit –Mini dal
Mill. This consists of a dehusking unit, an aspirator and a reciprocating sieve, all
run by a 1 HP motor. The mini dhal mill can process 100-150 kg of pre-conditioned

pulse per hour without causing much breakage and powdering. Dehusked split dal husk and brokens are collected at different points as in big dal mill. The product quality is comparable to that of commercial dhal mill and dal yield is 78-82%. The cost of processing is also low. This unit is highly suitable for dehusking of bolder grains like arhar (tur), bengal gram, peas, soybean, field bean etc, while only splits (unhusked dal) could be obtained from green gram and black gram.

• The pre-milling treatment as practiced in rural technology (soaking and sun-drying) is retained, since it is easily carried out on rural surroundings. However, duration of soaking is standardized to suit the variety of pulse. Since not all the grains soak uniformly, separation of soaked and swollen grains is essential in order to get good quality product. For this a specially designed grader also has been developed for grading the soaked pulse which can also be used as a pre-cleaner-cum-grader. This unit is run by a half HP motor.

Common pulse processing avenues

The Pulses can be processed and used in the following ways:

- Cooking
- Dehulling -Dal
- Germination- Cooking
- Puffing
- Cooking-Sambar
- Wet grinding- Idli, Vada, Dosa
- Dry Grinding- Sev, Bajji, Bonda
- Some eaten raw

> Problems of traditional dal milling industry

There are about 7000 registered dhal mills in India and about 5000 small or cottage scale dhal mills. The problems are \acute{o}

- Long processing time for pulses (5-6 days)
- Lower yield of dal (72-74%) and more broken (12-15%)
- Lack of skilled labour / trained personnel
- Dust pollution

> Factors influencing the milling are –

a) Raw Material Characteristics

- Size and shape of pulses
- Husk content and its thickness
- Adherence of husk to the cotyledons
- Moisture content of the grains
- Extent of infestation

b) Pre-Milling Treatment

• Wet pre-milling treatment (Soaking in water & Sun drying)

• Dry pre-milling treatment (Pitting, oil mixing, Sun drying, water addition, Sundrying).

17.9 Domestic Machineries Developed

Under the R&D in PHT on Oilseeds, Pulses and maize Mini Mission II of TMOP, domestic processing machines were developed. CFTRI, CSIR, SAUs and ICAR institutes, under this programme, developed processing technologies.

(Table-17.3): Processing technology developed under R&D in PHT

Name of the Domestic	Brief Features	Advantages
Machinary/(Institutes)		
Mini Dal Mill (CFTRI, Mysore)	 Application - Promotion of village dal milling by traditional rural processors Capacity-100 to 150 Kg / hr. Space - 2 x 4 meters Power: - Mill - 1.0 HP	 Easy to operate, maintain and repair Simple pre-milling treatment Low capital investment Ideal for Cottage scale rural industry By-products useful as cattle feed Low cost processing Supplied under subsidy programme
Name of the	Brief Features	Advantages
Domestic Machinary		
(Institutes)		
Versatile	• Capacity: 250-300 Kg / hr.	Suitable for small scale
Dal Mill	• Power required: 15 HP	processing
(CFTRI, Mysore)	 Space Required: 8 x 12 Meters Utility: Can process all types of pulses Dehulling: 98-99% Yield of dal: 75-78% Breakage: 2-3% 	 Good quality dal at competitive price By-products ó valuable animal feed Transportation cost reduced Employment generation Filling to advance technology base for rural processing
Modern Dal Mill (CFTRI, Mysore)	 Capacity: One tonne per hr. Power: 100 HP (Including 60 HP for Electrical for Heating and conditioning) Space: 15 x 30 Meters Utility: Can process all types of pulses Processing Time: Less than 2 days 	 Independence from climatic conditions Higher recovery of dal Automatic process for round the clock production Reduced time of processing

Table Gota Separator (CFTRI, Mysore)	 Yield of dal: 77-80% Dehusking: 98-99% Utility: Can separate gota (pearled tur from whole grain) Principle: Works on surface 	A A	Suitable for incorporation in large scale dal mills Additional annual recovery of 8 tonnes of
	 Frinciple: Works on surface resilience differences of grains Capacity: 500 kg/hr. Power: 2 KW Space required: 4 x 4 meters 	>	first grade dal.valued Rs.2 lakhs. Saving of power to the tune of 20%
Hand-Operated Pulse Dehusker (CFTRI, Mysore)	 Capacity ó 40 kg per hour Power ó Nil Utility ó can process bold pulses, suitable for Home/cottage scale 	A A A	Suitable for small scale processing Good quality dal at competitive price By-products ó valuable animal feed

17.10 Milling methods of pulses

In India there are two conventional pulse milling methods; (i) wet milling method, and (ii) dry milling method. The latter is more popular and used in commercial mills.

17.10.1 Traditional dry dal milling method

There is no common processing method for all types of pulses. However, some general operation of dry milling method such as cleaning and grading, rolling or pitting, oiling moistening, drying and milling have been described below:

- Cleaning and grading: Pulses are cleaned from dust, chaff, grits, etc., and graded according to size by reel type or rotating sieve type cleaner.
- Pitting: The clean pulses are passed through an emery roller machine. In this unit, husk is cracked and scratched. This is to facilitate the subsequent oil penetration process for the loosening of husk. The clearance between the emery roller and cage (housing) gradually narrows from inlet to outlet. As the material is passed through the narrowing clearance, mainly cracking and scratching of husk takes place by friction between pulses and emery. Some of the pulses are dehusked and split during the operations which are then separated by sieving.
- **Pre-treatment with oil:** The scratched or pitted pulses are passed through a screw conveyor and mixed with some edible oil like linseed oil (1.5 to 2.5 kg/tonne of pulses). Then they are kept on the floor for about 12 hours for diffusion of the oil.
- Conditioning: Conditioning of pulses is done by alternate wetting and drying. After sun drying for a certain period, 3-5 per cent moisture is added to the pulses and tempered for

about eight hours and again dried in the sun. Addition of moisture to the pulses can be accomplished by allowing water to drop from an overhead tank on the pulses being passed through a screw conveyor. The whole process of alternate wetting and drying is continued for two to four days until all pulses are sufficiently conditioned. Pulses are finally dried to about 10 to 12 per cent moisture content.

- Dehusking and splitting: Emery rollers, known as Gota machine are used for the dehusking of conditioned pulses. About 50 per cent pulses are dehusked in a single operation (in one pass). Dehusked pulses are split into two parts also. The husk is aspirated off and dehusked, split pulses are separated by sieving. The tail pulses and unsplit dehusked pulses are again conditioned and milled as above. The whole process is repeated two to three times until the remaining pulses are dehusked and split.
- **Polishing:** Polish is given to the dehusked and split pulses by treating them with a small quantity of oil and/or water.

Flow-Diagram of milling process is indicated in box




Fig.-17.1: Flow diagram of dry milling of pulses

17.10.2 Milling techniques of different pulses

Pulses like tur, black gram, green gram and horse gram are generally difficult to dehusk
while pulses like Bengal gram, peas, lentil and khesari are easy to dehusk. This
difference in milling behavior is mainly due to the extent of adherence of the husk to the
cotyledon. Actual commercial practices generally followed for some of the individual
pulses are described as follows:

i) Dehulling of Tur (Arhar)

• Arhar poses greatest difficulty in milling since the husk is tightly adhered to the cotyledons. Generally only dry method is followed throughout the country for milling of arhar or tur. Fig. 1 gives a flow chart for its milling. The cleaned and size graded grains are pitted in smooth roller machines smeared with oil (0.2 \(\dectrice{0} \) 0.5%) (linseed, cashew or

any other cheap oil) tempered for about 12-24 hours, sun dried for 1-3 days, followed by spraying with water (2-3%), thoroughly mixed, heaped overnight and then passed through the rollers for dehusking. This type of operation is repeated 3-4 times. After each dehusking operation, the husk, powder and brokens are separated from dhal and gota (mixture of dehusked and unhusked grains). The dhal thus obtained is considered as II grade since its edges are rounded-off due to scouring. The gota obtained is again mixed with water as above, equilibrated and sun dried. The sun dried gota is either passed through the roller machine or split in horizontal or vertical chakki or using a patka machine. The dhal obtained from the gota is considered as I grade dhal since it does not have any chipped edges and has a better consumer appeal. In some places both I and II grade dals are mixed and marketed. The yield of dhal varies from 70 to 75% depending upon the variety and the method followed. The present survey has revealed that in large scale mills sun drying is being replaced gradually with batch type bin drier. As a result these units are able to work throughout the year.

Processing of Arhar is mainly done in the states of Madhya Pradesh, Chhattisgarh, Uttar Pradesh, Maharashtra, Gujarat, Karnataka, Tamilnadu, Bihar and Uttrakhand.

Dehulling of Tur – Large Scale (Wet Method)




Fig.-17.2: Dehulling of tur-large scale

• Dehulling of Tur-Small Scale (Dry Method)

Small scale duhulling of tur are following the two process viz. (i) dry method and (ii) wet method. Dry and wet method at small scale is exibited under *flow-diagram fig. 2 and fig 3*.




Fig.-17.3: Dehulling of tur- dry method




Fig.-17.4: Dehulling of tur- wet method

ii) Dehulling of Bengal Gram (Chickpea)

This pulse is comparatively easy to mill. The cleaned and size graded grains are pitted in smooth rollers at low peripheral speed. After pitting the grains are mixed with about 5% water in a mixer and heaped for a few hours to allow the water to seep in. The wetted grains are sun dried for a day or two. The dried pulse is then passed through either horizontal or vertical chakki. Here dehusking and splitting take place simultaneously. The dhal is separated from the husk and brokens. Any remaining unhusked grains are dehulled by repeating the above operation till all the grains are dehulled. Processing of Chickpea is confined mainly to Rajasthan, Delhi, Uttar Pradesh, Madhya Pradesh, Chhattisgarh and Maharashtra (Fig.4).




Fig.-17.5: Process for dehulling of Bengal gram

iii) Dehulling of Black Gram

The cleaned and size graded grains are pitted using emery rollers in 2 or 3 passes, so that complete pitting is effected. After each pitting operation the husk and powder is separated. The pitted grains are then mixed with about 0.5% oil and heaped overnight for absorption. The grains are then sun dried for 2 days. In some mills mechanical dryers are used. After drying, the grains are given a spray of water (2 to 3%), equilibrated and passed through the rollers twice for dehusking. The split dhal obtained is termed as II grade dhal. The dehusked <u>gota</u> is passed through Burr mill for splitting. The dhal obtained from <u>gota</u> is considered as Ist grade dal. The split dhal is õpolishedö with soapstone powder at the final stages. This is believed to give luster to the dhal and enhance their market value.

Processing of Urdbean in the states of Andhra Pradesh, Odisha, Tamilnadu, Karnataka, Maharashtra, Chhattisgarh, Madhya Pradesh, Uttar Pradesh and Delhi.




Fig.-17.6: Process for milling green gram and blackgram

iv) Dehulling of Green Gram

The husk of green gram is thin, soft and slippery. While the husk is tightly adhering to the grain surface, the two cotyledons are loosely attached and separate out easily. Hence, splitting into dhal occurs even before good dehusking can be effected. During the dehusking operation, there is also scouring of the cotyledons resulting in large losses in the form of broken and powder. The method generally followed is pitting, oiling (0.2-0.5%), sun drying followed by dehulling and splitting in roller machines (fig.5). In some states like West Bengal, because of the demand for smaller sized Dhal, general practice is to go on scouring the Dhal which results in loss of valuable proteinaceous material in the form of powder.

Processing of green gram is largely done in Rajasthan, Madhya Pradesh, West Bengal, Uttar Pradesh, Andhra Pradesh, Odisha and Maharashtra.




Fig.-17.7: Green gram/black gram splits-flow chart

v) Dehulling of Peas, Lentil and Khesari

Processing of these pulses is fairly easy as in the case of bengal gram. General practice involves initial scouring, moisture application, heaping and sun drying, followed by dehusking and splitting in roller machines. After separating the Dhal, the unhusked grain is treated a second time as in the first pass, and repeated till all grain are dehusked and split (Fig.7,8,9).

Processing of lentil is generally practised in Uttar Pradesh, Bihar, Jharkhand, West Bengal, Madhya Pradesh and Delhi while milling of peas (yellow peas) is restricted to the state of Uttar Pradesh, West Bengal, Maharashtra, Delhi, Chhattisgarh, Madhya Pradesh. Khesari pulse is processed mainly in Madhya Pradesh, Chhattisgarh and Bihar States.




Fig.-17.8: Process for dehulling peas




Fig-17.9: Process for milling of masoor (lentil)




Fig.-17.10: Process for dehulling khesari pulse (lakh)



Chapter-18

Policy Interventions

18.1 Projects/programme on pulses development

With the unabated population increase in the Country, pulses production, the main source of protein/balanced diet particularly for the rural mass also thought to be paralleled in proportionate to population growth. Accordingly the Department of Agriculture, Cooperation & Farmers Welfare launched various development programmes on pulses during different Plan periods.

Plan interventions in the pulses sector were brought by the Govt. of India, Department of Agriculture, Cooperation & Farmers Welfare since Fourth Five year Plan with more focused approach since VIth Plan onwards as under:

õPulses Development Schemeö a Centrally Sponsored Scheme, was initiated from the IVth Plan (1969-70 to 1973-74). The focused area was the introduction of production technologies and improved varieties amongst the farmers.

- Seventh plan (1985-90): conceived the National Pulses Development Project (NPDP), merging all the earlier centrally sponsored schemes on pulses.
 To further supplement the efforts under NPDP, a õSpecial Food Grain Production Programme (SFPP) on Pulsesö was also implemented during 1988-89 on a 100% Central assistant basis.
- Technology Mission on Oilseeds (TMO 1985-86): To ensure the accelerated development of certain priority areas of economic and social concern, the Government of India adopted a compressive approach and launched *Six Technology Missions*viz. i) Rajiv Gandhi National Drinking Water Mission ii) Immunization Mission iii) National Literacy Mission iv) Tele-communication Mission v) Dairy Development (Operation Flood-II): and vi) Maximization of indigenous production of vegetable oilseeds/oils etc.
- For accelerated development and successful implementation of the mini-missions approach, three strategic Committees were also set up for Structural innovation viz. (i) Empowered Committee (EC) (ii) Technical and Advisory Committee (TAC) (iii) Standing Committee (SC).
- The TMO remained operational under the supervision of ICAR till 1987-88. From 1988-89 onwards, the implementation and responsibilities were transferred to Department of Agriculture and Co-operation to harness the best of production, processing management technologies harmonizing the interest of farmers, consumers and accelerate self-reliance in oilseeds and edible oils. The TMO pursued a Mission-Mode-Approach by forming a consortium of concerned department and stake holders.

- TMOP (1990-91): Pulse development programmes were brought to the ambit of the Mission in August 1990. Thereafter Oilpalm (1992-93) and Maize (May,1995) also became the part of it, renaming the TMO as Technology Mission on Oilseeds, Pulses and Maize (TMOP&M). The Seventh Plan ongoing interventions under National Pulses Development Project (NPDP) became the part of TMOP&M.
- TMOP&M had four-pronged strategy approach under its four Mini Missions involving the concerned department and agencies to facilitate the task of handling specialized focused areas of development viz. MM-I Crop Protection Technology: DARE with ICAR as nodal deptt., Department of Bio-Technology and SAUs as implementing agencies; MM-II Post Harvest Technology: Department of Scientific & Industrial Research with CSIR as nodal deptt. and Department of civil supplies as participating agencies; MM-III- Input and service support to farmers: DAC as nodal agency with SDAs, NDDB, NABARD and NOVOD Board, as implementing agencies and MM-IV-Price support, storage, processing and marketing: DAC as nodal deptt. with participating agencies as NCDC, NDDB, NAFED, Department of civil supplies KVIC and NOVOD Board.
- **ISOPOM** (2004-05 to 2009-10): From April 2004 to March 2010, on the advice of the Planning Commission, õIntegrated Schemes of Oilseeds, Pulses, Oilpalm and Maize (ISOPOM)ö has been under implementation by merging 4 ongoing schemes of NPDP, OPP, OPDP and AMDP. The ISOPOM had a more focussed and integrated approach. To strengthen the market invention and effective pricing policies were some of the added features of this programme.
- NFSM-Pulses (2007-08): From 2007-08 (Rabi), in pursuance of the resolution adopted in 53rd meeting of National Development Council, a Centrally Sponsored Scheme onö National Food Security Mission was launched. It was resolved to enhance the production of rice, wheat and pulses by 10, 8 and 2 million tonnes, respectively by the end of XII Plan. The implementation of the NFSM scheme is continued beyond the XII Plan*i.e.* 2017-18.
- The NFSM aimed at increasing production of rice, wheat and pulses through area expansion and productivity enhancement; restoring soil fertility and productivity; creating employment opportunities; and enhancing farm level economy to restore confidence of farmers of targeted districts. The basic strategies were implementation of interventions in a mission mode through active engagement of all the stake holders at various levels. These interventions includes promotion and extension of improved technologies i.e., Seed, Integrated Nutrient Management (micro-nutrient, soil amendments), IPM and resource conservation technologies along with capacity building of farmers. Flow of fund closely monitored to ensure that intervention reach the target beneficiaries on time, Interventions proposed were integrated with the district plan and target for each identified district was fixed. Constant monitoring and concurrent evaluation were done for assessing the impact of the interventions for a result oriented approach by the implementing agencies.

• NFSM + Special initiatives (2010-11 to 2013-14): To accelerate the pulses production, a centrally sponsored Accelerated Pulses Production Programme (A3P) (2010-11 to 2013-14)-cluster demonstration approach from; Special initiatives for õpulses and oilseeds in dry land areaö under RKVY during 2010-11; Integrated development of 60000 Pulses villages in Rainfed Areas under RKVY during 2011-12 and õSpecial plan to achieve 19+ million tonnes of Pulses production during Kharif 2012-13ö were also been implemented.

Strong Research and Development efforts during XI Plan had spectacular achievement realising more than 20% increase in the production of Pulses at the terminal year of XI Plan (2011-12).

• NFSM-Pulses XII Plan: During 2017-18, the Pulses development scheme under NFSM was under implementation in 29 states viz. Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Chhattisgarh, Goa, Gujarat, Haryana, Himachal Praedsh, Jharkhand, J&K, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Mizoram, Nagaland, Odisha, Punjab, Rajasthan, Sikkim, Tamil Nadu, Telangana, Tripura, Uttar Pradesh, Uttarakhand and West Bengal with additional production target of 4 Million tonnes by the end of XII Plan (2016-17).

During 12th Plan, the NFSM with the other four Missions, viz. NMAET, NMSA, NMOOP & MIDH is continued. The pattern of Central assistance under NFSM has been 60:40 per cent up-till 2016-17.

The Twelfth Plan NFSM (2012-13 to 2016-17), revamped from 2014-15 and is under implementation with five components viz.i) NFSM- Rice, ii) NFSM-Wheat, iii) NFSM-Pulses, iv) NFSM-Coarse Cereals (millets) and v) NFSM-Commercial Crops (Jute, Cotton, Sugarcane).

- A target of an additional production of 25 million tonnes of food grains i.e. from 259.29 MT to 284.29 over the base year of XI Plan (i.e. 2011-12) comprising Rice-10 million tonnes, Wheat 08 million tonnes, Pulses 04 million tonnes & Coarse Cereals-03 million tonnes, is targeted to be achieved at the end of 12th Plan (2016-17).
- The existing Centrally Sponsored Scheme have also been rationalized and 03 schemes viz. (i) Krishi Unnati Yojana (ii) National Crop Insurance Programme (NCIP) and (iii) Pradhan Mantri Krishi Sinchai Yojana (PMKSY) are operational since 2015-16. NFSM-2015-16 is a part of KrishiUnnatiYojana (State Plan). From 2016-17, the revamped NFSM under State Plan Scheme ó Krishi Unnati Yojana (State Plan) with interim sharing pattern of 60:40 for plains and 90: 10 for hilly states between Centre and State is under implementation in 29 states. A Central Share of Rs. 1700 Crores has been approved during 2016-17.

- The basic strategy of the Mission is to focus on low productivity high potential districts, promote and extend improved technology package, implementation of cropping system centric interventions on technological package, agro-climatic zone wise planning and cluster approach demonstrations, Further 30% of total demonstrations would be Cropping System Based Demonstration (CSBD) with technical backstopping of ICAR/State Agricultural Universities (SAUs)/ on Rice, Wheat, Pulses; distribution of certified HYV seeds/Hybrid seeds, Resource Conservation Technology (RCT) tools, irrigation machineries/MIS, trainings and undertaking local initiatives to the tune of 9% of total budgetary allocation to improve productivity.
- Special emphasis has also to be given by targeting reclamation of problematic soils, water logging areas and mitigation of adverse effects of climate change for high productivity areas, value chain integration (FPOs), assistance to Custom Hiring Centre (CHCs). 30% of budgetary allocation has to be earmarked for women beneficiaries.
- To ensure equity, of the total budgetary allocation to a district proportionate expenditure under Special Component Plan (SCP) for SCs, Tribal Sub Plan (TSP) ó SMF and Women farmers at 16%, 8%, 33% and 30% respectively is mandatory.
- Assistance for various interventions like cluster demonstrations on improved package of
 practices, demonstrations on cropping system, cropping system based training of
 farmers, seed distribution of HYVs, manual sprayer, power sprayer, tractor mounted
 sprayer, chiseller (deep ploughing), water carrying pipes, mobile raingun, sprinkler set,
 pump set (up to 10 HP), seed drill, zero till seed drill, multi crop planter, zero till multi
 crop planter, ridge furrow planter, rotavator, multi crop thresher, laser land leveller,
 plant protection chemical and bio pesticides, weedicides, gypsum/phospho-gypsum,
 bio-fertilizers, micro nutrients, local initiatives are provided under NFSM-Pulses
 programme.
- Concerted efforts are being made for promotion of cultivation of pulses as inter-crop
 with cereals, oilseeds, commercial crops.At least 30% of the cluster demonstrations
 under NFSM and BGREI are being conducted by adopting cropping system approach to
 promote pulses as second crop in rice fallow areas.
- Formation of Farmer-Producer Organizations (FPOs) is also being promoted particularly to support the small and marginal farmers to offer collective strength for seed production, procurement and access to improved technologies. Besides, for primary processing of pulses, assistance is provided for establishment of mini *dal* mills under NFSM. State Agriculture Universities/ Indian Council of Agricultural Research Institutes/ International Research Organizations are also involved to address various researchable issues of pulses and demonstrations of latest technologies for better yield realization at farmersøfield.

- Government of India has allocated Rs.2201.23 crores (CS-1395.00 Cr + SS-806.23 Cr) for NFSM for 2017-18, out of which an amount of Rs. 1371.11 (CS- 850.00 Cr + SS-521.11 Cr) crores is earmarked for pulses.
- FLD on Pulses through ICAR-IIPR: Rs. 0.97 crore
- Establishment/Strenghtening of Biofertilizer and Bio-control Production Units (24 centres/Institutes): Rs. 29.61 crores

• *FPOs: (111)* : Rs. 52.1084 crores

• **Seed Minikit**: Total allocation Rs. 150 Crore

(Table-18.1): Plan-Wise Intervention (VIII th to XII th Plan)

Sr.	Plan Period (VIII th To XII th Plan)	States
No.		Covered
	VIII th -IX th and X th Plan	
1.	National Pulses Development Project (NPDP) (1990-91 to 2003-04)	28 + 02
		UT
2.	Integrated Scheme of Oilseeds, Pulses, Oilpalm and Maize (ISOPOM)- Pulses	14
	(2004-05 - 2006-07).	
	XI th Plan	
1.	Integrated Scheme of Oilseeds, Pulses, Oilpalm and Maize (ISOPOM)- Pulses	14
	(2007-08 - 2009-10)	
2.	National Food Security Mission-Pulses (Rabi, 2007-08 to 2011-12)-Pulses	16
	component of ISOPOM merged with NFSM w.e.f.1.4.2010	
3.	Accelerated Pulses Production Programme (A3P) (2010-11 to 2011-12)	16
4.	Special initiatives for pulses and oilseed in dry land areas under RKVY (2010-11)	07
5.	Integrated Development of 60000 Pulses villages in Rainfed Areas under RKVY	11
	(2011-12)	
6.	Macro Management of Agriculture (MMA) (2004-05 onwards)	Other than
		NFSM
	XII th Plan (2012-13 to 2016-17)	
	2012-13 to 2013-14	
1.	National Food Security Mission (NFSM)óPulses	16
2.	Accelerated Pulses Production Programme (A3P)	16
3.	Special Plan to achieve 19+ million tonnes of Pulses prod. during <i>Kharif</i> 2012-13	08
	2014-15 to 2017-18	
1.	National Food Security Mission (NFSM)óPulses 2014-15	27
2.	National Food Security Mission (NFSM)óPulses 2015-16	27
3.	National Food Security Mission (NFSM)óPulses 2016-17	29
4.	National Food Security Mission (NFSM)óPulses 2017-18	29
5.	Seed Hub-ICAR	150
6.	Breeder Seed Production Programme -ICAR	08
7.	Seed Minikit	NFSM
		States
8.	Cluster FLDs ó Pulses through 549 KVKs	31366 ha
9.	Establishment/strengthening of Bio-fertilizer and Bio-control Production Units	24 Nos.
10.	Farmer Producer Organization (FPOs)	111 Nos.
L	<u> </u>	

(Table-18.2): Recent policy initiatives/interventions taken (2015-16 to 2017-18)

S. No.	Road Map	Year/	Allocation	Remark
	(Interventions/ Initiatives)	Inception	(Rs. in Cr)	
1.	Enhanced allocation-NFSM	2015-16	640.16	> 50% of Total NFSM
	Pulses			Allocation
2.	Additional Allocation	2015-16	440.00	In addition to regular Rabi
	(Spring/Summer Pulses)	2016-17	346.00	Programme
		2017-18	577.67	
3.	Modification in BGREI- Pulses	2015-16	With	To include cluster
	area expansion in Rice Fallows		BGREI	demonstration under Cropping
			allocation	System approach (CSBD) in
				Rice fallows involving pulses.
4.	CFLDs through	2015-16	12.00	Promotion/adoption of new
	ICAR/ATARI/KVK	2016-17	25.29	varieties
		2017-18	26.11	ICAR-IIPR : 0.98 Cr. Since
				2016-17
5.	MSP enhanced substantially	2015-16		(Gram, Lentil) -Marketing
				season 2016-17
6.	Seed availability (Strengthening	_		
a)	Enhancing Breeder Seed	2016-17	20.39	To increase BSP from 10000
	Production (EBSP)	to		qtls. to 14000 qtls. (2016-17),
	(8 states/ 12 locations)	2018-19		15000 qtls. (2017-18) and
				16000 qtls. (2018-19)
b)	Seed Production Incentives	2016-17	200.00	@Rs. 2500 per qtls. on <10
	(To increase SRR/ VRR)	2017-18	111.50	yrs old varieties
c)	Pulse Seed Hubs	2016-17	225.00	150 Nos. @ 1000 qtls. per
,	(24 states/ 150 centres)	to		seed hub annum seed
		2018-19		production target.
d)	Pulse Seed Minikits distribution	2016-17	100.00	For 0.4 ha demonstration
	(To ensure varietal			(Urd, Mung, Arhar-@ 4 kg
	Replacement)			Gram -@ 16 kg; Lentil-@ 8
				kg)
e)	Seed Village Programme	2017-18		(29 states, 638 districts)
				60,000 village, 19.55 lakh
				farmers
7.	Technology demonstration			
a)	Increased Cluster demonstration	2015-16	Within	2015-16 - 5.10 lakh ha
	(StatesøAAP)	2016-17	NFSM Pulse	2016-17 - 5.50 lakh ha
			AAP	(for bridging the yield gaps)
b)	Promotion of minor pulses	2016-17	From within	ICAR to Strengthen BSP and
	(Rajmash, Cowpea, Fieldpea,		EBSP/ Seed	identification of varieties (for
	Horsegram etc.)		Hub and	exploiting minor pulses)
			CFLDs	(for bridging the yield gaps)
c)	CDP (Tur on Rice bunds)	2016-17	From within	Area Expansion
	(Target- 02 Lha)		NFSM/	
			BGREI	

S. No.	Road Map	Year/	Allocation	Remark
	(Interventions/ Initiatives)	Inception	(Rs. in Cr)	
d)	CDP (demonstrations on Ridge	2016-17	From within	Popularization of Good
	and Furrow cultivation,		NFSM/	Agricultural practices (GAP)
	summer mung, Tur		BGREI	
	transplanting & intercropping)			
e)	Critical Irrigation (Provision of	2016-17	Within total	Sprinkler sets and water
	irrigation under PMKSY,		allocation of	carrying pipes (to Improve
	50% allocation reserved for		PMKSY	Water use efficiency (WUE)
	pulses)			
8.	Remunerative Prices to pulse g			
a)	Loan against warehouse	2016-17	Within Interest	Interest free loans against
	receipts		Subvention	warehouse receipts
	(Pledge Loan)		Scheme	
b)	MSP and Procurement (Credit	2016-17	NAFED,	45.57 Lakh MT
	guarantee raised from Rs. 9000		SFAC and FCI	Rs. 10.57 Cr
	Cr to Rs. 19000 Cr)			
9.	Extension			
a)	Advisory on pulses (monthly)	2016-17		DPD, Bhopal ó IYOP 2016
				celebrated with training
				pamphlets on all pulses in
				Hindi/English
b)	Creation of IT based			IYOP 2016 round the year
	information portal			trainings/ workshops were
	(http//dpd.gov.in)			organized.
				Pulse Bulletins/ pamphlets
				(bilingual) developed and
				distributed across the country.
	Effective monitoring	2015-16	Within NFSM	• Senior Officers Meeting
	Č	onwards		(SOM) under the
				chairmanship of Secretary,
				DAC&FW- weekly.
				• Committee for monitoring
				actions of road map- under
				the chairmanship of CEO,
				NRAA, Govt. of India-
				fortnightly & monthly.
				 Video Conferencing by
				DAC&FW- weekly.
				 CDDs Review Meeting
				under the chairmanship of
				JS (Crops) - bimonthly.
				(F)

S. No.	Road Map	Year/	Allocation	Remark
	(Interventions/ Initiatives)	Inception	(Rs. in Cr)	
10.	Effective monitoring	2015-16 onwards	Within NFSM	 Monthly field visits by CDDs & consultants. Field visits/ monitoring by Director, ATARI; DES (SAUs), Director, IIPR. ICAR-AGM on pulses/ interface/ seminar/ workshop/ meetings by Directorate of Pulses Development, Bhopal. Monitoring of seed minikits distribution and FLD/ CFLD by CDDs. Monitoring of Seed Hubs,
11.	Soil test based promotion of INM/ IPM	2015-16	Within NFSM- Pulses	EBSP by CDDs. 2015-16 - 29 Lha 2016-17 - 41 Lha
12.	Targeting Rice Fallow Area (TRFA) (Area Expansion)	2016-17	8.85 (2016-17) 203.64 (2017-18)	06 eastern states of Assam,
13.	FPOs ó Value Addition Chain Development (Marketing/ Value Addition)		52.10	11 States (119 FPOs on pulses)
14.	Enhanced MSP of Pulses	2015-16 to 2017-18	Kharif 2018-19 MSP Arhar-Rs.5675 Urd-Rs.5600 Mung-Rs.6975	The real term exponential growth rates >4% in all five pulses
15.	EXIM policy	2017-18	Import of pulses during 2017-18 declined by 03 MT resulting saving of foreign exchange of Rs. 7698 Cr.	Government has imposed import duties on pulses for the first time in this decade. All varieties of pulses, including organic pulses, have been made :freeøfor export.
16.	District Agriculture Contingency Plan (DACPs) involving pulses	2016-17	-	By CRIDA involving 46 SAUs and 8 ICAR institutions. 676 districts.

(Table-18.3): Interventions under NFSM-Pulses

Sr.	Head		Interventions
No.			
1.	Technology Demonstrations	•	Cluster demonstrations
		•	Cropping system based demonstrations
		•	Front Line Demonstrations by ICAR/SAUs
2.	Seed	•	Distribution of HYVs seed
3.	Integrated Nutrient Management (INM)	•	Micro-nutrients
		•	Lime/Gypsum/80% WG Sulphur
		•	Lime
		•	Bio-fertilizers
4.	Integrated Pest Management (IPM)	•	Distribution of Plant Protection chemicals
		•	Weedicides
5.	Resource Conservation	•	Power Knap Sack Sprayers
	Technologies/Tools	•	Manual Sprayer
		•	Zero Till Seed Drills
		•	Multi Crop Planter
		•	Seed Drills
		•	Zero Till Multi Crop Planters
		•	Ridge Furrow Planters
		•	Rotavators
		•	Chiseller
		•	Laser Land Levelers
		•	Tractor mounted sprayer
		•	Multicrop Thresher
6.	Efficient Water Application Tools	•	Sprinkler Sets
		•	Pump Sets
		•	Pipe for carrying water from source to the field.
		•	Mobile Rain guns
7.	Cropping System based trainings	•	Four Sessions in a crop season (One before
			Kharif and Rabi Season & one each during Kharif
			and Rabi Crops).
8.	Miscellaneous Expenses (Project		Project Management Team & other miscellaneous
	Management Support & Monitoring)		expenses at District and state level
9.	Local Initiatives		On project basis, up to 9% of the total allocation
			to the state
10.	Other		Specialized projects for high productivity areas
			Support to institute/organizations including
			NGOs in remote areas.
			Value chain integration of small producers
			Assistance to Custom Hiring Centres
		•	Marketing support for pulses

(Table-18.4): Summary of research project funded under of NFSM-Pulses in year 2017-18

(Rs. in lakh)

		T	T			T	. et		S. in takn)
S. No.	Project Title	Implementing Agency	Project Duration	Total Allocation	Allocation for 2017-18	Unspent balance/ Revalid .2016-17	1st Release	2 nd Release	Location/ Varieties
1	Enhancing breeder seed production for increasing indigenous production of pulses in India (8 states/ 12 locations)	IIPR, Kanpur	2016-17 to 2018-19	2039.00			407.80 (20 % of allocation)		Rel-2016-17 Rs. 815.60
2	Creation of seed óhubs for increasing indigenous production of pulses in India (24 states/150 centres)	IIPR, Kanpur	2016-17 to 2017-18	22531.08 (150 hubs)	11164.26	768.18	4895.50		
3	Generation advancement and development of new genotypes through pre-breeding in Lentil and Kabuli Chickpea"	ICARDA	2013-14 to 2016-17	320.196 (Revised)		4.89			
4	Enzymatic pre treatment in the processing of Pigeonpea	JAU,Junagarh (Gujarat)	2014-15 to 2016-17			0.21			Junagarh / Var. BDN-2
5	Enhancing productivity through introduction of new high yielding varieties, production technologies in chickpea, green gram, black gram & cowpea.	UAS, Dharwad , Karnataka	2016-17		11.02				
6	Enhancing mothbean and mungbean productivity through high yielding varieties, nutrient management and IPM practices in Western Rajasthan	SKRAU, Bikaner	2014-15 to 2016-17			1.83623			
7	Development of suitable technology for increasing the production of pulses in rice fallows	OUAT, Bhubaneswar	2014-15 to 2016-17	80.37		1.24532			Greengram, Blackgram and Gram
8	Scaling up and popularization of high yielding pigeonpea hybrids for enchancing productivity of small and marginal farmers of	ICRISAT, Hyderabad, Telangana	2016-17	77.965				19.49 (committed liability of	ICPH 2740
	Maharashtra, Karnataka & Odisha States of India	ICRISAT, Hyderabad, Telangana	2018-19	649.685				2016-17)	Approved for 2018-19 ICPH 2740 & ICPH3762
9	Addressing phytophthora blight disease : An emerging threat of pigeonpea expansion and production	ICRISAT, Hyderabad	2013-14 to 2016-17	400.923 (Revised)		3.33 (committed expen. of 2016- 17)	80.0		
10	Identification of salt tolerant chickpea varieties for coastal regions of Gujarat.	NAU, Navsari (Dr. P.B. Patel)	2014-15 to 2016-17	32.123 (Revised)		2.11788			

18.2 Performance over-view: (First to Twelfth Plan)

An analysis to pre, post TMOP and during NFSM interventions

For comparative analysis, the average area covered, the production, productivity and percentage of pulse area under irrigation have been taken into consideration on Five Year Plan basis. Annual plans (1966-69, 1979-80 and 1990-92), during which the five year plans could not be enforced/implemented, have, however, been excluded for the purpose of analysis. The analytical review of pulses status prior to TMOP, during the TMOP and during NFSM period is briefly analyzed as below:

(Table- 18.5): Plan-wise trend of growth in pulses

{Area-Mha., Production- MTons, Yield- kg/ha}

Plan	Average Area	Average Production	Average yield	Average % of irrigation coverage
Pre-TMOP Periods				
I st Plan (1951-56)	21.09	10.04	475.2	9
II nd Plan (1956-61)	23.71	11.75	494.8	8
II rd Plan (1961-66)	23.85	11.14	466.8	9
IV th Plan (1969-74)	22.21	10.90	491.4	9
V th Plan (1974-79)	23.32	11.71	501.4	8
VI th Plan (1980-85)	23.08	11.77	509.8	8
VII th Plan (1985-90)	23.08	12.54	543.0	9
Post-TMOP Period (inclu	udes two ann	ual plans (1990	-92)	
VIII th Plan (1992-97)	22.47	13.34	593.6	12
IX th Plan (1997-02)	21.97	13.15	597.4	13
X th Plan (2002-07)	22.44	13.35	593.8	14
NFSM Plan Period				
XI th Plan (2007-2012)	23.97	15.85	662	16
XII th Plan (2012-2017)	25.28	18.84	745	19

18.2.1 Area Expansion

- During the first five year plan (1951-56), the average pulse acreage of 21 million hectares maintained an increasing trend till Third plan (1961-66) where an area of about 24 million ha was occupied. However, there was a slight drop in area coverage i.e. 22.21 million hectares during the Fourth plan (1969-74) despite the introduction of first centrally sponsored Pulses Development Scheme. It is also a fact that the normal average area of pulses enhanced to about three million ha during IInd five year plans, the periods when average per cent coverage under pulses was about 8-9 percent.
- It is observed that the role of plan funds had catalytic role especially in stabilization of area coverage under pulses as beyond the IIIrd five year plan, the normal five year plan area has been between 22-23million hectares, a visible two million hectares increase over the Ist plan period.
- Another most important observation is stability in pulse area from eighth plan(1992-97) period to tenth plan period (2002-07) and significantly increased eleventh to twelfth plan period (2007-12 to 2012-2016). The plan period had the critical intervention in pulses sector through the Technology Mission (TMOP) and National Food Security Mission (NFSM) with the increase in irrigation coverage, 16% and 19 % of total pulses stablized in irrigated area.

18.2.2 Production Enhancement

- During the initial phase of the *IstFive year plan (1951-56)*, the production of pulses was 10 million tonnes. There was a slight fall during the *IVth Plan (1969-74)* from the *IIIrd plan* recording the average production of 10.90 Million tonnes. However, there was a homogeneous increase thereafter. With the inclusion of pulses development under TMOP during August 1990, the beneficial impacts were realized during the *VIIIth Plan (1992-97)* and *Xth Plan (2002-07)* the country witnessed an average plan period production of 13.34 Million tones and 13.35 million tonesrespectively, the maximum ever achieved during the pre TMOP Five year plans periods.
- **During** X^{th} **plan** (2002-07), inspite of the consecutive droughts/flood in the major pulses growing states of Madhya Pradesh, Rajasthan, Uttar Pradesh, Bihar Andhra Pradesh and Maharashtra and stagnant area coverage, the country harnessed an average production of 13.35 lakh tonnes which may be attributed to TMOPs critical intervention and Central funding support under **NPDP/ISOPOM** making a dent on seeds/irrigation and other infrastructural support to farmers.
- During the course of implementation of *NFSM XIIth plan (2013-14& 2016-17)*, the country witnessed a significant increase in production of pulses *i.e.* 19.25 *Million tons* and 22.95 *Million tons* respectively, the maximum ever achieved during 2016-17.

18.2.3 Productivity

- Productivity of pulses has also increased during the TMOP period. Pre-TMOP plan period average yield during the first plan (1951-56) was only 475 Kg/ha and the Third Plan (1961-66) even exhibited minimum productivity of about 467 kg/ha whereas, maximum average yield was recorded (598 Kg/ha) during the Nineth Plan (1997-2002), Approximately 131 Kg/ha increase in productivity levels between the Pre-TMOP (1961-66) and during the TMOP period recorded.
- Similarly, during the *NFSM plan period*, XIth and XIIth plan, productivity was achieved 662 kg/ha and 744 kg/ha respectively. Although this productivity is still below the world¢s average productivity of 909 kg/ha and as also what has been realized under the frontline demonstrations of ICAR. A productivity gap of 56% under total pulses between the FLDs and State average yield is the existing potential and a challenge for both the research and development agencies to harness.

18.2.4 Irrigation

- The production and productivity increase during TMOP period against the stagnant area coverage under pulses may be attributed to adoption of modern technology based package of practices, more coverage of area under irrigation including various inputs. These could be possible because of the launching of TMOP in 350 districts of 30 states/Union Territories, nation-wide. It can be assumed that if the pace of pulses production in the country is constantly maintained, the pulse requirement could be easily met in the long run.
- In-adequate irrigation facilities, especially the supply of critical irrigation, are the main cause of low production of Indian pulses. Taking the average of the five years of the

first Five Year Plan (1951-56), the coverage of area under irrigation was hardly 9.18%. There was a decreasing trend till the Fifth plan. However, the increasing trend was restored from the Sixth plan with the maximum coverage of 12-13% from eighth plan onward i.e. the initial phase of the launching of TMOP, attributing the productivity enhancement i.e. about $600 \text{ kg/ha}(VIII^{th}-X^{th} \text{ plan})$.

• During the NFSM plan period, irrigation increased upto 19%, attributing the productivity enhancement from 594 kg/ha during Xth Plan (Before NFSM) to 745 kg/ha during XIIth Plan.

18.3 Strategies Adopted

18.3.1 Area expansion

Rabi pulses to bring additional production from additional area coverage in rice fallows, crop-wise strategies were adopted viz., *Gram* - CG, West Bengal, Bihar, Jharkhand, Odisha, Assam, Andhra Pradesh, Tamil Nadu; *Lentil* - Chhattisgarh, West Bengal, Bihar, Jharkhand, Assam and *Mung/Urd* in rice fallow costal region; *Intercropping* - Gram with barley, *Mustard and Linseed* in Rajasthan, UP, Bihar, Vidarbha (Maharashtra); *Intercropping* - Gram/Lentil with autumn planted ratoon sugarcane in UP, Maharashtra, Bihar.

Kharif pulses - additional production from additional area coverage (diversion to other crops like cotton, oilseeds, coarse cereals), cultivation of kharif pulses as intercrop, planting of red gram on rice bunds, cultivation of minor pulses in niche areas.

Spring/Summer pulses Punjab, Haryana, Madhya Pradesh, Uttar Pradesh, Bihar, West Bengal, Gujarat, Jharkhand, TN and AP were chosen.

During 2017-18 total allocation under additional pulses production programme of Rs. 577.67 Crores (350 Crore GoI + 227.67 State Share) is operational in 17 states.

(Table-18.6): Area coverage under spring/summer pulses

(Area-Lakh ha)

States	2014	2015	2016	2017	2018
Tamil Nadu	1.83	1.73	2.38	2.31	2.39
Bihar	0.00	2.58	0.73	2.95	1.47
Uttar Pradesh	1.16	2.04	1.56	1.36	1.40
Andhra Pradesh	1.53	0.44	0.26	0.74	0.15
Gujarat	0.56	0.51	0.43	0.40	0.38
Haryana	0.44	0.17	0.06	0.06	0.06
Karnataka	0.13	0.15	0.19	0.09	0.09
Madhya Pradesh	2.24	1.69	1.67	1.51	0.92
Punjab	0.43	0.65	0.32	0.14	0.15
West Bengal	0.40	0.39	0.52	0.61	0.86
Others	0.16	0.32	0.21	2.35	1.08
Total	8.88	10.67	8.33	12.52	8.95

Source: WWWR Report, DPD, Bhopal




Fig.-18.1: Area Coverage under Total Summer Pulses




Fig.18.2: Major state-wise Area Coverage under Spring/Summer Pulses

18.3.2 Targeted rice fallow area (TRFA)

- Under area expansion of rabi pulses in rice fallows, TRFA programme has been initiated since 2016-17 in 06 Eastern States of Assam, Bihar, Chhattisgarh, Jharkhand, Odisha and West Bengal. Against a target of 4.5 Mha rice fallow under pulses, an area of > 2.6 Mha has so far been achieved uptill 2017-18.
- During 2016-17 the TRFA was implemented in 15 districts of 6 states to cover 19.14 lakh ha (pulses-15.31 lakh ha + oilseed-3.83 lakh ha). During 2017-18 the scheme has been extended to 40 districts and 4000 villages with a view to cover 15.00 lakh ha (pulses-12.00 lakh ha + oilseeds-3.0 lakh ha) with support for cluster demonstrations, minikit distribution and training to the farmers etc. According to the report an area of 10.72 lakh ha were covered (pulses-9.13 lakh ha + oilseeds-1.60 lakh ha) in rice fallows covering 43 districts and 3739 villages. The additional area coverage resulted a production of 9.04 lakh tonnes of pulses and oilseeds as against the production target of 10.00 lakh tonnes. Most of the rice fallows were covered under pulses *viz.* pea, lentil, black gram, green gram, chickpea, arhar and lathyrus.

- The total allocation under TRFA-Pulses for the year 2017-18 was Rs. 98.27 crores which was fully released to the states. However the expenditure reported so far is Rs. 62.33 crores which is 63% of the release.
- During 2018-19 the total allocation 220.80 Crore out of this Rs. 148.28 Crores of Central share comprising 133.48 Crore for demo. & Rs. 14.80 Crore for Minikit.

(Table-18.7): Progress of TRFA

(Area- Lha, Allo Amount- Crores)

State	Area	Pulses C	Coverage	2018-19	A	rea Cov	ered	Allocati	on (CS)	Intervent
	under	No. of	Districts/	(Target)	2017	'-18	2018-19	2017-18	2018-19	ions
	Rice	Villages	(2017-18)				(Target)			
	Fallow	Target	Ach.		Target	Ach.				
Assam	10.42	8/800	8/800	8/800	1.84	0.92	2.10	21.48	28.80	Cluster
Bihar	3.00	5/380	5/380	7/700	0.24	0.75	0.35	12.23	16.80	Demo.
Chhattisgarh	28.56	5/427	5/427	9/900	2.80	4.49	3.95	12.23	21.60	Minikit
Jharkhand	4.75	4/378	4/378	5/500	0.24	0.18	0.35	8.37	12.00	Distributi
Odisha	29.61	9/854	9/854	11/1100	3.44	2.05	4.20	21.98	26.40	on, and
West Bengal	12.00	12/900	12/900	10/1000	3.44	0.75	3.85	21.96	24.00	Farmers
Total	88.34	43/3739	43/3739	50/5000	12.00	9.14	14.80	98.25	129.60	Training

(Table-18.8): State-wise production of pulses under TRFA during 2017-18

(Production in lakh tonnes and Yield kg/ha)

States	Area	Production	Yield	<u> </u>
Assam	0.92	0.82	897	
Bihar	0.75	0.71	940	
Chhattisgarh	4.49	4.48	997	
Jharkhand	0.18	0.19	1079	
Odisha	2.05	1.07	524	
West Bengal	0.75	0.65	867	
Total	9.14	7.92	867	

- Short duration high yielding rice varieties with its earlier planting as dry seeding/ DSR and early transplanting were introduced under NFSM to bring pulses under Rice fallow.
- Rice fallow covered under additional pulse programme by Relay sowing (uttera cropping) of lentil, khesari, small seeded chickpea and pea also solved the problem of late sowing.

Recommendations for improving pulses productivity under rice fallow

Under mentioned recommendations and policy issues indicated in the NAAS Policy Paper (64) on "improving productivity of rice fallows", is relevant to harness the potential of rice fallows, and may be the part of ongoing programmes of pulses development under NFSM- Additional Pulse Programme and RKVY-TRFA.

- Mechanization of field operations: Residual soil moisture in surface layer at the time of planting rabi crops is the major constraint in rice fallows. Relay cropping in standing rice is often practiced but with use of combine for rice harvesting, the option is now shifting for direct seeding using zero-till drill or turbo type Happy Seed drill which need to be designed for different situations. For harvesting and threshing, appropriate machines need to be designed and developed.
- ii) Scaling-up crop management practices: Tillage and plant population management, application of nutrients and weed management in rabi crops pose serious challenges in rice fallows. Early-maturing crop varieties, relay cropping, higher seed rate, seed priming, seed inoculation with Rhizobium culture, seed pelleting, mulching, foliar spray of nutrients etc. are recommended practices which need to be further refined and standardized for different ecosystems. Work on development of short-duration, highyielding varieties, appropriate seeding techniques, water harvesting and recycling, post-emergence herbicides, biotic and abiotic stresses etc. need to be strengthened.
- iii) Crop-specific information on area expansion: Based on biophysical conditions, farm resources and market demand, likely coverage of area under each crop in different states/ region need to be estimated. This would facilitate area expansion in phased manner by arranging critical inputs.
- iv) Periodic GIS mapping: In order to monitor impact of R&D efforts on area expansion in rice fallows under different crops, cropping systems and soil health, periodic monitoring through GIS is required.
- v) Creation of community water reservoirs: Despite heavy rains during *kharif* season, soil moisture becomes the most critical limiting factor for raising second crop during winter as most of the runoff is wasted. It is, therefore, necessary to create farm pond and community water reservoirs in the area well supported by Government. This will serve as important source for life-saving and supplemental irrigation. Further, the loss of soil and plant nutrients from productive lands will be reduced.
- vi) Quality seeds: Timely availability of quality seeds is often a major constraint for delayed planting and poor yields. Hence, community-based seed production programmes need to be launched with appropriate processing and storage facilities. The national and state seed Corporations should strengthen their activities in these areas.
- vii) Ensuring timely availability of other critical inputs: Traditionally, the winter crops on residual soil moisture are grown using local varieties without application of plant nutrients, bio-fertilizers, fungicides and other agro-chemicals due to their nonavailability. Since crop productivity is the driver for area expansion, which in turn is influenced by better crop management, emphasis needs to be placed on timely availability of all critical inputs.
- viii) Marketing infrastructure: Marketing plays a key role in enthusing farmers for crop production. Well organized marketing and processing of farm produce need attention.
- ix) Protection from stray cattle: Blue bull and other stray cattle cause heavy damage to pulses and thus discourage farmers to grow winter crops. Appropriate policies are

needed to tackle this menace. To avoid crop damage by stray cattle, open grazing lands at panchayat level should be earmarked. These activities should be the part of state level planning.

• Recommendations based on performance/experience

Under NFSM-pulses, additional area coverage programme of spring/summer season and RKVY-TRFA, the development efforts on increasing productivity of pulses in rice fallow areas includes identification of suitable varieties, planting methods, foliar nutrition and plant protection, refining and packaging improved technologies etc are based on the past experience, to address different problems.

- i) Selection of crops and varieties: should be decided on the basis of winter temperature, soil texture, soil moisture content etc. (In lentil and gram, small seeded varieties due to better contact with soil, less rotting be selected).
- ii) **Seed priming and optimum seed rate:** Overnight soaking of seeds (seed priming), hastens seed germination and crop establishment under relay cropping. Adoption of 20-25% higher seed rate over the recommended rate is recommended ensures desired plant stand.
- iii) Foliar nutrition: Since application of fertilizers under relay cropping is not feasible, seed pelleting and foliar application of nutrients should be practiced. Foliar application of 2% urea at flowering and pod formation significantly improves yields of chickpea under rainfed conditions by increasing leaf N content and making them photo synthetically more active. Seed pelleting with micronutrients like Zn and Mo is also recommended as a part of nutrient management strategy in rice fallows.
- iv) **Planting strategy:** In rice fallows, planting is generally delayed. Under relay planting, seeds should be broadcast 2-5 days before harvest of rice. Zero-till seed-cum-fertilizer drill should be used wherever feasible when planting is done after harvest of rice. It is necessary to use short to medium maturing varieties of rice for timely planting of *rabi* crops.
- v) Plant protection: Since post-emergence herbicides are not commercially available specially for crops like chickpea and lentil and inter-cultivation is difficult due to hard soil, hand pulling of weeds is the only option which should be done at an early stage. Post-emergence herbicide (Imazethapyr @ 50 g/ha) has been found quite effective against seasonal grassy weeds in crops like groundnut, urdbean and mungbean. It should be applied at 3-4 leaf stage. Similarly, quizalofop can be used to check ratooning of rice stubbles which cause substantial moisture loss. Insect-pests and diseases should be promptly controlled. Seed dressing with fungicides like carbendazim should be done.
- vi) Issues based major technological interventions and region specific varieties are summarized under *Table 18.9 (a) and (b)*.

(Table – 18.9 - a): Major technological interventions

Issues	Interventions	Action
Lack of suitable	Development of high-yielding varieties	ICAR-IIPR
cultivars	with appropriate maturity duration	
Poor crop stand and	Tillage machines, sowing methods, seed	SDA/SAUs
establishment	priming, higher seed rate, timely planting, seed treatment with fungicides	
Diseases and pests	Development of IPM modules	SDA/SAUs/NCIPM
Weed menace	Post-emergence herbicides like	SDA/SAUs/DWR
weed menace	Quizalophop ethyl and Imazethapyr	
Nutrient management	Foliar spray of urea/DAP to supplement N and P	SDA/SAUs
Micronutrient		SDA/IISS
deficiencies	Mo, B, Zn as seed pallets	~
Terminal	Residue mulching	SDA/SAUs/CRIDA
moisture/heat stress		
Non-availability of	Informal and formal seed production and	SDA/SSC/NSC
quality seeds	supply systems	
Lack of	Tillage machines, zero-till planter and	SDA/SAUs/CIAE
mechanization	harvester	
Poor transfer of	Innovative farmerøs participatory	SDA/SAUs/KVKs
technology	approach	

(Table -18.9-b) Performance based recommendation

Region Rec. Crops & Varieties	Eastern Plains	Central region	Coastal Region
Lentil	Rust- A major threat Small seeded lentil varieties having resistance to rust WBL-77, KLS- 218, PL-8, NM-1, DPL-15	 Chickpea var Pusa-372, PG- 186, Udai Small seeded chickpea var. JSC-55, JSC-56, JG-14, vijay, JG- 315, JAKI-9516 Lathyrus- var- Ratan, Prateek, Mahateora 	Powdery Mildew a major threat 1. Urdbean-var powdery mildew resistance var., LBG-17, LBG-602, LBG-623 Urdbean Normal planting var. (Mid Nov to Mid Dec) LBG-402, LBG-611, LBG-22, LBG-648, LBG-685, LBG-645, LBG-709, LBG-752 2. Late Planting Urdbean (II nd fortnight ofDec) var. LBG-22, LBG-645, LBG-709, LBG-752 3. Mungbean varieties (Normal Planting) LGG-460, LGG-410, LGG-450, LGG-407, IM-96-3, Pusa 9072, NARM-1,2 and 18.

Source: Policy Paper 64: Improving Productivity of Rice Fallows

18.3.3 Productivity Enhancement

• Promotion of Sulphur & Zinc

Wide spread deficiency of sulphur and zinc noticed in pulse growing regions constraints the productivity of pulses. In major pulse growing areas, 44 districts have shown 40-60% sulphur deficiency and 82 districts with 50-60% zinc deficiency. In view of encouraging response to application of S and Zn with cost benefit ratio of 10-21%, their application was vigorously pursued after 2014-15.

• Promotion of Rhizobium & PSB

About 40% pulse growing regions have low to medium population of native Rhizobium. Seed inoculation with bio-fertilizer (Rhizobium and PSB); low cost inputs; are known to increase pulse productivity by 10-12%. Rhizobium + PSB provided under NFSM.

• Technology demonstration conducted

The frontline demonstrations were conducted in different agro-climatic regions on important pulse crops with a view to demonstrate and assess the benefits of new varieties and technologies under diverse cropping systems have revealed the existing potential of productivity to be exploited through technological interventions.

A package technology like improved cultivar, Rhizobium inoculation, use of Sulphur, INM, application of weedicide, foliar spray of urea, IPM etc were vigorously pursued.

For good crop establishment, seed priming (soaking the seeds over night in water surface, drying and sowing next day), seed treatment with effective Rhizobium strain, sowing of seed into deeper moist soil (in case of chickpea), lime pelleting for acidic soil and gypsum in saline areas was encouraged under NFSM pulses/CFLDs.

Government focused on key areas like seeds of improved varieties, irrigation tailored to pulses (especially micro-irrigation), bringing new niche areas under pulse cultivation, attractive minimum support price (MSP) and market that allow farmers to increase their profitability aligned to improved farmer welfare.

The Government of India released amount of Rs. 97.50 lakh for 1300 no. of demonstration during 2017-18 and Rs. 117.00 Lakh for same nos. of demonstration during 2018-19.

(Table – 18.10): Promotion of improved varieties

States	Crop	Leading varieties						
Andhra	Tur	LRG-41						
Pradesh	Gram	JG-11-KAK-2						
	Mung	LGG-460, Urd ó PU-31, LBG-752						
Bihar	Mung	HUM-16, IPM-02-03, Panth Mung ó 5						
	Kulthi	DV-7						
Jharkhand	Tur	NDA-2, MAL-13, ICPH-2671,						
	Gram	Kabulichana ó 2, JAKI-9218						
	Mung	TM-99-21, HUM-12 (MalviyaJanchetna), HUM-16, HUM-668,						
		IPM-02-03						
	Urd	PU-31, Sujata, WBU-109, Sekhar-3,						
	Kulthi	Birsa-Kulthi-1, Madhu,						
	Lentil	WBL-77, HUL-57						
Karnataka	Tur	BRG-2, TS-3R, BSMR-736						
	Gram	JG-11, JAKI-9218						
	Mung	BGS-9, SML-668						
	Urd	T-9, TAU-1						
Maharashtra	Tur	PKV-Tara,BDN-711,BDN-708, BSMR-853, ICPL-8863 (Maruti),						
		ICPL-87119 (Asha), ICPL-87 (Pragati						
	Gram	Akash, Digvijay, JAKI-9218, Virat, ICCC-37,KAK-2,Vishal, vijay						
	Mung	BM-2002-1, PKV-AKM-4, BM-2003-2, Utkarsh, Kopargaon						
	Urd	AKU-15, TAU-1						
Odisha	Tur	Asha, Lakshmi, UPAS-120, LRG-41						
	Mung	PDM-139 (Samrat), HUM-12 (MalviyaJanchetna), SML-668, IPM-						
		02-3, IPM-2-14						
	Urd	PU-30,31, Shekhar-2, IPU-02-43, Azad Urd						
Rajasthan	Gram	RSG-44, GNG-1581 (Gangaur), GNG-1958 (Marudhar), CSJ-515,						
		GNG-2144 (Teej), RSG-974						
	Mung	SML-668, GM-4, MH-421, IPM-02-3,IPM 2-14, HUM-16, HUM-						
		12 (Malviya Jan Chetna), PDM-139, MH-2-15 (Satya), MH 2-14						
	Urd	PU-30,31, IPU-94-1, TAU-1,2, KU-300, KPU-07-08, KPU-405, K-						
		96-3						
	Moth	RMO-40,225,423, CAZRI-2, RMB-2						
Tamil nadu	Arhar	LRG-41, CO (Rg)-7, BRG-1,2; Urd- VBN-5,6, ADT-3						
	Mung	CO (GG) 912/CO-7, CO-8						
	Kulthi	Paiyur-2						
Telangana	Arhar	LRG-41, PRG-176, ICPL-87119 (Asha)						
	Gram	JG-11, JAKI-9218						
	Mung	LGG-460, MGG-295, IPM-2-14						
	Urd	LBG-752, T-9, IPU-2-43, PU-31						
Uttar Pradesh	Arhar	UPAS-120,PUSA-992,Narendra Arhar-1,PUSA-9,PDA-11, Bahar						
	Gram	JAKI-9218, Shubhra, Ujjawala, DCP-92-3, Avrodhi, Uday						
	Mung	IPM-2-3, Narendra Mung-1, MalviyaJyoti (HUM-1),						
	<u></u>	MalviyaJanchetna (HUM-12), PDM-139 (Samrat).						
	Urd	IPU 91-1 (Uttara), IPU-2-14, PU-31, PU-35, Shekhar-3, IPU-2-43,						
		T-9						
	Lentil	PL-4,5, DPL-62, Narendra Masoor-1, K-75, L-4076						

Source: SDAs of Agriculture.

18.3.4 Marketing Strategy

Enhanced procurement of pulses: (2014-15 to 2017-18)

NAFED has done record procurement of pulses jointly with SFAC and FCI during the year 2017-18 under Price Support Scheme (PSS) and Price Stabilization Funds (PSF), funded by Ministry of Agri. & FW. It procured more than 2008.52 thousand MT of Pulses (Gram 188.59 thousand MT, Masoor 27.07 thousand MT, Moong 407.74 thousand MT, Urd 290.62 thousand MT and Tur 1094.49 thousand MT).

NAFED has been involved in creating buffer stocks and stabilizing the prices of Pulses. A substantial quantity from buffer stock has been proposed to be supplied to Para-military and Defense forces. Also the supplies will be made to state governments as per their requirements under PDS and other such schemes. Consequently, the procurement agency has positioned itself as the Pulse Armøof the government.

The total monetary investment towards procurement of total pulses during 2017-18 is Rs. 10.57 Cr by alone NAFED. Crop-wise and state-wise procurement of total pulses is given in (*Table 18.11 to 18.12*)

(Table-18.11): Crop-wise procurement of pulses enhanced

(Oty. in 000 MT, Rs. in Crore)

Crop	2014-15		2015-16		2016-17		2017-18	
	Qty.	Amt.	Qty.	Amt.	Qty.	Amt.	Qty.	Amt.
Arhar/Tur	6.30		92.85	0.40	1567.37	4.68	1094.49	5.79
Urd			33.07	0.05	183.19	0.56	290.62	1.56
Mung					430.41	1.08	407.74	2.23
Gram	775.13	1.12			287.67	0.36	188.59	0.88
Lentil					80.02	0.35	27.07	0.11
Total	781.43	1.12	125.92	0.45	2548.66	7.03	2008.52	10.57

Source: NAFED, New Delhi.




Fig.-18.3: Quantity Procured under MSP




Fig.-18.4: Values of Procured Pulses under MSP

(Table-18.12): State-wise procurement of pulses under MSP (PSS)

State		ed (000 Million nnes)	Value (Rs. Crore)			
State	2001-02 to 2013-14	2014-15 to 2018-19	2001-02 to 2013-14	2014-15 to 2018-19		
Andhra Pradesh	49.05	213.00	172.22	970.29		
Gujarat	29.70	270.95	41.99	1385.71		
Karnataka	28.77	791.15	68.48	4043.19		
Madhya Pradesh	260.63	953.26	377.38	4322.05		
Maharashtra	104.68	956.72	334.25	4962.82		
Rajasthan	92.87	967.48	153.48	4648.05		
Telangana	0.00	303.35	0.00	1549.79		
Uttar Pradesh	71.49	55.10	148.06	294.13		
West Bengal	7.66	6.79	16.94	36.67		
All India	5057.86	3236.57	1171.75	16360.42		

Source: NAFED, New Delhi.




Fig.-18.5: State-wise Procurement of Pulses under MSP (PSS)

(Table-18.13): Crop-wise procurement of pulses under MSP (PSF)

(Quantity- 000 Million tonnes)

Crop	2015-16	2016-17	2017-18	Total
Arhar/Tur	45.53	1165.90	0.00	1211.43
Urd	4.89	88.49	0.00	93.38
Mung	0.00	209.93	0.00	209.93
Gram	0.00	0.00	60.25	60.25
Lentil	0.00	0.00	27.07	27.07
Total	50.42	1464.33	87.33	1602.08

Source: NFSM, Cell, Min. of Agri. & FW (DAC&FW), New Delhi.




Fig.-18.6:Crop-wise procurement of pulses under MSP (PSF)

18.3.5 Farmer Producer Organization: Empowerment Through Group

In its resolve to provident to end to end solution to the various issues relating to pulse sector, the govt. has provisioned to formulate farmer interest groups (FIGs) and FPOs. As of now against the total 847 FPOs, 119 in major 07 pulse producing states (Gujarat, Karnataka, Madhya Pradesh, Maharashtra, Telangana, and Uttar Pradesh) are exclusively involved in the pulses sector.

The total registered FPOs in India about 847. Out of 190 FPOs under taken activities of pulse, total targeted farmers are > 8.0 lakh. About 90% farmers are mobilized. The statewise detail is as under:

(Table-18.14): State-wise progress of FPO promotion (As on 30.06.2018)

S.	State	1	No. of Farm	ers		No. of FPOs		No. of
No.		Mobiliz	Under	Total	Registe	Under the	Total	FPOs on
		ed	Mobiliz	Targeted	red	Progress of		Pulses
			ation	Farmer		registration		
1	Andhra Pradesh	6792	208	7000	7	0	7	-
2	Arunachal Pradesh	1750	1000	2750	2	2	4	-
3	Assam	5647	1853	7500	12	3	15	-
4	Bihar	25685	8315	34000	24	11	35	-
5	Chhattisgarh	29135	0	29000	26	2	28	-
6	Delhi	3535	0	3500	4	0	4	-
7	Goa	1810	0	1750	2	0	2	-
8	Gujarat	19166	834	20000	20	1	21	7
9	Haryana	13240	510	13750	23	4	27	-
10	Himachal Pradesh	4887	0	4850	5	0	5	-
11	Jammu & Kashmir							
	Jammu (Division)	3694	287	3981	1	2	3	-
	Srinagar (Division)	3120	960	4080	1	3	4	-
12	Jharkhand	10009	0	12000	8	2	10	-
13	Karnataka	118218	4282	122500	118	2	120	14
14	Madhya Pradesh	126584	18416	145000	135	9	144	28
15	Maharashtra	88348	13152	101500	85	17	102	24
16	Manipur	5671	1279	6950	4	4	8	-
17	Meghalaya	2990	760	3750	3	1	4	-
18	Mizorun	1700	1000	2700	1	2	3	-
19	Nagaland	1750	0	1750	2	0	2	-
20	Odisha	39463	0	38900	41	0	41	-
21	Punjab	6288	0	6000	7	0	7	-
22	Rajasthan	49617	883	50500	40	2	42	16
23	Sikkim	16279	0	15750	29	1	30	-
24	Tamilnadu	10945	55	11000	11	0	11	-
25	Telangana	24548	0	23998	20	0	20	10
26	Tripura	2874	0	2750	4	0	4	-
27	Uttarakhand	6004	0	6000	7	0	7	-
28	Uttar Pradesh	35746	0	49000	34	16	50	20
29	West Bengal	72266	0	88500	68	19	87	-
	Total	737761	53794	820709	744	103	847	119



Chapter-19

Vision 2030

- 1. The current population (2018) of the country is 1.36 billion which is expected to be 1.51 billon by 2030. As per the Indian Council of Medical Research (ICMR), the recommended dietary allowance (RDA), the protein requirement for a working male and female is 60 g and 55 g per capita per day respectively. In vegetarian population, the protein requirement is primarily contributed by pulses with supplementary sharing by other food ingredients. The domestic demand of pulses is 23 million tons *i.e.* 52 g/capita/day, which has increased to almost 53 g/capita/day.
- 2. To feed 1.51 billion population, the projected demand of pulses by 2030 is likely to be 35 million tons as per the behaviouristic approach (consumption ó 28.70 Mt + seed post harvest losses-5.72 Mt). This necessitates an annual growth rate of 3.57 *per cent*.
- 3. To meet the projected requirement and sustain the balanced production in pulses, the existing actual yield gaps of 439 kg/ha or 65 *per cent* under total pulses shall have to be a bridged besides bringing an additional area of 5-6 Mha under pulses.
- 4. Eradicating hunger and malnutrition is one of the great challenges of our time. One in three-suffer from some form of malnutrition. Of the 17 global goals of the world Food Programme 2015 adopted by global community for sustainable development to improve people lives by 2030, Goal-2-Zero Hunger- pledges to end hunger, achieve food security, improve nutrition and promote sustainable agriculture.
- 5. India is self sufficient in cereals and has achieved quite balanced, almost self-sufficient, pulse production during 2017-18. However, by 2030 for meeting out the 35 Mt requirement of pulses the existing (2017-18) productivity of 835 kg/ ha shall have to be raised to 1030 kg/ha in addition to bring additional coverage of pulses in a area of 5-6 Mha over the existing normal area. The per annum average growth in area and productivity shall have to be ensured at 1.7 *per cent* and 1.95 per cent respectively.
- 6. Besides WTO and Intellectual Property Regime (IPR) implications, pulses have a greater significance in the economy of rainfed regions *i.e.* 73 Mha area of the country and the prosperity of the youth, small and marginal farmers, which would mean better nutritional security, improvement in the production base through conservation of natural resources, livelihood generation high net returns to farmers though value addition and lowering the cost of production with an eye on export.
- 7. The rainfed eco system with more than 40 *per cent* of total population and 2/3rd or 66 *per cent* of livestock of the country is a targeted group for the government. The DAC&FW, in its resolve to double the farmersø income by 2022, envisions to sustain self sufficiency in pulse production, improve competitiveness through knowledge based technological interventions for improving nutritional security and sustainability of the production system.

19.1 Road Map-2030

In order to sustain the growth of pulses at various levels i.e. among the states, districts, within districts and to bridge the yield gap between FLDs and farmersø practice, DAC&FW has envisioned a road map with two pronged strategy:

I. Horizontal Expansion through bringing additional area under pulses, and diversification of rice-wheat system in Indo-gangetic plains (IGP) through popularization of short duration varieties of pigeonpea, kabuli chickpea, field pea and summer mungbean; Bringing additional area under pulses through promoting urdbean/mungbean cultivation in rice fallow in peninsular India and chickpea lentil in NEPZ and Chhattisgarh; promotion of pulses in intercropping viz., short duration thermo-insensitive varieties of mungbean/urdbean with spring sugarcane; pre-rabi chickpea with mustard/linseed; pigeonpea with groundnut/soybean/millets, etc.; development and popularization of urdbean/mungbean for late planting (mid Aug-early Sept in north India).

(Table-19.1): Potential of additional area under pulses

S. No.	Potential Crops/Cropping System/Niches	Specific areas	Potential Area	Target Area (2030)
			(Million	n ha)
1.	Intercropping			
	Mungbean with Sugarcane (Irrigated) and with cotton and millets (Rainfed uplands)	Western, central and eastern UP, Bihar, MH, AP and Tamilnadu	0.70	0.50
	Pigeonpea with soybean, sorghum, cotton, millet and G. Nut (Rainfed upland)	A.P., Malwa Plateu of MP, Vidarbha of MH, North Karnataka, Tamilnadu	0.50	0.50
	Chickpea with barley, mustard, linseed and safflower (rainfed)	South óeast Rajasthan, Punjab, Haryana, UP, Bihar, Vidarbha of MH	0.50	0.30
	Chickpea/Lentil with Autumn planted/ ratoon sugarcane	MH, UP, Bihar	1.00	0.60
2.	Catch Crops	l		
	Mungbean : Spring/Summer	Western & Central UP, Haryana, Punjab, Bihar, WB	3.00	2.00
3	Rice fallow			
	Chickpea	Eastern UP, Bihar, Jharkhand, Odisha, CG, WB	0.40	0.40
	Urdbean/Mungbean	AP, Tamilnadu, Odisha, Karnataka	0.50	0.40
	Lentil	Eastern UP, Bihar, WB, Assam, Jharkhand	0.30	0.30
	Lentil/Fieldpea	North east	0.10	0.10
4	Kharif Fallow			
	Urdbean/Mungbean	UP, Bundelkhad MP	1.20	1.00
	Total		8.20	6.10

II. Vertical Expansion through increasing productivity and bridging the yield gaps; development of high yielding short duration varieties having multiple and multiracial resistance to diseases; development of new and efficient plant types; development of input use efficient genotypes; exploitation of hybrid vigour in pigeonpea; popularization of improved crop management practices and bridging yield gaps.

Improving yield stability

Development of transgenic against drought and gram pod borer; efficient water management in rainfed areas; rainwater harvesting and recycling through farm ponds and community reservoirs; promoting short duration varieties in drought prone areas; promoting micro irrigation system; adoption of moisture conservation practices; development of location specific suitable varieties and ensure availability of quality seeds.

• Development of resilient pulse crops to climate adversaries

Development of resilient pulse crop varieties to mitigate the impacts of climate change; critical monitoring of diseases and pest dynamics with reference to climate change; production and supply of quality seeds through seed hub and ensuring seed production accountability to SAUs/ ICAR to organise location specific recommended latest varieties of pulses in their jurisdiction; active involvement of private sector, NGOs, & farmersø help groups in production of quality seeds; mandatory target to public sector seed corporations; popularization of seed village concept with buyback system; more incentive on production of seeds of new varieties; promotion of farmer to farmer exchange of seeds and seed village programme.

Reducing post harvest losses

Refinement and popularization of harvesters, threshers and graders; development of stored grain pest resistant varieties; popularisation of storage bins and mini dal mills; strengthening of FPOs and establishment of processing units; development and popularization of low cost safe storage structures.

• Ensuring attractive price to producers

Announcement of MSP well in advance; assured procurement and creation of procurement centres in production zones; development of organized markets for pulses; linking farmers with FPOs, aggregations and E-NAM (markets); promotion of export of pulses like lentil and kabuli chickpea and arid legumes; production of value added products and use of by-products; branding of produce and promotion of organic pulse production.

• Ensuring timely availability of critical inputs and advisory

Promotion of IPM technologies against Helicoverpa; ensuring timely availability of quality bio-pesticides- HaNPV, Trichoderma & herbicides e.g. Pendimethalin etc.; seed dressing of fungicides for controlling seed borne diseases; providing safe storage structures like Pusa Bin and warehouse facility; creation and maintain/ sustain of production units of quality bio-fertilizers and bio-pesticides; fortification of fertilizers

with specific nutrients like S, Fe, Zn, B etc., in specific regions; popularization of sprinklers and micro irrigation techniques in rainfed areas; establishment of single window input supply centres for cluster of villages; advanced forewarning and forecasting systems for pest and disease outbreaks.

• Efficient transfer of technology

Organizing farmers training and exposure visits; popularization of improved technology through mass media; close interaction of research organizations, state departments of agriculture and private agencies; market led initiatives for organized village level seed production to exploit the high demand for improved varieties of pulse crops as well as branding of local germplasm ex. Baigani Arhar in tribal belts of MP; exploiting the led commercial pulse processing units at village level; promotion of pulse production as cash crops in unconventional areas like hills, coastal and tribal belts of country; promotion of farmer information and communication technologies based pulse knowledge management to increase production and productivity of small holder farmers.

Advance Seed Plan

The DAC&FW has made advance planning for Quality Seed Production of Pulses for three years (2018-19 to 2020-21) in order to ensure sustainability growth of pulses production in the country.

(**Table – 19.2**): Seed Plan-Total Pulses (2018-19 to 2020-21)

(Oty. in qtls.)

Crops		2018-19			2019-20			2020-21	
	Normal	Seed	Total	Normal	Seed	Total	Normal	Seed	Total
	Req.	Bank		Req.	Bank		Req.	Bank	
Andhra Pradesh	52890	0	52890	54490	200	54690	58490	200	58690
Assam	13938	2789	16727	0	0	0	0	0	0
Bihar	5680	0	5680	6532	0	6532	7384	0	7384
Chhattisgarh	29553	299	29852	30417	251	30668	31249	316	31565
Goa	3.00	1.80	4.80	3.90	2.00	5.90	4.80	2.10	6.90
Haryana	42875	1550	44425	61375	2725	64100	42875	1925	44800
Gujarat	42480	4248	46728	57224	5722	62946	62640	6264	68904
Karnataka	94700	4735	99435	94700	4735	99435	95200	4760	99960
Maharashtra	100870	8200	109070	100669	11350	112019	102704	13950	116654
Madhya	95580	2215	97795	105630	2530	108160	112400	2650	115050
Pradesh									
Odisha	2100	0	2100	2150	0	2150	2200	0	2200
Rajasthan	219808	4263	224071	282828	4336	287164	306366	5362	311728
Tamil Nadu	13827	0	13827	14355	0	14355	15081	0	15081
Telanagana	77928	2200	80128	79487	3950	83437	81077	0	81077
Tripura	1881	0	1881	1944	0	1944	2074	0	2074
Uttar Pradesh	55992	0	55992	58270	0	58270	60805	0	60805
Uttarakhand	1355	62	1417	1400	62	1462	1450	62	1512
West Bengal	5950	160	6110	6210	160	6370	6415	160	6575
Total	857410	30722	888133	957685	36023	993709	988415	35651	1024065

Source: Seed Division, DAC&FW, N. Delhi




Fig.-19.1: State-wise seed plan total pulses

(**Table – 19.3**): Seed Plan: Kharif Pulses (2018-19 to 2020-21)

(Qty. in qtls.)

Crops		2018-19	•		2019-20			2020-21	giy. in qus.)
	Normal	Seed	Total	Normal	Seed	Total	Normal	Seed	Total
	Req.	Bank		Req.	Bank		Req.	Bank	
Andhra Pradesh	480936	0	480936	480936	0.00	480936	480936	0	480936
Assam	28829	5765	34594						
Bihar	41200	0	41200	47380	0	47380	53560	0	53560
Chhattisgarh	167450	1691	169142	156214	1578	157792	174430	1762	176192
Haryana	8349	2350	10699	9016	2450	11466	9916	2650	12566
Gujarat	52250	5225	57475	59400	5940	65340	67100	6710	73810
Karnataka	161950	8098	170048	162160	8108	170268	162500	8125	170625
Madhya Pradesh	612600	5675	618275	780460	6125	786585	949500	7311	956811
Odisha	42400	2000	44400	40900	2000	42900	42100	2000	44100
Rajasthan	230850	12438	243288	260000	12516	272516	289960	13073	303033
Tamil Nadu	50688	0	50688	53378	0	53378	55704	0	55704
Telanagana	85681	8000	93681	89964	8400	98364	94462	8400	102862
Tripura	2797	0	2797	2935	0	2935	3085	0	3085
Uttar Pradesh	435724	0	435724	448953	0	448953	459537	0	459537
Uttarakhand	1558	0	1558	2779.6	0	2779.6	3442.56	0	3443
West Bengal	47767	910	48677	49430	910	50340	52805	910	53715
Total	2451029	52152	2503181	2643906	48027	2691933	2899037	50941	2949978

Source: Seed Division, DAC&FW, N. Delhi




Fig.-19.2: State-wise seed plan kharif pulses

(**Table – 19.4**): Seed Plan: Rabi Pulses (2018-19 to 2020-21)

(Qty. in qtls.)

Crops		2018-19			2019-20			2020-21	y. III qus.)
-	Normal	Seed	Total	Normal	Seed	Total	Normal	Seed	Total
	Req.	Bank		Req.	Bank		Req.	Bank	
Kharif Pulses									
Redgram	293480	11311	304791	298460	13214	311674	306206	13387	319593
Greengram	217514	9498	227012	220814	8806	229620	249921	11961	261882
Blackgram	204355	7270	211624	205104	8681	213785	215252	7123	222375
Horsegram	4142	35	4178	4295	37	4332	4451	38	4490
Cowpea	16402	555	16957	16432	565	16997	16727	565	17292
Mothbean	30798	913	31711	40825	927	41752	45858	1029	46887
Guar	91800	1140	92940	153756	2193	155949	150000	1547	151547
Total Kharif	858490	30722	889213	939685	34423	974109	988415	35651	1024065
Rabi Pulses									
Redgram	1173	0	1173	1180	0	1180	1849	0	1849
Greengram	74312	2751	77063	73940	2569	76509	77676	2652	80328
Blackgram	100661	3910	104572	98848	3289	102138	101269	3290	104559
Bengal gram	1712765	113018	175093	197315	39655	201281	2176463	42236	2218700
			0	9		4			
Lentil	165394	2278	167673	170772	602	171374	183993	750	184744
Fieldpea	243665	3614	247279	252386	471	252857	282267	563	282830
Horsegram	4290	0	4290	4290	0	4290	4290	0	4290
Cowpea	1960	65	2025	2092	65	2157	2620	65	2685
Lathyrus	65101	1368	66469	67415	1376	68791	69732	1385	71117
Rajmash	535	0	535	562	0	562	590	0	590
Total Rabi	2369321	127005	2421473	2644082	48027	2692109	2900159	50941	2951100
Total Pulses	3227811	157727	3310686	3583767	82450	3666218	3888574	86592	3975165

Source: Seed Division, DAC&FW, N. Delhi




Fig.-19.3: State-wise seed plan rabi pulses




Fig.-19.4: Season-wise seed plan of pulses




Fig.-19.5: Crop-wise seed plan of pulses

19.2 Value Addition/Strategy Processing

- There is a need for systematic listing of available various value additions/processing
 Technologies developed by various research institutes (ICAR/CSIR/CFTRI/SAUs etc)
 and publication of these for wider circulation in vernacular languages. Steps to avoid
 wasting the vast potential of the post harvest business in pulses sector need to be
 urgently addressed.
- Development of cheaper and acceptable Dal supplements/substitutes to ease out pressure on pulses through PHT.
- Export oriented crop cultivation and milling, need diversification and modernization of
 post harvest technology with special emphasis on export oriented processing. The
 varieties having export value e.g. bold-seeded lentil (sagar masra) and chickpea
 varieties like Gulabi chana and õkateela chanaö and special Baigani arhar (grown in
 Mandla, Baiga tribes), need popularization.
- Cost effective processing plants/units need to be set up at potential pulse pockets to avoid farmersøhardships for transportation and carriage.
- HRD programmes on scientific technologies/storage may be imparted for food preservation, value addition of by-products.
- There is need for cost effective/easy to access containers and chemicals for processing which should be within the reach of the poor farmers.
- Possibilities of import of tin containers, whose cost appears to be prohibitive in the indigenous market or its alternative, need to examine.
- Various incentives and social amenities need to be provided to the farmers to encourage them for maximum production of the raw-materials (pulses) to feed the processing industries for corresponding output. This may be achieved through formulations of Self Help Groups (SHGs) in the potential pulse areas.

- Modern techniques of pulse production is the foremost thing without which, processing
 and value addition of Pulses are not possible. Farmers, therefore, need up-to-date
 techniques along with proper transfer of technology.
- Appropriate food processing facilities cannot only avoid the wastage of food, but will also lead to value addition thereby, income generation in the centralized fashion in rural areas.
- Traditional food processing technologies as well as high-tech and environment friendly production technology should be encouraged.
- The food processing industry/machinery has to conform to high hygienic standard. Strict adherence to the standard prescribed by competent authorities has to be ensured.
- There is need to make better coordination mechanism between State Departments of Agriculture, marketing, mandi boards and Food Ministry at state level so that even marginal/small farmers could process their own produce without going to the far-flung bigger industries/plants. Small scale pulse mills could also be installed at community level through FarmersøInterest Groups (FIGs)/SHGs etc.
- Pulse growers must be provided with storage bins and other equipment required in post
 harvest operations to increase the durability of produce that will further go to the
 processing units for value added by-products.

19.3 Strategy related to research issues

- To break the yield barriers, development of physiologically efficient plant types, **Use of Biotechnology** for speedy transfer of genes, conferring resistance to important diseases and pests for e.g. transfer of Bt gene in chickpea and pigeonpea for control of pod borer, pre-harvest sprouting of mungbean and also the mutation breeding.
- Exploitation of heterosis breeding by way of use of CMS in pigeonpea, need aggressive research to develop and refine the process cost effective seed production Technology.
- Integrated approach for the management of diseases, pests, drought, nutrient etc. need multi-disciplinatry research, development of multiple disease resistant varieties, transgencies for Helicoverpa pod borer and drought in chickpea and pigeonpea and MYMV in urdbean and mungbean and development of varieties having tolerance to temperature extremitie, etc are urgently needed to address down and unstable yieldø constraints in pulses.
- Research on validation and refinement of technologies, development crop modules, forecasting and fore-warning the incidence of pests/diseases need to be initiated and developed.
- **Research emphasis on minor pulse** (mothbean, cowpea, horsegram, fababean, rajmash and lathyrus) need to be strengthened on regional basis alongwith development of technology dissemination modules for different situations.
- Research back-up needed for change from low point input to optimum input technology for various cropping systems as well as for cultivation as sole crop alongwith the need for development of nutrient and water use efficient genotypes.
- **Pulse Ideotype requirement** for Irrigated Medium stature semi-erect and compact, responsive to high input and high HI.

- For **multiple cropping**, quick growing, short statures and synchronous in maturity.
- Under rainfed conditions, erect, tall, main stem with open canopy early flowering, larger size and number of leaflets with low osmotic adjustments are more desirable traits.

(Table-19.5): Crop Specific Strategy/Recommendations (var./ plant protection)

	Top Specific Strategy/Recommendations (var., plant protection)
Pigeonpea	- Early maturing pigeonpea can be grown in irrigated tracts of north-
	west Rajasthan, Haryana, Punjab and western U.P. and as post-rainy
	season crop in September in U.P., Bihar, Odisha, southern Gujarat,
	A.P. and West Bengal.
	- In case of M.P. inter crop or mixed crop with Soybean (late variety)
	in un-irrigated area may be taken especially in Vidisha, Raisen,
	Sehore, Bhopal and Narsinghpur districts.
	- Popularization of Dharwad system (transplanting Pigeonpea),
	Seedlings are raised in polythene bags, transplanted in main field at
	45 days with the apacing of 5 feet X 3 feet under rainfed condition
	 and 6 feet X 3 feet in irrigated condition. Nipping (2 times) of pigeonpea after 45 and 55 days.
	- Nipping (2 times) of pigeonpea after 45 and 55 days. - The productivity is proposed to be increased by adoption of
	improved crop production technology i.e. use of improved seed,
	NPV for control of Heliothis, providing irrigation at critical stages,
	use of gypsum and bio-fertilizers etc.
	- For reducing the vagaries of diseases Integrated Pest Management
	comprising of deep summer ploughing, mixed cropping with
	sorghum, discouraging ratooning or perennial cropping, planting in
	well drained field, seed treatment with Benomyl or thiram or
	carbendazim+thiram @ of 2-3 g/kg for reducing incidence of wilt
	and root rot and selection of diseases resistant varieties for
	cultivation. For biological control of wilt and root-rot, seed dressing
	with standard formulations of <i>Tricodermaviridae</i> or <i>T. harzianum</i> @
	4 g/kg should be promoted.
	- Use micro irrigation (precision) through drip irrigation.
	- Cultivation of pigeonpea on raised beds by using Broad Bed Furrow
	(BBF) Planter.
	- Use Sterlity Mosaic Resistant Var. (BDN 708, GTH-1, BRG 2, BDN
	711, Bahar, BSMR 736, Sharad, Pusa 9, BSMR 853), Phytophthora
	resistant (TJT 501, CORG 9701, JKM 189, Pant Arhar 291 etc.) Wilt
	resistant (VL Arhar 1, Vipula, GT 101, Maruti, BDN 2, BSMR 736,
	MA 6).
	- Use Pre-emergence herbicides like Pendimethalin @ 750-100 g/ha,
Pigeonpea	- Metribuzin 250-100 g/ha for weed control.
	- GTH-1 is the hybrid variety for tasgenic (Cry gene) against pod
CIT	borer.
Chickpea	The frontline demonstrations conducted by ICAR have clearly
	shown the potential to exploit the gram yields with the available
	technology. Improved varieties, use of recommended does of
	fertilizers, providing irrigation at critical stages of crop growth, application of gypsum/bio-fertilizers, use of NPV for control of
	Heliothis.
	- For Integrated Diseases Management (IDM), deep summer
	- 101 micgrated Diseases Management (IDM), deep summer

ploughing, crop rotation with non legumes, deep or late sowing, wider spacing and inter-cropping with any one among wheat, barley or mustard for effective control of wilt, root rot, ascochyta blight and other soil borne diseases, seed treatment with Benlate, Benomyl, Carbendazim or Thiram @ 2-3 g/kg is recommended with the *T. viridae or Bacillus subtilis or Gliocladiumvirens*@ 4 g/kg of seed and select wilt resistant (JAKI 9218, JG 6, WCG 3, RVG 201, RVG 101, PKV Harita (AKG 9303-12), Wilt tolerant varieties (RSG 902, BGD 103, Phule G 0517, GJG 3, PKV Kabuli 4, RVG 203) and Ascochyta blight resistant varieties (GJG 0809, Samarat (GNG-469) PBG-5, CSG 515) supplemented with management practices for wilt and root rot are the best options. Varieties suitable for saline areas GG-2 & PKV-2.

- Heat tolerant chickpea cultivars would be required for all late sown conditions (in rice-fallows; after a short season catch crop, such as potato and vegetables, in rabi season.
- The heat tolerant desi chickpea variety JG 14 was evaluated under late sown condition in UP, Bihar, Jharkhand, MP, Chhatisgarh and Odisha. JG 14 gave 10 to 25% higher yield than the check cultivars n late sown conditions.
- Use Pre-emergence herbicides like Oxyfluoren @ 150-250 g/ha , Pendimethalin 750-1000 g/ha for weed control.
- Use short duration varieties Desi: JG-11, JG 14, JG 16, JAKI 9218 &Kabuli: IPCK 2002-29, IPCK 2004-29, KAK 2, JGK-1 and for Rice fallow condition Pant G 186, BG 372, Rajas, RSG 963, Pusa 547, Vaibhav.

Blackgram (Urd)

- Yellow mosaic virus resistant varieties, namely VBN 6, IPU 94-1, Mash 391, LAM 752, Mash 479, IPU 2-43, LBG 625, LBG 685, Improved early maturing varieties with a large number of clusters like Mash 1008 and Pant U-30; Azad Urd 1,; PDU 1 Variety for spring season in north India PDU-1, Azad Urd 1, Shekhar 2 (KU 300), WBU 109, Mash 414 powdery mildew resistant variety CO 6, VBN 4 & 7, Gujarat Urd 1, IPU 2-43, WBG 26 and LBG 402, LBG 625, 685 & 623 (Prabha), KU 301, TU 94-2 for Rabi season. CO 6, ADT 5, Vamban 6 for rice fallows condition.
- To enhance the kharif productivity selection of appropriate variety resistant to YMV, in-situ moisture conservation to escape terminal drought, IPM, application of gypsum, use of bio-fertilizer.
- In case of **summer urd**, crop has to be grown under better management conditions, mostly inter-cropped with sugarcane and sunflower. It is necessary to use only recommended varieties for summer cultivation, seed treatment, use of gypsum, etc. In rice-fallow areas during Rabi, varieties resistant to powdery mildew are required to give more thrust in addition to other agronomic practices.
- **IPM Management**: Seed treatment with Thomethoxam 35 FS @ 2g/kg seed; installation of yellow sticky trap @ 20 /acre; sowing of 1 row of maize or tur after every 30 rows of mungbean as a barrier crop; removal of weeds and disease plant from the crop; spray of NSKE 5 % after 25 DAS or on

Blackgram (Urd)

- appearance of pest; foliar spray of thiomethoxam @ 0.3 g or Trizophos @ 4.0 ml/lit water.

- Control for **Tobacoco caterpillar**: Novaluron 10 EC @ 150 ml or Acepate 75 SP @ 800 g or Chloropyriphos 20 EC @ 1.5 lit. using 100 lit. of water/acre at the appearance of pest and repeat after 10 days if necessary.
- Use Pre-emergence herbicides like Oxyfluoren @ 150-250 g/ha , Imazethapyr 75-100 g/ha for weed control.
- Apply Pendimethalin + Imazethapr (Pre mix) 0.9 kg/ha as pre emergence or Imazerhaapyr 100 gm /ha as eraly post or Imazethapyr + Imazamox 50g/ha as early post for effective control of weeds in mostvof the pulsecrops.
- Hoeing of 20 & 40 DAS recorded higher weed control efficiency and it was comparable with EPOE Imaethapyr + Imazamox (RM) 70-80g/ha.
- Proper water management-border irrigation under flood and sprinkler/micro-irrigation under limited water availability particularly at pod filling stage.
- Timely availability of quality seeds of recommended varieties.
- Good tillage and crop stablshment practices-laser land levelling, use of new type seed-drills, Zero-tillage sowing in proper moisture, residue retension of previous crop.
- Use of pre-emergence herbicides (Pendamethalin @ 0.75-1.50 kg ai/ha) and one need based hand weeding.
- Use of phosphorus and Sulphur particularly after wheat and intercropping with sugarcane.
- Control of Thrips in Mungbean at pre-flowering (use of Dimethoate or emidacholoprid).
- Promising varieties against White fly OBG 33, KUG 503, AKU 10-2.

Greengram (Moong)


- In cropping system manipulation, sugarcane can be intercropped with mungbean in U.P. and northern Bihar cotton, pearl millet and groundnut can be inter-cropped in rainfed uplands of Maharashtra, Karnataka and Tamilnadu.
- The increase in productivity during kharif season is to be achieved by use of improved seed, seed treatment, use of weedicides, control of insects/pests through IPM, application of gypsum, providing irrigation in absence of rains, wherever possible. The average productivity obtained under the Frontline Demonstrations is about 7.8 qtl per ha suggested that the present productivity can be improved further with the use of available technology.
- Early sowing during spring (around 15th March), soil application of insecticide like Phorate or Carbofuran G. @ 1.0 kg a.i./ha for effective control of YMV and fungal diseases or chemical (7 Carbendasim + Thiram) seed treatment for reducing incidence of wilt and root rot disease.
- **IPM Management**: Seed treatment with Thomethoxam 35 FS @ 2g/kg seed; installation of yellow sticky trap @ 20 /acre; sowing of 1 row of maize or tur after every 30 rows of mungbean as a barrier crop; removal of weeds and disease plant from the crop; spray of NSKE 5 % after 25 DAS or on appearance of pest; foliar spray of thiomethoxam @ 0.3 g or Trizophos @ 4.0 ml/lit water.
- Control for **Tobacoco caterpillar**: Novaluron 10 EC @ 150 ml or

	75 90 0 000 911 11 20 79 0 15 11
	Acepate 75 SP @ 800 g or Chloropyriphos 20 EC @ 1.5 lit. using 100 lit. of water/acre at the appearance of pest and repeat after 10
	days if necessary.
	- Select short duration YMV resistant varieties of Mungbean like
	HUM 16, IPM 2-3, IPM 02-14, Pusa 0672, SML 668, Samrat (PDM-
	139), Pant mung-2, 4 & 6, IPM-99-125 (Meha) and having a
	potential to increase area in
Greengram	- spring/summer in U.P., Bihar, West Bengal, MP, Rajasthan, Punjab
(Moong)	and Haryana. Powdery mildew resistant varieties like TJM 3, VBN
	3, AKM 9904, PKV Green Gold TM 96-2, TARM-1 & 2, TARM-
	18, JM-721. Large seeded Pant M-5, Pusa Vishal, SML 668, HUM
	16, TMB 37.
	- Promising varieties against Thrips SML 1807, 1814, 1810, 1836,
	1837, LGG 486 and for White fly ML 1774, ML 1779.
	- Use Imazethapyr 30 g/ha as a Pre & Post-emergence & Imazapic 10
	g/ha as a post-emergence for weed control whereas, Co-7, Vamban 3, ADT 5 for Rice fallow condition.
	- Proper water management-border irrigation under flood and
	sprinkler/micro-irrigation under limited water availability
	particularly at pod filling stage.
	- Timely availability of quality seeds of recommended varieties.
	- Good tillage and crop stablshment practices-laser land levelling, use
	of new type seed-drills, Zero-tillage sowing in proper moisture, residue retension of previous crop.
	- Use of pre-emergence herbicides (Pendamethalin @ 0.75-1.50 kg
	ai/ha) and one need based hand weeding.
	- Use of phosphorus and Sulphur particularly after wheat and
	intercropping with sugarcane.
	- Control of Thrips in Mungbean at pre-flowering (use of Dimethoate
	or emidacholoprid).
Lentil	- Bold seeded varieties namely, DPL 15 and DPL 62, DPL 4046,
	Sapna, Priya, Pant L 5, Mallika, JL 3, IPL 81. Rust resistant
	varieties with different plant types ó VL-126, IPL 406, Pusa Masur 5, Shekhar Masur 2 & 3, Pant L-024, PL-8. Wilt resistant variety
	viz .VL 125, Moitree WBL 77, Pant L-6, VL Masur 129 & VL-133.
	Small seeded varieties Pant L 4, IPL 406, Pusa vaibhay, Pant L 406
	& 639, KLS 218, HUL 57. Pusa vaibhav, KLS 218, Pant L 639, DPL
	62, Pant L 5.
	- Provide seeds of improved varieties resistant to wilt and rust, seed
	treatment with fungicide and Rhizobium culture, irrigation at critical
	stage (pod stage) of crop growth, use of gypsum, as a source of
	sulphur and use of IPM for the control of pest/diseases.
	- Use Pre-emergence herbicides like Oxyfluoren @ 150-250 g/ha,
Peas	Imazethapyr 75-100 g/ha for weed control. - Use of leafless dwarf types of Peas with high yield for closer
1 Cas	planting (JAY (KPMR 522), HFPD 24, KPMR 400). Powdery
	mildew resistant varieties (IPF 99-25, IPFD 1-10, Paras, Pant Pea 14,
	VL Matar 42, Pant Pea 25 & 42 and Rust resistant variety Swarna
	Trapti, VL Matar 47, Aman (IPF 5-19).
	- Under the Frontline Demonstrations, yield levels to the tune of 1790
	kg/ha have been reported which is almost double the normal yield

	levels. Field peas normally receive better management and thus farmers pay adequate attention to this crop. However, the targeted productivity would be achieved by providing seeds of better varieties resistant to powdery mildew, seed treatment, application of gypsum, managing rust disease and providing irrigation, etc. - Early sowing (during 1 st week of October) to escape onset of powdery mildew and rust diseases in NEPZ. - Fungal seed treatment to reduce incidence of seed rot and root-rot, two-three foliar spray of wettablesulphur (0.3%) for control of powdery mildew and rust. - Use Pre-emergence herbicides like Oxyfluoren@150-250 g/ha, Pendimethalin 750-1000 g/ha and Metribuzin 250-100 g/ha for weed control.
Lathyrus	 Lathyrus is most commonly grown as Utera in rice. Important states are Chhattisgarh, Odisha, Maharashtra, Madhya Pradesh, Bihar and West Bengal. Increase in productivity in case of lathyrus would be obtained by better management of utera cultivation. Variety Bio L 212 (Ratan), Prateek (ODAP-0.109%) & Mahateora (ODAP-0.074%) a low toxin Lathyrus can be grown in rice fallows of Uttar Pradesh, Bihar, Odisha, West Bengal and Chattisgarh. Strees tolerant varieties Prateek, GNG 1581, Mahateora.
Moth	 Adoption of improved crop production technology i.e. use of improved seeds, NPV, irrigation IPM, INM disease resistant varieties, weed management and other package of practices at critical stage. Yellow Mosaic resistant varieties TMV-1, Rajasthan Moth 257, RMO 2004, CZM-45 & 99, JMM-259 and for Drought tolerant is RMO 423.
Rajmash	 The cultivation of Rajmash may be promoted mainly in North East Plain Zone. Rajmash and other beans can be grown profitably in irrigated areas of Uttar Pradesh, Maharashtra and Gujarat Varieties suitable for the plains of northern India for Rabi season available in different grain colours, namely, Variegated (PDR 14 or Uday), Red (HUR 137) and White (HUR 15). Anthracnose resistant varieties are Phule Surekha (KOF B-4), Varun (ACPR 94040), VL Rajma 125, VL Bean 2 and rresistant to Bacterial Blight Mosaic Virus (BCMV)- Amber (IPR 96-4), IPR 98-5 (Utkarsh), IPR 98-3-1 (Arun) etc.,

19.4 Policy Related Strategy

- In order to make a break-through in expansion of area under pulses, short duration varieties of pigeonpea to need based replacement of soybean, and short duration early maturing chickpea varieties for late sown conditions after paddy harvest, need popularisation, through demonstration.
- Better package of practices especially the inter-cropping Package, developing effective and adaptive IPM against major disease and Crop Management etc. need to be documented and popularized across the country.
- Development/promotion of perfect technology for *utera* cultivation with a view to divert an existing area of about 6 to 7 lakh ha under lathyrus towards chickpea lentil, cultivation.
- Dove-tailing and convergence concept should be materialized; provisions of assured irrigation in rabi and summer/spring season should be made on priority-basis.
- Developing strong seed production and distribution chain to achieve seed replacement rate of 33% by 2016-17 for all pulses.
- Creation of seed banks to meet seed shortage needs and for calamity situations by associating public as well as private sector seed companies. Monitoring of seed hub by SDA.
- An area of approximately 1.3 million hectares of a large tracts of Rice-fallow land (because of unirrigated conditions and properties of soils to hold moisture for shorter duration), and 2.47 million hectares under inter cropping in different cropping situations may be brought under pulses through aggressive crop coverage campaign.
- Delineation of un-exploited potential belts in non-traditional areas like watershed, introduction of pulses during non-traditional seasons under irrigated conditions, inter/mixed-cropping, summer cropping etc.
- Creation of production units for Nuclear Polyhydrosis Virus (NPV) with all the KVKs and integration of development and research at district level.
- Emphasis on sprinklers and micro-irrigation systems to promote pulses in irrigated area with efficient water management.
- Provide an effective market mechanism to pulses by minimizing the price fluctuations. FPOs, Self-help groups (SHGs), Farmers Interest Groups (FIGs) for effective market improvement can be organized.
- Development and Dissemination of location specific agronomic package of practices by SAUs, Skill development packages *etc.*, by aggressive ToT programmes.



Chapter - 20

Way Forward-2030

- 20.1 It is an established fact that a human body requires a daily intake of about 55 to 60 g of protein. The protein malnutrition and other health indicators like maternal and infants mortality rates, anaemia etc, is a matter of concern for the government. Nutritious food has direct bearing on health and affects work performance of the people. Out of the 22 amino acids required in the human diet, the body supplies 14. The remaining eight have to come from food. If all the eight amino acids are present in a single food item, it is called a complete protein food. Since all proteins from animal sources are complete proteins, it is easy to meet the dietary protein requirements of people with non-vegetarian food as the main diet. However, for vegetarian population the main sources of protein are leguminous plants to which the pulses belong. However, in general, pulses have lower concentrations of protein than animal sources. Besides, none of the pulses, except soybeans, are complete proteins. Therefore, combinations of two or more pulses are needed in a vegetarian diet. Dairy products, which are complete proteins, may also be used to supplement pulse proteins in vegetarian diets.
- 20.2 Keeping in view the production issues and recent pulse scenario in India, it is now established that the domestic supply of pulses is able to meet the growing demand of our consumers. Availability, price and the dietary preference for specific types of pulses in different parts of the country is largely responsible for this. About 29 million hectares of land is under pulses producing about 24 million tons annually. Our population is increasing as well as demand will also increase which we have to resort to increase the production. It is a challenge for us to ensure supply of pulses as pulse crops are primarily taken up for cultivation under rainfed condition in poor soils and are prone to production losses due to moisture stress.
- 20.3 Two major issues have emerged in respect to pulses production in the country. First, the limited genetic potential for high yields and second their vulnerability to pests and diseases. Compared to other food grains crops, yield potential of the pulses has been rather low. Newer varieties of pulses need to be developed so that the crop cycle fits well into cropping systems that the farmers adopt. Another important issue is limited mechanization potential, especially for planting and harvesting of the crop. Suitable plant types need to be developed for mechanical harvesting with pods above the canopy and sturdier plants.
- 20.4 Due to the high protein content in pulses, the crop is highly vulnerable to pests and diseases. It is estimated that about 30% of pulses crops are lost on account of pest attacks and diseases every year. Attack by pod borer and pod fly is so severe that the entire standing crop is devastated at several times. Research efforts need to be prioritized in this direction by employing modern biotechnological tools for crops for developing pest resistant varieties of pulses.

- 20.5 Weed is one of the problematic areas for limiting yield both in kharif and rabi pulses. Manual weeding appears to be difficult in future to manage this weed menace, therefore development of herbicide tolerant pulses as well as varieties suitable for mechanical harvesting is the necessity of the time to minimize the labour-intensive crops.
- 20.6 Another problem peculiar to the Indo Gangetic plains is the menace of large scale grazing by blue bulls. We need to support the efforts of the farmers for higher acreage under pulses crops without contravening the legal provisions of the Wild Life Protection Act which prohibits killing of these animals. There is thus a huge possibility and potential of bringing innovative solutions to save the pulses crops and encourage more intensive promotion of production technologies.
- Besides varietal research, we need to address the issues relating to farmersø preference 20.7 for the competing crops to pulses through development and promotion of crop production and crop protection technologies. Under National Food Security Mission, BGREI and CDP, these aspects have been taken up for more aggressive promotion of available technologies through cluster demonstration to ensure that the farmers are able to harvest better crops. The Government significantly increased the Minimum Support Price of the pulses and strengthened pulses procurement mechanism by designating additional central agencies to support the farmers. In fact, the minimum support price has been doubled in last three year with quantum jump given this year with an increase of more than 150% of cost of production during Kharif 2018. Use of drip irrigation in pigeonpea and agronomic practices like transplantation and nipping of branches are showing very encouraging results. Our import & export policies are linked to our ability for better crop forecasts and the principle of balancing the farmer interest in a manner that the prices are not distorted and the Indian farmer continues to get a good return for their produce. This will hopefully help in better planning and management of supply chain. With better yields through development of pest and disease resistant varieties, increased MSP support and lessons learnt from exchange of knowledge, the Indian farmer will definitely adopt pulse based cropping systems to produce more pulses with increased acreage. This will significantly ease the supply side constraints in the years to come.
- 20.8 To meet the challenges faced by the pulses sector, government has given emphasis on research efforts for developing biotic stress resistant and stress tolerant varieties, to be encouraged along with public-private initiatives for better logistics planning and handling of pulses. The crop cycle should be such as to fit into the overall cropping system that the farmer takes during the year. Aggressive promotion of available technologies under the cluster front line demonstration has been taken up. The pulse procurement mechanism has been strengthened by designating additional central agencies and the minimum support price (MSP) has been significantly increased to more than 150% of the production cost. Thus, by balancing the farmer interest in such a way that the domestic prices are not distorted and the Indian farmer continues to get a good return for his produce, the use of new production technologies and agronomic practices, and government support put together could maintain to self sufficiency.

- 20.9 Aggressive promotion of available technologies under the cluster front line demonstration has been taken up. The pulse procurement mechanism has been strengthened by designating additional central agencies and the minimum support price (MSP) has been significantly increased to more than 150% of the production cost. Thus, by balancing the farmerøs interest in such a way that the domestic prices are not distorted and the Indian farmer continues to get a good return for his produce, the use of new production technologies and agronomic practices, and government support put together could maintain to self sufficiency.
- 20.10 Given the important role that pulses play in the human diet, their availability needs to be increased indigenously. The possibility of improving pulse productivity two to three times through existing varieties and available package of technologies is well demonstrated in scientific experiments.

Focusing on individual pulse crop group

20.11 Enhancing chickpea production

- Chickpea has tremendous scope for expansion in rice fallows of eastern U.P., Bihar, Orissa and West Bengal, Bhal areas of Gujarat and through diversification of rice-wheat system in Indo-gangetic plains. This will not only increase the area of chickpea but also help in sustaining cereal based cropping system in long run. An estimated additional area of 1.0 to 1.5 million ha area can be brought under chickpea cultivation. Under intercropping of chickpea with barley, mustard, linseed and safflower (south-east Raj., Punjab, Haryana, UP, Bihar, Vidarbh of MS), Autumn-planted/ Ratoon sugarcane (MS, UP, Bihar), chickpea under rice fallow (eastern UP, Bihar, Jharkhand, Odisha, CG, WB), an additional area of 02 Mha may be brought.
- A large number of improved varieties and matching production technologies have been developed in chickpea which showed its impact on increasing the production and productivity but there are some critical gaps which could not be properly addressed so far.
- In the last 4 decades, the area, production and productivity of chickpea have fluctuated widely. Some of the states like Punjab, Haryana, Uttar Pradesh and Bihar have lost considerable area where as the states like Andhra Pradesh, Maharashtra, Karnataka have brought additional area.
- The growing condition is altogether different in southern states from northern states, there is need to develop more varieties and production and protection technologies to address the critical gaps in the region. Still there is lot of scope of increasing production and productivity in these states by developing high yielding varieties with short maturity duration, terminal heat and moisture stress tolerance. There is also scope of refinement in production technologies for better establishment of crop and increasing the productivity.
- The chickpea programs need to be dynamic and efficient in rapidly developing varieties needed for adaptation in existing and evolving cropping systems and have the traits

preferred by farmers, industries and consumers. We need to also consider the environment and the efficiency in use of available resources. Traits like phosphorus acquisition efficiency (PAE) and biological nitrogen fixation (BNF) efficiency need due consideration. Though chickpea is a good source of protein, carbohydrates, dietary fibre, minerals (Fe, Zn, Ca and Mg) and other important nutrients that are essential for human health, there is a scope for further improving nutritional quality of chickpea.

- Protein content of present day cultivars is usually in the range of 18ó22% whereas, much larger variability (14ó32%) exists in the germplasm which could potentially be exploited to develop high protein (×25%) varieties. Similarly, wide variation has been observed for iron and zinc contents in the germplasm, which can be exploited for developing high iron and zinc varieties. It is important to develop nutritionally enhanced varieties as the consumers would get a higher amount of protein and other nutrients from the same amount of chickpea consumed.
- The applied research and efforts towards short duration varieties for bringing additional area in eastern India, abiotic and stress tolerance, machine harvestable and herbicide tolerant varieties, extra-large kabuli chickpea for premium domestic and international market, diversification of rice-wheat system, efficient cropping systems, varieties with enhanced phosphorus acquisition efficiency, management of pod borer (*Helicoverpa armigera*), incorporation of multi-racial and multiple disease resistance, nutritionally enhanced varieties, popularization of extra-large seeded kabuli varieties and machine harvestable varieties among farmers etc. shall have to be continued.

20.12 Enhancing pigeonpea production

- Pigeonpea or redgram, the second most important crop in total pulses and the top ranking kharif season pulse crop, is a preferential pulse across the country. Owing to its characteristic behaviour as a contingent and risk management crop, the two pronged strategy i.e. area expansion and bridging the yield gaps (61 percent or 530 kg/ha over state average yield) shall be the way forward.
- Expansion in area (Horizontal expansion): There is less scope in direct expansion of pigeonpea area overall in the country. However, a new concept of rabi pigeonpea is developing in Chhattisgarh state parts of eastern UP, Bihar, West Bengal, Odisha, Gujarat and eastern Madhya Pradesh state in those areas where paddy cultivation has a limitation due to irrigation water resources or after early paddy and hood prone areas where field get waterlogged during rainy season, there can be a scope of rabi pigeonpea. There is a great demand of early pigeonpea (120-130 days) by the soybean growing farmers especially of Malwa region where soybean is a dominating kharif crop. Because of its early harvest farmers yet cannot afford to replace soybean as land can be made free for rabi cultivation (double cropping) as soon soybean is harvested. Chickpea and wheat fetches higher income to the farmers of Madhya Pradesh. Therefore, alternate replacement crop for soybean has not yet paved its way in soybean areas in spite of decreasing productivity of soybean crop in the state. The ways to increase the pigeonpea area is intercropping of pigeonpea with soybean in kharif season. Under NFSM Cluster

Demonstration and CFLDs, farmers are being demonstrated to include one or two lines of pigeonpea alternated after 6 or 8 lines of soybean. Govt. of India is also promoting intercropping of pigeonpea with soybean in Madhya Pradesh by taking Frontline Demonstrations of AICRP Programmes. The practice is being adopted in some areas like Bhopal and Raisen districts where underground water is scarce.

- Making availability of early maturing: Pigeonpea naturally is a perennial crop. However, through intensive research efforts scientists could bring down its maturity upto 120 days and some varieties like UPAS 120, ICP 151, ICPL 87, TT 401, TJT 501 have been developed and released for cultivation. However because of its natural habit, any change in micro climate results drastic change in growth behavior of pigeonpea crop. Varieties maturing in 120 days in South zone, gains 140-150 days in Central zone (especially in Madhya Pradesh). The state has released Mid-early duration variety pigeonpea variety TJT 501 in 2009 and this variety has gained large popularity due to its stability in yield. However some minor changes in maturity cannot be avoided due to inherent nature of the crop. Looking to the immense need of early duration variety (~120-130 days).
- During kharif 2018, early duration varieties Pusa-16 (125 days), ICPL-151 (140 days), ICPL-87 (130 days) and other short maturing varieties of southern region have been taken for seed production and multi-location trials at 06 locations of MP (KVKs/ ZRS-Shajapur, Rajgarh, Dewas, Khargone, Sehore, Ujjain).
- It is proposed to consider re-notification of varieties in central region of MP & Vidharbha of Maharashtra, AP, North Karnataka and Tamil Nadu under intercropping (pigeonpea with soybean, sorghum, cotton, millet and rainfed upland groundnut with 01 row of pigeonpea + 6, 8 or 10 rows of principal crop). Potential area of 0.5 Mha has been envisioned by 2030 in addition to ever highest area coverage of 5.39 Mha during 2016-17 (existing normal area of 4.2 Mha).
- The Bhabha Atomic Research Centre (BARC), in association with RVSKVV, Gwalior (SAU) has been roped in for mutation breeding to bring 120 days variety for MP, the 3rd ranking Tur state in the country.

Adoption of recommended Package of Practices

Row spacing: Keeping recommended row spacing in sole pigeonpea cropping is an important factor affecting the grain yield. In spite of scientific recommendations of 60 cm for early, 75 cm for Mid-early and 90 cm for medium duration pigeonpea varieties, farmers are adamant to plant pigeonpea in a closer spacing of not more than 45 cm especially in Madhya Pradesh. In addition to it within row minimum spacing of 20 cm is not adopted. This results in very close and high plant population, limits the crop growth with lower productivity.

Ridge sowing: Madhya Pradesh is experiencing high rainfalls during past 2-3 years. Continuous rainfall increases not only atmospheric humidity, but continuous moisture

(water stagnation) during early crop growth stages increase incidence of Phytophthora disease in pigeonpea causing heavy toll of seedlings, reducing optimum plant population. Scientific method of ridge planting can save the pigeonpea seedlings, allowing water drainage and reduces the intensity of mortality due to Phytophthora disease.

• Development of early duration varieties/hybrids

- Hybrid pigeonpea ICPH-2671 and ICPH-2740, in 2000 on farm trials have exhibited 47% and 42% yield advantage over the best local varieties, would have to be taken forward with effective hybrid seed production training programmes through KVKs/SAUs.
- Since last 4 years with the collaboration of ICRISAT, efforts are being made to develop early duration pigeonpea hybrids in pigeonpea. One hybrid RVICPH 2431 has reached to the level of AVT 2 and is likely to be released in 2-3 years.
- Collection and identification of early pigeonpea genotypes.
- Although there are many extra early and early genotypes/varieties released and available, these genotypes have found unproductive in Madhya Pradesh condition. The major constraint of low yield is heavy incidence of pod borer complex including Marucca from flowering itself. Some accession of determinate (DT) and non-determinate (NDT) early pigeonpea have been acquired from ICRISAT and tested since 2-3 years. Although someof them flower and pod in south condition, the pod setting of these super-early accessions in Madhya Pradesh condition is yet under adaptation. In addition to it early duration CMS lines are also being exploited for development of

In addition to it early duration CMS lines are also being exploited for development of early maturing pigeonpea hybrids.

• Research Projects on Early pigeonpea:

- Research funding to ICRISAT (2017-18 to 2019-20, Rs. 8.90 Cr) for enhancing genetic gains in chickpea & pigeonpea and scaling up and popularization of pigeonpea in Karnataka, MS, Telangana, AP and Odisha (2018-19, Rs. 6.50 Cr), has been done to ensure sustainable growth to these two crops.

20.13 Enhancing Mungbean, Urdbean and Lentils Production

- Mungbean, urdbean, lentil, lathyrus, rajmash and fieldpea together accounts for about 37% of total pulses production in India. Therefore, these crops assume greater significance in bringing self sufficiency in pulses in the country. Research and input supports are crucial in further advancing the production and productivity of these crops. It is envisioned that the new initiatives will not only improve the productivity of these crops, but also help in promotion of pulses in different production nitches. These efforts are likely to bring 2-2.5 Mha additional area under pulses which can provide 1-1.5 mt additional production to national pulses basket in next three years.
- In mungbean, low and instable productivity due to severe incidence of MYMV, Jassids
 and Thrips and poor crop management often noticed in major growing states like
 Rajasthan, Maharashtra, Madhya Pradesh and Uttar Pradesh. IPM strategies for control
 of pests use of resistant cultivars, seed dressing with systemic insecticides and timely
 spray of recommended insecticides and use of bio-fertilizers, sulphur (@20 kg/ha) and

- zinc (@ 15 kg/ha) and foliar spray of 2% DAP/urea at flowering stage can help improve the crop management.
- Mechanical harvesting in mungbean is prevalent in some parts of MP, Rajasthan, Punjab, Haryana and western Uttar Pradesh. Other states shall have to be brought under mechanical harvesting in mungbean with the development of suitable genotypes and appropriate machines. There is adequate scope of horizontal expansion under mungbean and urdbean to the tune of >3 Mha by promotion of the crop under intercropping with cotton, pigeonpea and millets in the states like Karnataka, Odisha and UP, Madhya Pradesh; with spring sugarcane in Uttar Pradesh and bringing additional area under rice fallow in rabi season in Tamil Nadu, Andhra Pradesh and Odisha.
- In lentil, inadequate availability of quality seeds of improved varieties in major states like Uttar Pradesh, Madhya Pradesh and Bihar has been the major constraint. Promotion of quality seed production through seed hubs/ formal and informal seed production systems shall have to be promoted. Wilt in Uttar Pradesh and Madhya Pradesh and Rust in Uttar Pradesh and Bihar have been crucial diseases which can be controlled through IPM with the use of resistant varieties and seed dressing with bio-agents like Trichoderma. Availability of good quality bio-agents are important to tackle menace of root diseases in the crop. Bio-agents and bio-fertilizers units need be strengthened/ installed in these states for timely availability of such crucial inputs. Terminal moisture stress for cultivation of lentil in Uttar Pradesh and Madhya Pradesh further aggravated the problem of realizing productivity. Promotion of extra short duration varieties (~ 100 days maturity), adoption of in situ moisture conservation measures, water harvesting and recycling need to be popularized and strengthened. Provision of life saving irrigation through sprinklers can boost up the productivity of rabi pulses enormously. Increased yield of lentil in utera cultivation in Uttar Pradesh and Bihar can be obtained by seed priming, enhanced seed rate and inoculation with rhizobium and foliar spray of 2% DAP / Urea at flowering and pod development stage. Use of lime in acidic soils of West Bengal, Bihar and Assam can be popularized in programme mode.
- Pulses for rice-fallow cultivation: Promotion of pulses such as lentil in Assam, West Bengal, Jharkhand, Bihar and eastern Uttar Pradesh and Mungbean in coastal Andhra Pradesh, Tamil Nadu, and Odisha under *utera* cultivation after the harvest of short season rice can provide ample opportunity to produce additional food from fallow lands. This requires appropriate varieties and matching production technology of lentil, Mungbean and Urdbean which may suit to rice-fallow ecology. Seed priming, foliar nutrition, planting techniques using zero-till seed-cum fertilizer drill and post-emergence herbicides may play vital role in improving productivity of rice-fallow pulses.
- Use of post-emergence herbicides: Lentil is poor competitor of weeds. Similarly, weeds
 are serious menace for kharif Mungbean and Urdbean. Application of pre-emergence
 herbicides could provide protection from weed from 20-25 DAS. One or two manual
 weeding requires further protection during growth stages of the crop. Non-availability
 of manual labour for weeding in these crops is crucial for realizing good crop growth.

Successful application of post-emergence herbicides like imazethapyr and quizalofop in soybean encouraged to identify molecules which can provide protection against weeds in lentil, Mungbean and urdbean. Therefore, efforts are being made to use of post-emergence herbicides for MULLaRP crops.

- Improved adaptation to climate change: In central India, lentil suffers heavily from terminal moisture and heat stress. Similarly, the productivity of Mungbean and Urdbean affected due to sudden increase in temperature during flowering. Therefore, efforts are being made to identify suitable genotypes tolerant moisture stress and high temperature. The tolerant genotypes will be used in varietal development programme for improved adaptation to climate change.
- Increased attention will also have to be paid to development of sustainable production systems that protect the natural resource base as well. Recent evidence of resource degradation and declining productivity in some intensively cropped areas is of particular concern. Also population driven intensification of agriculture without the use of external inputs, is leading to a serious problem of decline in soil fertility.
- Resource use efficiency in agricultural production has been a major concern in India. Due to decrease in resources (land and water) and increasing demand by other sectors limit the availability of resources for agriculture and that too for pulse crops. Thus, any decrease in marginal returns as predicted by the law of diminishing returns is more or less compensated by the benefits of other technological changes. Therefore, development of resource use efficient genotypes, precisely irrigation water and fertilizer delivery system, slow release fertilizers, integration of bio-fertilizers with other fertilizer sources, pulse based cropping systems including intercropping and other resource conservation agronomic practices may help to utilize available resources to the maximum possible level. While developing resource efficient technology, due consideration may be given for õeco-efficientö agriculture.
- Improved agricultural technology, irrigation, livestock sector and literacy will be most important instruments for improving the nutritional security of the farm-households. Watershed development and water conservation techniques will have far reaching implications in increasing agricultural production in the rainfed areas. Need based and location specific community programs, which promise to raise nutritional security, should be identified and effectively implemented. Small-mechanized tools, which minimize drudgery but do not reduce employment and add value to the working hours, are needed to enhance labour productivity. Special safety nets should be designed and implemented for them. There is need to disseminate widely post-harvest handling and agro-processing and value addition technologies not only to reduce the heavy post-harvest losses but also improve quality through proper storage, packaging, handling and transport.
- Farmersø perception and literature reviewed both revealed that inclusion of legumes in dominated cereal-based cropping system increased the yield of subsequent cereal crops

and reduce the fertilizer cost too resulting decrease in cost of production and increase profitability. Inclusion of pulses in various ways is technically feasible as well as economically viable in long run.

20.14 Convergence approach of programme implementation

- The various initiatives taken by the government between 2015-16 to 2017-18 under all CSS/CS on crop and agricultural development shall have to be converged. The infrastructure created on water resources, customs hiring centres, seed hubs and EBSPs, and the capacity buildings of famers. FPOs, SHGs and extension workers would be needed to fully utilized in favour of the growth of the pulse sector.
- To sustain the production and fulfill the demand as also to achieve the target of Zero Hunger by 2030, the DAC&FW has formulated the strategy by delineating the districts of the country into four categories: i) High area high productivity districts (HAHP) ii) High area low productivity districts (HALP) iii) Low area high productivity districts (LAHP) iv) Low area low productivity districts (LALP).
- The Government has resolved to make strategic interventions in all four category districts involving the approach of horizontal and vertical expansion, inclusion of traditional wisdom as well as recent improved technologies.
- On the based on yield during 2016-17 workout crop-wise and state-wise categories of district having LALP (Low Area Low Productivity), LAHP (Low Area High Productivity), HALP (High Area High Productivity), HAHP (High Area High Productivity) than its average area and productivity, which is attached *Annex-VI*.
- This Directorate also categorized districts on the basis of above and below national average yield for further need strategies and follow the increasing production of the pulses. List of districts is enclosed with *Annexure VII*.
- The major ten districts having highest production under individual pulse crop category and the production analysis to state as well as all India is attached with *Annexure-VIII*.

Glimpses of Activities/ Outreach Programmes on Pulses Development



Closing Ceremony of IYOP at Agra



Honøble Agri. Minister, *M.P. (Shri. Gauri Shankar Bisen)* and Union RDMinister (Shri. Narendra Singh Tomar) at Tikamgarh: BundelkhandSrajan 2017-National Agriculture Expo.

GLIMPSES OFACTIVITIES/REVIEW



Secretary (DAC&FW), GoI, visiting ZARS, Pawarkheda, Hoshangabad, (M.P.)

Secretary (DAC&FW), Review of Mechanization/CHC, CFMTTI, Budni, (M.P.)



Secretary, (AC& FW) inaugurating E-KisanSarthi portal under CHC-MP



Secretary (DAC& FW) adressing Krishak Sammelan at Sonkachch, Distt. Dewas

Overseas Consultations on Pulses





FAO Regional Initiative on Zero Hunger Challenge
Regional Inception Workshop for Regional TCP on Creating Enabling
Environments for Nutrition-Sensitive Food and Agriculture to Address
Malnutrition

GLIMPSES OF ACTIVITIES/REVIEW



IYOP-2016: National Workshopõ Pulses Opportunities in Central and Southern Statesö (Feb, 03-04, 2016) (ICAR-CIAE-M.P)



IYOP-2016: "National Workshop/Brainstorming Session on Promotion of Pulses in New Niches: Summer Cultivationö (Feb. 9-10, 2016)

(ICAR-IIPR, Kanpur)

GLIMPSES OF ACTIVITIES/REVIEW/FIELD VISITS



IYOP-2016:Farmers Training õProduction and Protection Technology of Pulses, KVK-Ichhawar, Sehore







IYOP-2016:Farmers Training õProduction and Protection Technology of Pulses KVK- Raisen

NATIONAL LEVEL MONITORING TEAM (NLMT) VISITS



Propagation of Hybrid cultivation : Tur (Var-ICPH-2740) Dist.-Gulberga, Karnataka

Area expansion through intercropping:
Tur (Transplanted) + Maize (Sweet corn)
District-Chhindwara, (M.P.)



Resource Use Efficiency: Arhar (Transplanting + MIS-Drip)
District-Betul (M.P.)

Organic Farming: Pigeonpea Nursery for Transplanting District- Narsinghpur (M.P.)

NFSM ACTIVITIES/MONITORING



Mechanization: NFSM-RCT, Tractor District-Harda, (M.P.)



Local Initiative : Spiral Grader, District- Indore (M.P.)



Conservation Agriculture: Happy Seeder, District- Betul, (M.P.)



Local Initiative: Processing Unit -Dal Mill, Chhattisgarh

NFSM ACTIVITIES/MONITORING



Crop- Chickpea, Variety JG-14 Harvesting, Madhya Pradesh

Seed-hub Production of Mungbean at Dantewada, Gujarat



Cluster Demo.-Gram: District-Harda, M.P.

NFSM-Demo., Gram (JAKI 9218),Block-Lohandiguda, District-Bastar, CG

NFSM ACTIVITIES/MONITORING



Crop- Pigeonpea, Variety TJT-501 Hand dibbling on ridge, Madhya Pradesh

Crop- Pigeonpea, Variety ó Rajiv Lochan Hand dibbling on ridge, Chhattisgarh



Innovation: Intercultural operation of Gram District-Indore, Madhya Pradesh



NFSM-Outreach Programme in Naxal Affected Area , Bastar, Chhattisgarh District- Bastar, Chhattisgarh

NFSM-PULSES IN TRIBAL DISTRICT- (BAIGA TRIBES)



Vill. Silpidi: Kisan Gosthi/Baiga Tribes, District-Mandla, M.P.

Vill. Silpidi: Indigenous Storage Bin/ Baiga Tribes, District-Mandla, M.P.



RKVY- Solar Fencing, District-Mandla, M.P.

BGREI (Check-Dam), Block-Tokapal, District-Jagdalpur (Bastar), C.G.

1. NFSM - PULSES

1	*Demonstrations on Improved Technologies: (a) Cluster Demonstrations (of 100 ha each) (b) Intercropping Demonstration	Rs.9000/-ha Rs.9000/-ha
	(b) Intercropping Demonstration	
		Rs 9000/-ha
		135.7000/ Hu
	(b) Cropping System based Demonostration	Rs.15000/-ha
2	Distribution of Seed minikits on Pulses & Oilseeds	Free of cost
3	Distribution of Certified Seeds:	
	(a) HYVs seeds (less than 10 year)	Rs.5000/-Qtls or 50 % cost
	(b) HYVs seeds (more than 10 year) limited to 20 % of total seed distribution	Rs.2500/-Qtls or 50 % cost
	Production of Seeds: (for varieties less than 10 year)	Rs.5000/-Qtls or 50 % cost
	Integrate Nutrient Management:	
	(a) Micro-nutrients	Rs.500/-ha
	(b) Gypsum/80% WG Sulphur	Rs.750/-ha
	(c) Lime	Rs. 1000 /- ha
	(d) Bio-fertilizers	Rs.300/-ha
4	Integrated Pest Management (IPM)	
	(a) Distribution of PP Chemicals	Rs.500/-ha
	(b) Weedicides	Rs.500/-ha
5	Farm Implements & Equipments:	
	(a) Manual Sprayer	Rs. 600/Unit
	(b) Power Knap Sack Sprayer	Rs.3000/Unit
	(c) Zero Till Seed Drill	Rs.15000/Unit
	(d) Multi Crop Planter	Rs.15000/Unit
	(e) Seed Drill	Rs.15000/Unit
	(f) Zero Till Multi Crop Planter	Rs.15000/Unit
	(g) Ridge Furrow Planter	Rs.15000/Unit
	(h) Chiseller	Rs.8000/Unit
	(i) Rotavator	Rs.35000/Unit
	(j) Laser Land Leveler	Rs.150000/Unit
	(k) Tractor mounted sprayer	Rs. 10000/Unit
	(i) Multi crop thresher	Rs. 40000/Unit
6	Efficient Water Application Tools:	
	(a) Sprinkler Sets	Rs.10000/-
	(b) Pump Sets	Rs.10000/Unit
	(c) Pipe for carrying water from source to the field	Rs. 15000 / per farmer /
		benificiary
	(d) Mobile Rain gun	Rs. 15000/Unit
7	Cropping System based trainings	Rs.3500/ Session Rs.14000/ Training

^{*}Subsidy for SC/ST small, marginal and women farmers 50% of cost and other farmers 40% of cost in farm implements & equipments.

District Covered (Identified) under NFSM - Pulses

SI.No.	States	Total Number of Districts in states	Total Number of Districts covered under - NFSM
1	Andhra Pradesh	13	13
2	Arunachal Pradesh	17	17
3	Assam	27	27
4	Bihar	38	38
5	Chhattisgarh	27	27
6	Goa*	2	2
7	Gujarat	26	26
8	Haryana	21	21
9	Himachal Pradesh	12	12
10	Jammu & Kashmir	22	22
11	Jharkhand	24	24
12	Karnataka	30	30
13	Kerala*	14	14
14	Madhya Pradesh	51	51
15	Maharashtra	35	33
16	Manipur	9	9
17	Meghalaya	11	11
18	Mizoram	8	8
19	Nagaland	11	11
20	Odisha	30	30
21	Punjab	22	22
22	Rajasthan	33	33
23	Sikkim	4	4
24	Tamil Nadu	32	30
25	Telangana	10	9
26	Tripura	8	8
27	Uttar Pradesh	75	75
28	Uttarakhand	13	13
29	West Bengal	19	18
	Total	644	638

^{*}Goa & Kerala States Information updated Ministry Letter No 1-8/2014-NFSM as on Dated:03/02/2016

Pulses: All India- Crop Calender: Package Of Practice

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ
Crop					
Chickpea					
	Punjab, Haryana,	East Uttar Pradesh,	M.P., Chhattisgarh,	A. P., Kerala,	J&K, H.P., Uttrakhand
	Delhi, Rajasthan,	Bihar, West Bengal,	Maharashtra, Gujarat and	Karnataka, Tamil	NEH States (Sikkim, Nagaland,
States	West Uttar Pradesh,	Jharkhand, Assam,	(Bundelkhand Region of	Nadu, Odisha	Meghalaya, Manipur, Mizoram,
	Plains of Uttrakhand	J&K	UP)		Tripura, Arunachal Pradesh)
Sowing time					
Rainfed Areas	1 st fortnight of Oct.	1 st fortnight of Oct2 nd fortnight of			
					Oct.
Irrigated Areas	Last week of Oct 1 st	Last week of Oct 1 st	Last week of Oct 1 st	2 nd fortnight of Nov	1 st fortnight of Nov 1 st fortnight
	week of Nov.	week of Nov.	week of Nov.	1 st fortnight of Dec.	of Dec.
Late sown	1 st fortnight of Dec	1 st fortnight of Dec	1 st fortnight of Dec.	2 nd fortnight of Dec.	2 nd fortnight of Dec.
Areas	2 nd fortnight of Dec.	2 nd fortnight of Dec.			
Seed Rate					
Small size	60-70 kg/ha	70-80 kg/ha	70-80 kg/ha	70-80 kg/ha	70-80 kg/ha
Medium size	80-90 kg/ha	80-90 kg/ha	80-90 kg/ha	80-90 kg/ha	80-90kg/ha
Bold size	100- 120 kg/ha	90-100 kg/ha	100- 120 kg/ha	90- 100 kg/ha	100- 120 kg/ha
Spacing					
Timely Sown	30 X 10 cm				
Late Sown	25 X10 cm				
Irrigated	45 X 10 cm	45 X 10 cm	45X 10 cm	45 X 10 cm	45 X 10 cm
Seed Treatmen	nt				
Fungicide	2 gm Thiram + 1 gm	2 gm Thiram + 1 gm Carbendazim			
	Carbendazim or	Carbendazim or	Carbendazim or	Carbendazim or	or Carboxin (Vitavax) 2 gm/kg of
	Carboxin (Vitavax) 2	Carboxin (Vitavax) 2	Carboxin (Vitavax) 2	Carboxin (Vitavax) 2	Seeds
	gm/kg of Seeds	gm/kg of Seeds	gm/kg of Seeds	gm/kg of Seeds	
Insecticide	Thiamethoxam 70	Thiamethoxam 70 W.P.	Thiamethoxam 70 W.P.	Thiamethoxam 70	Thiamethoxam 70 W.P. @ 3 gm/kg
	W.P. @ 3 gm/kg Seed	@ 3 gm/kg Seed	@ 3 gm/kg Seed	W.P. @ 3 gm/kg Seed	Seed
Rhizobium	Rhizobium 5 gm +	Rhizobium 5 gm + PSB	Rhizobium 5 gm + PSB 5	Rhizobium 5 gm +	Rhizobium 5 gm + PSB 5 gm/kg
	PSB 5 gm/kg	5 gm/kg	gm/kg	PSB 5 gm/ kg	

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ
Crop	1,,,,,,,	1,222	02	~2	1,222
	oil test based applied as Basal	(1)		1	
Macronutrients	N:P:K:S	N:P:K:S	N:P:K:S	N:P:S	N:P:K:S
	18-20:40-60:20:20	15-20:40-60: 20 : 20	20:40:20:20	15:30-40:20	20-25 :40:20:20
Micronutrients	ZnSo ₄ @ 25 kg /ha,	ZnSo ₄ @ 25 kg /ha,	ZnSo ₄ @ 25 kg /ha,	ZnSo ₄ @ 25 kg /ha,	ZnSo ₄ @ 25 kg /ha,
	Mo (Na-Molybdate) @	Mo (Na-Molybdate)	Mo (Na-Molybdate) @	Mo (Na-Molybdate) @	Mo (Na-Molybdate) @
	3.5g for seed treatment	@ 3.5g for seed treatment	3.5g for seed treatment	3.5g for seed treatment	3.5g for seed treatment
Foliar Spray(Need	Urea @ 2% at 45-55 DAS,	Urea @ 2% at 45-55 DAS,	Urea @ 2% at 45-55	Urea @ 2% at 45-55	Urea @ 2% at 45-55
based)	DAP @ 2%	DAP @ 2%	DAS, DAP @ 2%	DAS, DAP @ 2%	DAS, DAP @ 2%
	Boron @ 0.2% at	Boron @ 0.2% at flowering	Boron @ 0.2% at	Boron @ 0.2% at	Boron @ 0.2% at
	flowering (50-60 DAS)	(50-60 DAS)	flowering (50-60 DAS)	flowering (50-60 DAS)	flowering (50-60 DAS)
Irrigation	Two irrigations 1 st at	Two irrigations 1 st at	Two irrigations 1 st at	One irrigation at pre	One irrigation at pod
	Branching (40 -50 DAS) &	Branching (40 -50 DAS) &	Branching (40 -50 DAS)	flowering (45-55 DAS),	development stage
	2 nd at pod initiation (70-80	2 nd at pod initiation (70-80	& 2 nd at pod	and 2 nd at pod	(70-80 DAS)
	DAS)	DAS)	development (70-80	development stage (70-	
			DAS)	80 DAS)	
Weed Manageme	ent				
Manual	One hand weedings at 25-	One hand weedings at 25-	One hand weedings at	One hand weedings at	One hand weedings at 25-
	30 DAS	30 DAS	25- 30 DAS	25- 30 DAS	30 DAS
Chemical	Pendimethalin at PE stage	Pendimethalin at PE stage	Pendimethalin at PE	Pendimethalin at PE	Pendimethalin at PE
	@ 1-1.25 Kg a.i. /ha or	@ 1 ó 1.25 kg a.i. /ha	stage @ 1 ó 1.25 Kg	stage @ 1 ó 1.25 Kg	stage @ 1 ó 1.25 Kg
	Fluchloralin @ 0.75 kg a.i.		a.i./ha	a.i./ha	a.i./ha
	/ha.				
Maturity/ Harve	sting				
Rainfed	130-140 DAS	120-140 DAS	120-140 DAS	120-140 DAS	120-140 DAS Mid -
	Early March- Early April	Mid -March- April	Mid -Feb- Mid March	Mid- Jan-Last March	March- Early April
Irrigated	120-140 DAS	120-140 DAS	120-140 DAS	120-140 DAS	120-140 DAS
	Mid- March- Mid April	Mid -March- Mid April	Mid- March- Mid April	Early March-Last April	Mid -March- Mid April
Late Sown	120-130	120-130	125-135	120-130	120-130 DAS
	Last March-Last April	Last March-Last April	Last March-Last April	Last March-Last April	Last March-Last April
Cropping System	Chickpea+ Barley (4:2	Chickpea + Mustard (4:1)	Chickpea+ Linseed,	Chickpea + Safflower	Chickpea + Mustard (4:2
	row) Chickpea + Mustard		Chickpea + Safflower	(2:1) & Chickpea+	and 6:2) Chickpea +
	(4-6:1 row)		(4:1)	Coriander (4:1)	Wheat (2:2)

360 Pulses in India: Retrospect and Prospects -2018 DPD, Govt. of India, Bhopal\

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ
Crop					
Pigeonpea					
States	Punjab, Haryana, Delhi, Rajasthan, West Uttar Pradesh, Plains of Uttrakhand	East Uttar Pradesh, Bihar, West Bengal, Jharkhand, Assam, J&K	M.P., Chhattisgarh, Maharashtra, Gujarat and (Bundelkhand Region of UP)	A. P., Kerala, Karnataka, Tamil Nadu, Odisha	J& K, H.P., Uttrakhand NEH States (Sikkim, Nagaland, Meghalaya, Manipur, Mizoram, Tripura, Aru.Pradesh)
Sowing time					
Normal	2 nd fortnight of May - 1 st fortnight of June Summer: Mid-April-1 st week of May	Early: 1 st fortnight of June Late: 1 st fortnight of July Rabi: Mid Sep Mid Oct.	Rainfed: 1 st fortnight of July Irrigated: 2 nd fortnight of June Rabi: Mid Sep Mid Oct.	Kharif: Onset of Monsoon/ 2 nd fortnight of June Rabi: Mid Sep Mid Oct.	Early: Mid May-Mid July
Transplant (Bidar/Dharwad)	-	-	1 st fortnight of May- 1 st fortnight of June	1 st fortnight of May- 1 st fortnight of June	-
Seed Rate & Spa	cing				
Normal	18-20 kg/ha; 45X15 cm	Early: 18-20 kg/ha,45X15cm Late - 12-15kg/ha ,60X10cm Rabi: 25-30 kg/ha;30X10 cm		Rabi:25-30kg/ha;30X10cm	Early:18-20 kg/ha,45X15cm
Transplant (Bidar/Dharwad)			Seed Rate- 2-5 kg/ha Spacing: Irrigated: 5ftX 3ft Rainfed 6ftX 3 ft (MP- Shahdol, Rewa, Maharashtra- Vidarbha and some parts of MH)	Seed Rate- 2-5 kg/ha Spacing: Irrigated: 5ftX 3ft Rainfed -6ftX 3 ft Karnataka- Gulbarga, Bijapur, Bidar, Dharwad etc. and some parts of AP, TN and Telangana)	
Seed Treatment					
Fungicide	Thiram @ 2gm + Carbendazim @ 1gm or Thiram @3 gm or Trichoderma 5 gm / kg	Thiram @ 2gm + Carbendazim @ 1gm or Thiram @3 gm or Trichoderma 5 gm / kg	Thiram @ 2gm + Carbendazim @ 1gm or Thiram @3 gm or Trichoderma 5 gm / kg	Thiram @ 2gm + Carbendazim @ 1gm or Thiram @3 gm or Trichoderma 5 gm / kg	Thiram @ 2gm + Carbendazim @ 1gm or Thiram @3 gm or Trichoderma 5 gm / kg

Pulses in India : Retrospect and Prospects -2018 DPD , Govt. of India, Bhopal\

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ
Crop					
Insecticide	Thiamethoxam 70 W.P.	Thiamethoxam 70 W.P. @	Thiamethoxam 70 W.P. @	Thiamethoxam 70 W.P. @	Thiamethoxam 70 W.P. @
	@ 3 gm/Kg Seed	3 gm/Kg Seed	3 gm/Kg Seed	3 gm/Kg Seed	3 gm/Kg Seed
Rhizobium	10gm/kg	10gm/kg	10gm/kg	10gm/kg	10gm/kg
Fertilizer (kg/ha)	N:P:S: ZnSo4	N:P :S: ZnSo4	N : P : S	N: P: S	N:P:S: ZnSo4
(Soil test based as	15-20:40:20:25	15-20:40:20:25	15:40:20	15:30:20	15-20:40 : 20:25
Basal dose)					
Irrigation	One irrigation pod	Early ó Pre-monsoon	One irrigation pod	One irrigation pod	One irrigation pod
	development stage if	irrigation as per	development stage if	development stage if	development stage if
	required (100-110 DAS)	requirement	required	required (100-110 DAS)	required (100-115 DAS)
		Rabi ó After 40 to 60 DAS	(100-110 DAS)		
		& 100- 110 days if required	at ad	at ad	
IC	-	-	1 st at 40-45 DAS & 2 nd at	1 st at 40-45 DAS & 2 nd at	-
Operation(Nipping			55-60 DAS	55-60 DAS	
)					
Weed Management					
Manual	Two hand weedings at 25	Two hand weeding at	Two hand weedings at 25	Two hand weedings	Two hand weeding at
	& 45 DAS	25 & 45 DAS	& 45 DAS	at 25 & 45 DAS	25 & 45 DAS
Chemical	Application of Pre-	Application of Pre-	Application of Pre-	Application of Pre-	Application of Pre-
	emergence Pendimethalin	emergence Pendimethalin	emergence Pendimethalin	emergence Pendimethalin	emergence Pendimethalin
	/Alachlor/Metachlor @	/Alachlor/Metachlor @	/Alachlor/Metachlor @	/Alachlor/Metachlor @	/Alachlor/Metachlor @
	1 ó 1.5 a.i. Kg/ha.	1 ó 1.5 a.i. Kg/ha.	1 ó 1.5 a.i. Kg/ha.	1 ó 1.5 a.i. Kg/ha.	1 ó 1.5 a.i. Kg/ha.
Maturity/Harvesting					
Kharif	190-200DAS OctDec.	190-200 DAS NovJan.	175-190 DAS OctJan.	150-170 DAS SepOct.	200-230 DAS OctDec.
Rabi	-	240-260 DAS Jan Feb.	200-230 DAS Jan Feb.	200-230 DAS Jan Feb.	-
Summer	190-200 DAS OctNov.	-	-	-	-
Dharwad	-	-	180-200 DAS May-June	180-200 DAS May-June	-
Cropping System	Pigeonpea-wheat	Early - Pigeonpea- wheat	Pigeonpea+ Groundnut (4	Pigeonpea+ Sorghum (2:1)	Pigeonpea:Sorghum (1:1)
	sequential cropping	Late óPigeonpea +Sorghum	:2)Pigeonpea + Soybean	Pigeonpea + Groundnut	Pigeonpea:Pearl millet
	Intercropping with	Moong/Urd/Sesame by	(4: 2) Pigeonpea +	(4: 2)	(1:1) Pigeonpea+
	Urdbean or Moongbean	pairing pigeonpea row at	Sorghum (2:1)	Pigeonpea+Mung/Urdbean	Cauliflower/Capsicum
	(1:1 row)	40/80 cm & planting one	Pigeonpea +Urdbean(1:1)	/ Cowpea (1:1)	
		row of intercrop			

Pulses in India: Retrospect and Prospects -2018 DPD , Govt. of India, Bhopal\

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ
Crop					
Mung/Urd					
States	Punjab, Haryana, Delhi, Rajasthan, West Uttar Pradesh, Plains of Uttrakhand	East Uttar Pradesh, Bihar, West Bengal, Jharkhand, Assam, J&K	M.P., Chhattisgarh, Maharashtra, Gujarat and (Bundelkhand Region of UP)	A. P., Kerala, Karnataka, Tamil Nadu, Odisha	J& K, H.P., Uttrakhand NEH States (Sikkim, Nagaland, Meghalaya, Manipur, Mizoram, Tripura, Aru.Pradesh)
Sowing Time					
Kharif	1 st fortnight of July/ Onset of monsoon	1 st fortnight of July/ Onset of monsoon	2 nd fortnight of June	2 nd fortnight of June	1 st week of July to 1 st week of Aug.
Rabi	-	-	2 nd fortnight of Oct 2 nd fortnight of Nov.	2 nd fortnight of Oct 2 nd fortnight of Nov.	-
Spring/ Summer	2 nd fortnight of March - 1 st week of April	2 nd fortnight of March -1 st week of April	2 nd fortnight of March -1 st week of April	Summer: 2 nd fortnight of March -1 st week of April Spring: 1 st fortnight of Dec.	-
Seed rate & Spacin	g				
Kharif	15-20 kg/ha; 45X10 cm	15-20 kg/ha;45X10 cm	15-20 kg/ha;45X10 cm	15-20 kg/ha;45X10 cm	15-20 kg/ha;45X10 cm
Rabi	-	-	25-30 kg/ha;30X10 cm,	25-30 kg/ha30X10 cm,	-
Spring/ Summer	30-35 kg/ha , 25X5 cm 20-25Kg/ ha ,45X10 cm	30-35 kg/ha , 25X5 cm 20-25Kg/ ha , 45X10 cm	30-35 kg/ha , 25X5 cm 20-25Kg/ ha , 45X10 cm	20-25 kg/ha; 30X10 cm	-
Seed Treatment					
Fungicide	Thiram @ 2gm + Carbendazim @ 1gm or Thiram @ 3 gm or Trichoderma 5 gm / kg	Thiram @ 2gm + Carbendazim @ 1gm or Thiram @ 3 gm or Trichoderma 5 gm / kg	Thiram @ 2gm + Carbendazim @ 1gm or Thiram @ 3 gm or Trichoderma 5 gm / kg	Thiram @ 2gm + Carbendazim @ 1gm or Thiram @ 3 gm or Trichoderma 5 gm / kg	Thiram @ 2gm + Carbendazim @ 1gm or Thiram @ 3 gm or Trichoderma 5 gm / kg
Insecticide	Thiamethoxam 70 W.P. @ 3 gm/Kg Seed	Thiamethoxam 70 W.P. @ 3 gm/Kg Seed	Thiamethoxam 70 W.P. @ 3 gm/Kg Seed	Thiamethoxam 70 W.P. @ 3 gm/Kg Seed	Thiamethoxam 70 W.P. @ 3 gm/Kg Seed
Rhizobium	Rhizobium@5gm/kg seed	Rhizobium @5gm/kg seed	Rhizobium @5gm/kg seed	Rhizobium@5gm/kg seed	Rhizobium@5gm/kg seed

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ			
Crop								
Fertilizer (kg/ha	Fertilizer (kg/ha) (Soil test based applied as basal dose)							
Kharif	N:P:K:S	N:P:K:S	N:P:K:S	N:P:K:S	N:P:K:S			
	10:40:20:20 as (Basal)	10:40:20:20 as (Basal)	10:30-40: 20 :20 as (Basal)	15-20:30-40 : 20 :20 as	10:40:20:20 as (Basal)			
	Mo (Na-Molybdate) @	Mo (Na- Molybdate) @	Mo (Na-Molybdate) @ 3.5g					
	3.5g for seed treatment	3.5g for seed treatment	for seed treatment	@ 3.5g for seed treatment	(Na-Molybdate) @ 3.5g for seed treatment			
Rabi	-	-	N:P:K:S	N:P:K:S	-			
			20-25:30-40:20:20	20-25:30-40:20:20				
Spring/Summer	N:P:K:S- 10:30:20:20	N:P:K:S- 20:30:20:20	N:P:K:S- 20:30:20:20	N:P:K:S	-			
•				20-25:30-40:20:20				
Irrigation								
Rabi	-	-	As per requirement of crop	As per requirement of crop	-			
			in absence of rain	in absence of rain				
(Spring/Summer)	1 st at 25 DAS subsequent	1 st at 25 DAS subsequent	1 st at 25 DAS subsequent	1 st at 25 DAS subsequent	-			
	as per requirement	as per requirement	as per requirement	as per requirement				
Weed Managme	nt							
Manual	One hand weeding at 30	One hand weeding at 30	One hand weeding at 30	One hand weeding at 30	One hand weeding at 30			
	DAS	DAS	DAS	DAS	DAS			
Chemical	Pre-emergence	Pre-emergence application	Pre-emergence application	Pre-emergence application	Pre-emergence application			
	application of	of Pendimethalin @ 0.75-						
	Pendimethalin @ 0.75-	1Kg/ha.	1Kg/ha.	1Kg/ha.	1Kg/ha.			
	1Kg/ha.				-			
Maturity/Harves	sting							
Kharif	70-90 DAS; SepOct.	70-90 DAS; SepOct.	75-85; SepOct.	70-90 DAS; SepOct.	65-85 DAS; SepOct.			
Rabi	-	-	75-85 DAS; Jan-Feb	75-85 DAS; Jan-Feb	-			
(Spring/Summer)	70-90 DAS; May-June	70-90 DAS; May-June	70-90 DAS; May-June	70-90 DAS; May-June	-			
	_	·		70-90 DAS; MarApril				
Cropping System	Intercropping of Moong	Intercropping of Moong	Intercropping of Moong	Rice ó Rice-Greengram/	Intercropping of Moong			
	bean /Urdbean with	bean /Urdbean with	bean /Urdbean with	Blackgram	bean /Urdbean with			
	summer planted ó	summer planted ó	summer planted Sunflower	Mungbean/Urdbean+	summer planted ó			
	sugarcane (2:1) and	sugarcane (2:1) and	(6:2 row ratio)	Sugarcane(2:1)	sugarcane (2:1) and			
	Sunflower (6:2 row ratio)	Sunflower (6:2 row ratio)			Sunflower (6:2 row ratio)			
					Moong/Urd: Capsicum			

Pulses in India: Retrospect and Prospects -2018 DPD , Govt. of India, Bhopal\

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ
Crop					
					Lentil Continued
Maturity/	120-140 DAS	135-140 DAS	130-140 DAS	-	120-140 DAS
Harvesting	March-April	March-April	March-April		March-April
Cropping System	Inter cropping with	Rice- lentil sequential	Rice-lentil Utera cropping		Inter cropping with barely,
	barely, rape / mustard	cropping in Northern	in Chhattisgarh.	-	rape & mustard (2:1)
	(2:2) Inter cropping with	Bihar	Intercrop with Linseed,		Lentil + Vegetable crops
	autumn sugarcane (2:1).		Barley and Mustard		intercropping
Fieldpea					
States	Punjab, Haryana, Delhi,	East Uttar Pradesh, Bihar,	M.P., Chhattisgarh,	A. P., Kerala, Karnataka,	J& K, H.P., Uttrakhand
	Rajasthan, West Uttar	West Bengal, Jharkhand,	Maharashtra, Gujarat and	Tamil Nadu, Odisha	NEH States (Sikkim,
	Pradesh, Plains of	Assam, J&K	(Bundelkhand Region of		Nagaland, Meghalaya,
	Uttrakhand		UP)		Manipur, Mizoram,
					Tripura, Aru.Pradesh)
Sowing time	Third week of Oct. ó Ist	Third week of Oct. ó Ist	Third week of Oct. ó Ist	2 nd week of Oct- 2 nd week	Third week of Oct. ó Ist
	week of Nov.	week of Nov.	week of Nov.	of Dec	week of Nov
Seed rate & spac					
Tall	60-70 kg; 30x10 cm	60-70 kg; 30x10 cm	60-70 kg; 30x10 cm	60-70 kg; 30x10 cm	60-70 kg; 30x10 cm
Dwarf	80-100kg; 22X10 cm	80-100kg:22X10 cm	80-100kg; 22X10 cm	80 -90 kg/ha; 22X10 cm	80-100kg; 22X10 cm
Seed Treatment					
Fungicide	Thiram @ 2gm +	Thiram @ 2gm +	Thiram @ 2gm +	Thiram @ 2gm +	Thiram @ 2gm +
	Carbendazim @ 1gm or	Carbendazim @ 1gm or	Carbendazim @ 1gm or	Carbendazim @ 1gm or	Carbendazim @ 1gm or
	Thiram @3 gm or	Thiram @3 gm or	Thiram @3 gm or	Thiram @3 gm or	Thiram @3 gm or
	Trichoderma 5 gm / kg	Trichoderma 5 gm / kg	Trichoderma 5 gm / kg	Trichoderma 5 gm / kg	Trichoderma 5 gm / kg
Insecticide	Chlorpyriphos 20 EC @8	Chlorpyriphos 20 EC @8	Chlorpyriphos 20 EC @8	Chlorpyriphos 20 EC @8	Chlorpyriphos 20 EC @8
	ml/Kg of seed	ml/Kg of seed	ml/Kg of seed	ml/Kg of seed	ml/Kg of seed
Rhizobium	Rhizobium culture	Rhizobium culture	Rhizobium culture	Rhizobium culture	Rhizobium culture
	10gm/Kg	10gm/Kg	10gm/Kg	10gm/Kg	10gm/Kg
Fert. Dose (kg/ha)	N:P:K:S	N:P:K:S	N:P:K:S	N:P:K:S 20-40 : 60:20:20	N:P:K:S
(Soil test based	20-40:60:20:20	20-40:60:20:20	20-40 : 60:20:20		20-40:60:20:20
applied as Basal)					

Pulses in India : Retrospect and Prospects -2018 DPD , Govt. of India, Bhopal\

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ
Crop					
Weed management					
Manual	One hand weeding at 30	One hand weeding at 30	One hand weeding at 30	One hand weeding at 30	One hand weeding at 30
	DAS	DAS	DAS	DAS	DAS
Chemical	Pendimethalin @1 kg a.i.	Pendimethalin @1 kg a.i.	Pendimethalin @1 kg a.i.	Pendimethalin @1 kg a.i.	Pendimethalin @1 kg a.i.
	as pre-emergence	as pre-emergence	as pre-emergence	as pre-emergence	as pre-emergence
Maturity/	120-140 DAS	125-135 DAS	130-150 DAS	120-140 DAS	120-140 DAS
Harvesting	March-April	March-April	Mid Feb- Mid- March	Feb- March	Feb- March
Cropping System	Sequential cropping after	Wheat-Pea	Sorghum/soybean/Pearl	-	Sequential cropping after
	rice, maize or pearl millet		millet ó pea		rice, maize or pearl millet
					Field pea- Vegetable crops
Lathyrus					
Sowing time					
Rabi	-	Last week Oct. to early	Last week Oct. to early	-	Last week Oct. to early
		Nov.	Nov.		Nov.
Utera	-	-	Last week of Sep to First	-	-
			week to Oct		
Seed Rate /Spaci	ng				
Line Sowing	-	40-60 kg/ha	40-60 kg/ha	-	40-60 kg/ha
		(30X10 cm)	(30X10 cm)		(30X10 cm)
Utera(Broadcast)	-	70-80 Kg/ha	70-80 Kg/ha	-	70-80 Kg/ha
Seed Treatment					
Fungicide	-	Thiram @ 2gm +	Thiram @ 2gm +	-	Thiram @ 2gm +
		Carbendazim @ 1gm or	Carbendazim @ 1gm or		Carbendazim @ 1gm or
		Thiram @3 gm or	Thiram @3 gm or		Thiram @3 gm or
		Trichoderma 5 gm / kg	Trichoderma 5 gm / kg		Trichoderma 5 gm / kg
Insecticide	-	Chlorpyriphos 20 EC @8	Chlorpyriphos 20 EC @8	-	Chlorpyriphos 20 EC @8
		ml/kg of seed	ml/kg of seed		ml/kg of seed
Rhizobium	-	Rhizobium culture	Rhizobium culture	-	Rhizobium culture
		10gm/kg.	10gm/kg.		10gm/kg.

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ
Crop					
					Lathyrus Continued
Fertilizer (kg/ha) (Soil test based applied as basal)	ot based 0:40-6		N:P:K:S 0:40-60:0:0	-	N:P:K:S 0:40-60:0:0
Irrigation		One irrigation (60-70 DAS)	One irrigation (60-70 DAS)	-	One irrigation (60-70 DAS)
Weed Manageme	ent				
Manual (if soil condition permits)	-	one hand weedings at 30 ó 35 DAS	one hand weedings at 30 - 35 DAS	-	one hand weedings at 30 - 35 DAS
Chemical		Application of Pendimethalin as pre- emergence stage @ 1 \u00f3 1.5 kg a.i./ha	Application of Pendimethalin as pre- emergence stage @ 1 \u00f3 1.5 kg a.i./ha	-	Application of Pendimethalin as pre- emergence stage @ 1 \u00f3 1.5 kg a.i./ha.
Maturity/ Harvesting	-	110-120 DAS	95-100 DAS	-	110-120 DAS
Cropping System	-	-	Can be grown with Rice as relay/ Utera	-	-
Horsegram					
States	Punjab, Haryana, Delhi, Rajasthan, West Uttar Pradesh, Plains of Uttrakhand	East Uttar Pradesh, Bihar, West Bengal, Jharkhand, Assam, J&K	M.P., Chhattisgarh, Maharashtra, Gujarat and (Bundelkhand Region of UP)	A. P., Kerala, Karnataka, Tamil Nadu, Odisha	J& K, H.P., Uttrakhand NEH States (Sikkim, Nagaland, Meghalaya, Manipur, Mizoram, Tripura, Aru.Pradesh)
Sowing Time					
Kharif	Last June - last July	-	-	-	Last June ó last July
Rabi	-	Last Oct early Nov.	I st fortnight of Sepearly Oct.	I st fortnight of Sepearly Oct.	-
Seed Rate /Spaci	ng				
Line Sowing	22-30 Kg/ha 40-45X25-30 cm	22-30 Kg/ha 40-45X25-30 cm	22-30 Kg/ha 40-45X25-30 cm	22-30 Kg/ha 40-45X25-30 cm	22-30 Kg/ha 40-45X25-30 cm
Broadcast	40-50 kg/ha	40-50 kg/ha	40-50 kg/ha	40-50 kg/ha	40-50 kg/ha

Pulses in India: Retrospect and Prospects -2018 DPD , Govt. of India, Bhopal\

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ	
Crop						
Seed Treatment						
Fungicide	Bavistin@ 2g/kg or	Bavistin@ 2g/kg or	Bavistin@ 2g/kg or	Bavistin@ 2g/kg or	Bavistin@ 2g/kg or	
	Trichoderma @ 4g/kg	Trichoderma @ 4g/kg	Trichoderma @ 4g/kg of	Trichoderma @ 4g/kg	Trichoderma @ 4g/kg of	
	of Seed	of Seed	Seed	of Seed	Seed	
Insecticide	Chlorpyriphos 20 EC	Chlorpyriphos 20 EC	Chlorpyriphos 20 EC @	Chlorpyriphos 20 EC@	Chlorpyriphos 20EC @ 8	
	@ 8 ml/kg of seed	@8 ml/kg of seed	8 ml/kg of seed	8 ml/kg seed	ml/kg of seed	
Rhizobium	Rhizobium culture	Rhizobium culture	Rhizobium culture	Rhizobium culture	Rhizobium culture	
	10gm/Kg.	10gm/Kg.	10gm/Kg.	10gm/Kg.	10gm/Kg.	
Fertilizer (kg/ha)	N:P:K:S	N:P:K:S	N:P:K:S	N:P:K:S	N:P:K:S	
(Soil test based applied as	10:20:0:0	10:20:0:0 10:20:0:0		10:20:0:0	10:20:0:0	
basal)						
Irrigation	Grown as rainfed	Grown as rainfed	Grown as rainfed	Grown as rainfed	Grown as rainfed	
Weed Management						
Manual	one hand weedings at	one hand weedings at	one hand weedings at	one hand weedings at	one hand weedings at 20-25	
	20-25 DAS	20-25 DAS	20-25 DAS	20-25 DAS	DAS	
Chemical	Application of	Application of	Application of	Application of	Application of Fluchloralin	
	Fluchloralin at pre-	Fluchloralin at pre-	Fluchloralin at pre-	Fluchloralin at pre-	at pre-emergence stage @ 2	
	emergence stage @ 2	emergence stage @ 2	emergence stage @ 2	emergence stage @ 2	ml/l of water	
	ml/l of water	ml/l of water	ml/l of water	ml/l of water		
Maturity/ Harvesting	80-100 DAS	80-100 DAS	80-100 DAS	80-100 DAS	80-100 DAS	
_	SepOct.	JanFeb.	DecJan.	DecJan.	SepOct.	

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ	
Crop						
Ricebean						
States	Punjab, Haryana, Delhi, Rajasthan, West Uttar Pradesh, Plains of Uttrakhand	East Uttar Pradesh, Bihar, West Bengal, Jharkhand, Assam, J&K	M.P., Chhattisgarh, Maharashtra, Gujarat and (Bundelkhand Region of UP) A. P., Kerala, Karnataka, Tan Nadu, Odisha		J& K, H.P., Uttrakhand NEH States (Sikkim, Nagaland, Meghalaya, Manipur, Mizoram, Tripura, Aru.Pradesh)	
Sowing time						
Kharif (Grain Purpose)	2 nd fortnight of August	-	-	-	2 nd fortnight of August	
Rabi (Fodder Purpose)	Upto September	-	-	-	Upto September	
Seed Rate						
Grain Purpose	40-50 kg/ha; 45-60X5- 10cm	-	-	-	40-50 kg/ha 45-60X510cm	
Fodder	60-75 kg/ha; 45-60X5- 10cm	-	-	-	60-75 kg/ha; 45-60X5-10cm	
Seed Treatment						
Fungicide	Bavistin@ 2g/kg or Trichoderma @ 4g/kg of Seed	-	-	-	Bavistin@ 2g/kg or Trichoderma @ 4g/kg of Seed	
Insecticide	Chlorpyriphos 20 EC @8 ml/Kg of seed	-		-	Chlorpyriphos 20 EC @8 ml/Kg of seed	
Rhizobium	Rhizobium culture 10gm/Kg.	-	-	-	Rhizobium culture 10gm/Kg.	
Fertilizer(kg/ha) (Soil test based applied as basal)	Grown on residual soil fertility.	-	-	-	Grown on residual soil fertility.	
Irrigation	Grown as rainfed	-	-	-	Grown as rainfed	
Weed Manag. Manual	One hoeing @ 30 DAS is enough	-	-	-	One hoeing @ 30 DAS is enough	
Maturity/ Harvesting	Kharif: 120-130 DAS Dec-Jan; Rabi: Jan-Feb	-	-	-	Kharif: 120-130 DAS Dec-Jan; Rabi: Jan-Feb	
Cropping System	-	-	-	-	Mixed with Jute, Maize, Finger millets	

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ
Crop					
Mothbean					
	Punjab, Haryana, Delhi,	East Uttar Pradesh,	M.P., Chhattisgarh,	A. P., Kerala,	J& K, H.P., Uttrakhand
	Rajasthan, West Uttar	Bihar, West Bengal,	Maharashtra, Gujarat	Karnataka, Tamil	NEH States (Sikkim, Nagaland,
States	Pradesh, Plains of	Jharkhand, Assam,	and (Bundelkhand	Nadu, Odisha	Meghalaya, Manipur, Mizoram,
	Uttrakhand	J&K	Region of UP)		Tripura, Aru.Pradesh)
Sowing time	2 nd fortnight of July	-	2 nd fortnight of July	-	-
Seed Rate	10-15 kg/ha	_	10-15 kg/ha	_	_
	30-60-40X15cm	_	30-60-40X15cm		_
Seed Treatment					
Fungicide	Bavistin@ 2g/kg or	-	Bavistin@ 2g/kg or	-	-
	Trichoderma @ 4g/Kg		Trichoderma @ 4g/Kg		
	of Seed		of Seed		
Insecticide	Chlorpyriphos 20 EC	-	Chlorpyriphos 20 EC	-	-
	@8 ml/Kg of seed		@8 ml./Kg of seed		
Rhizobium	Rhizobium culture	-	Rhizobium culture	-	-
	10gm/Kg.		10gm/Kg.		
Fertilizer (kg/ha)	N:P:K	-	N:P:K	-	-
	10:40:0		10:40:0		
Irrigation	As per requirement	-	As per requirement	-	-
Weed Managem	ient				
Manual	One hoeing at 30 DAS	-	One hoeing at 30 DAS	=	-
Chemical	Pre Plant incorporation	-	Pre Plant incorporation	=	-
	of fluchloralin (Basalin)		of fluchloralin (Basalin)		
	@ 0.5 to 1 kg a.i./ha		@ 0.5 to 1 kg a.i./ha		
Maturity/	120-130 DAS	-	120-130 DAS	-	-
Harvesting	NovDec.		NovDec.		
Cropping System	Mothbean+Pearlmillet	-	Mothbean+Pearlmillet	-	-
	(2:1)		(2:1)		

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ
Crop					
Rajmash					
	Punjab, Haryana, Delhi,	East Uttar Pradesh,	M.P., Chhattisgarh,	A. P., Kerala,	J& K, H.P., Uttrakhand
	Rajasthan, West Uttar	Bihar, West Bengal,	Maharashtra, Gujarat	Karnataka, Tamil	NEH States (Sikkim, Nagaland,
States	Pradesh, Plains of	Jharkhand, Assam,	and (Bundelkhand	Nadu, Odisha	Meghalaya, Manipur, Mizoram,
	Uttrakhand	J&K	Region of UP)		Tripura, Aru.Pradesh)
Sowing time					
Kharif	-	-	-	-	last week June to first week of
					July
Rabi	2 nd fortnight of	2 nd fortnight of	2 nd fortnight of October	2 nd fortnight of	-
	October	October		October	
Spring	-	-	-	-	Spring (Lower hills) 2 nd fortnight
					of March
Seed Rate	·	1		<u>'</u>	
Kharif					100-125 kg/ha
	_	-	_	-	45-50X8-10 cm
Rabi/Spring	100-125 kg/ha	100-125 kg/ha	100-125 kg/ha	100-125 kg/ha	100-125 kg/ha
	Rainfed:40cmx10 cm	Rainfed: 40cmx10cm	Rainfed: 40 cmx10 cm	Rainfed: 40 cmx10cm	Rainfed: 40 cmx 10cm
	Irrigated: 30cmx10cm	Irrigated:30cm x10cm	Irrigated: 30cmx 10cm	Irrigated:30cmx10cm	Irrigated: 30 cm x 10cm
Seed Treatment					
Fungicide	Bavistin@ 2g/kg or	Bavistin@ 2g/kg or	Bavistin@ 2g/kg or	Bavistin@ 2g/kg or	Bavistin@ 2g/kg or Trichoderma
	Trichoderma @ 4g/Kg	Trichoderma @ 4g/Kg	Trichoderma @ 4g/Kg	Trichoderma @ 4g/Kg	@ 4g/Kg of Seed
	of Seed	of Seed	of Seed	of Seed	
Insecticide	Chlorpyriphos 20 EC	Chlorpyriphos 20 EC	Chlorpyriphos 20 EC @	Chlorpyriphos 20 EC	Chlorpyriphos 20 EC @ 8 ml/Kg
	@ 8 ml/Kg of seed	@ 8 ml/Kg of seed	8 ml/Kg of seed	@ 8 ml/Kg of seed	of seed
Rhizobium	Rhizobium culture	Rhizobium culture	Rhizobium culture	Rhizobium culture	Rhizobium culture 10gm/Kg.
	10gm/Kg.	10gm/Kg.	10gm/Kg.	10gm/Kg.	
Fertilizer(kg/ha)	N:P:K	N:P:K	N:P:K	N:P:K	N:P:K
(Soil test based	90-120:60-80:0	90-120:60-80:0	90-120:60-80:0	90-120:60-80:0	90-120:60-80:0
applied as basal) Irrigation	25 DAS followed by	25 DAS followed by	25 DAS followed by	25 DAS followed by	25 DAS followed by irrigation at
TITIGATION	irrigation at 75 DAS.	irrigation at 75 DAS.	irrigation at 75 DAS.	irrigation at 75 DAS.	75 DAS.
	2 to 3 irrigation as per	2 to 3 irrigation as per	3 to 4 irrigation as per	3 to 4 irrigation as per	2 to 3 irrigation as per
	requirement	requirement	requirement	requirement	requirement
	requirement	requirement	requirement	requirement	requirement

Pulses in India: Retrospect and Prospects -2018 DPD, Govt. of India, Bhopal\

Zone/States	NWPZ	NEPZ	CZ SZ		NHZ
Crop					Rajmash Continued
Weed Management	t				-
Manual	One hand weeding at 30-35 DAS	One hand weeding at 30-35 DAS	One hand weeding at 30- 35 DAS	One hand weeding at 30-35 DAS	One hand weeding at 30-35 DAS
Chemical	Pendimethalin @ 1 to 0.75-1 kg a.i./ha in 500- 600 liters as pre- emergence	Pendimethalin @ 1 to 0.75-1 kg a.i./ha in 500- 600 liters as pre- emergence	Pendimethalin @ 1 to 0.75-1 kg a.i./ha in 500- 600 liters as pre- emergence	Pendimethalin @ 1 to 0.75-1 kg a.i./ha in 500-600 liters as pre- emergence	Pendimethalin @ 1 to 0.75-1 kg a.i./ha in 500-600 liters as pre- emergence
Maturity/ Harvesti		<u>υ</u>	<u> </u>	<u> </u>	
Kharif	-	-	-	-	80-120 DAS NovDec.
Rabi	80-120 DAS FebMarch	80-120 DAS FebMarch	80-120 DAS FebMarch	80-120 DAS FebMarch	-
Spring	-	-	-	-	80-120 DAS AugSep.
Cropping System	-	-	-	-	Early Potato-Rajmash (2:2 or 2:3)

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ
Crop					
Cowpea					
States	Punjab, Haryana, Delhi, Rajasthan, West Uttar Pradesh, Plains of Uttrakhand	East Uttar Pradesh, Bihar, West Bengal, Jharkhand, Assam, J&K	M.P., Chhattisgarh, Maharashtra, Gujarat and (Bundelkhand Region of UP)	A. P., Kerala, Karnataka, Tamil Nadu, Odisha	J& K, H.P., Uttrakhand NEH States (Sikkim, Nagaland, Meghalaya, Manipur, Mizoram, Tripura, Aru.Pradesh)
Sowing time					
Kharif	Early June - End of July	Early June - End of July	Early June - End of July	-	-
Rabi	-	-	-	1 st fortnight of Oct1 st fortnight of Nov.	-
Summer	-	-	-	-	Grain Purpose: 2nd to 4 th week of March Fodder purpose: February Green Manuring: April- May
Seed Rate					
Pure crop	20-25 Kg/ha	20-25 Kg/ha	20-25 Kg/ha	20-25 Kg/ha	20-25 Kg/ha
Fodder/Green Manure	30-35 Kg/ha	30-35 Kg/ha	30-35 Kg/ha	30-35 Kg/ha	30-35 Kg/ha
Spacing					
Bushy	35X10 cm	35X10 cm	35X10 cm	35X10 cm	35X10 cm
Spreading	45X15 cm	45X15 cm	45X15 cm	45X15 cm	45X15 cm
Seed Treatment					
Fungicide	Bavistin@ 2g/kg or Trichoderma @ 4g/Kg of Seed	Bavistin@ 2g/kg or Trichoderma @ 4g/Kg of Seed	Bavistin@ 2g/kg or Trichoderma @ 4g/Kg of Seed	Bavistin@ 2g/kg or Trichoderma @ 4g/Kg of Seed	Bavistin@ 2g/kg or Trichoderma @ 4g/Kg of Seed
Insecticide	Chlorpyriphos 20 EC @ 8 ml/Kg of seed	Chlorpyriphos 20 EC @ 8 ml/Kg of seed	Chlorpyriphos 20 EC @ 8 ml/Kg of seed	Chlorpyriphos 20 EC @ 8 ml/Kg of seed	Chlorpyriphos 20 EC @ 8 ml/Kg of seed
Rhizobium	Rhizobium culture 10gm/Kg.	Rhizobium culture 10gm/Kg.	Rhizobium culture 10gm/Kg.	Rhizobium culture 10gm/Kg.	Rhizobium culture 10gm/Kg.

Zone/States	NWPZ	NEPZ	CZ	SZ	NHZ	
Crop						
					Cowpea Continued	
Fertilizer (kg/ha)	N:P:K	N:P:K	N:P:K	N:P:K	N:P:K	
(Soil test based	15-20:50-60:10-20	15-20:50-60:10-20	15-20:50-60:10-20	15-20:50-60:10-20	15-20:50-60:10-20	
applied as basal)						
Irrigation	As per requirement	As per requirement	As per requirement	As per requirement	As per requirement	
Weed Management	-					
Manual	One hand weeding at	One hand weeding at	One hand weeding at 20-	One hand weeding at	One hand weeding at 20-30 DAS	
	20-30 DAS	20-30 DAS	30 DAS	20-30 DAS		
Chemical	Pendimethalin @ 1 to	Pendimethalin @ 1 to	Pendimethalin @ 1 to	Pendimethalin @ 1 to	Pendimethalin @ 1 to 0.75-1 kg	
	0.75-1 kg a.i./ha in 500-	0.75-1 kg a.i./ha in 500-	0.75-1 kg a.i./ha in 500-	0.75-1 kg a.i./ha in	a.i./ha in 500-600 liters as pre-	
	600 liters as pre-	600 liters as pre-	600 liters as pre-	500-600 liters as pre-	emergence	
	emergence	emergence	emergence	emergence		
Maturity/ Harvesti						
Kharif	90-100 DAS	145-150 DAS	65-85 DAS	-	-	
	OctNov.	OctNov.	SepOct.			
Rabi	-	-	-	80-120 DAS		
				DecJan.	-	
Summer	-	-	-	-	Grain Purpose: 85-105	
					DAS (May-June)	
					Fodder Purpose: 90-110	
					DAS (April-May)	
					Green Manure: 90-120 (Sep	
					Oct.)	
Cropping System	Sorghum+ Cowpea	-	Pigeonpea+ Cowpea	Maize+ Cowpea (1:1)	Sorghum+Cowpea (1:1)	
	(1:1)		(1:1)			

ANNEXURE- IV SPECIFIC STANDARDS PRESCRIBED FOR CERTIFICATION AT FIELD STAGE FOR PULSES

SI No	Crop	Minimum number of inspection	dist in r	ation tance neter	pla ear	type ants/ heads	oth F	parable er crop olant	Objectionable weed plant		affe by s bo dise	ant cted seed rne ases	Remarks
1	Black gram Bengal gram Horse gram	2	FS 10	5 5	FS 0.1	0.2 0.2	FS -	-	FS -	CS -	FS -	CS -	-
3.	Green gram Cowpea fresh bean	2	10	5	0.1	0.2	-	-	-	-	0.1	0.2	Disease for cowpea Aschochyta stem blight and Anthracnose Aschochyta blight & Cowpea mosaic for French bean bacterial blight Anthracnose, As chochyta blight & Cowpea mosaic for French bean bacterial blight & Cowpea mosaic step
4.	Moth bean	2	10	5	0.1	0.2	-	-	-	-	-	-	-
5.	Lentil	2	10	5	0.1	0.2	-	-	-	-	-	-	-
6.	Peas	3	10	5	0.1	0.2	-	-	-	-	-	-	-
7.	Pigeon pea	2	250	100	0.1	0.2	-	-	-	-	-	-	-
8	Rice bean	2	50	20	0.1	0.2	-	-	-	-	-	-	-

SEED STANDARD

Сгор	1	ure (Min)	ma	Innert tter(max)		er crop (max)		nl weed l(max)	Object -ble w seed (r	eed	Germin (mi		Mois ordir conta	ary	Maxii vapo pro conta	our of	disting var	ther guishable ieties DV)
	F	C	F	C	F	C	F	C	F	C	F	C	F	C	F	C	F	C
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
Blackgram/ green gram	98.0	98.0	2.0	2.0	5/ Kg	10/Kg	5/ Kg	10/ kg	-	-	75	75	9.0	9.0	08	08	10/ kg	20/ kg
Cowpea	98.0	98.0	2.0	2.0	none	10/Kg	none	10/ kg	-	-	75	75	9.0	9.0	08	08	5/ kg	10/ kg
French bean	98.0	98.0	2.0	2.0	none	none	none	10/ kg	-	-	75	75	9.0	9.0	07	07	5/ kg	10/ kg
Gram	98.0	98.0	2.0	2.0	none	5 /Kg	none	none	-	-	85	85	9.0	9.0	08	08	5/kg	10/ kg
Horsegram	98.0	98.0	2.0	2.0	none	10/Kg	none	none	-	-	80	80	9.0	9.0	08	08	5/kg	10/ kg
Lentil/Khesari	98.0	98.0	2.0	2.0	5/ kg	10/Kg	10/kg	20/ kg	-	-	75	75	9.0	9.0	08	08	10/ kg	20/ kg
Redgram	98.0	98.0	2.0	2.0	5/ kg	10/Kg	5/kg	10/ kg	-	-	75	75	9.0	9.0	08	08	10/ kg	20/ kg

F-Foundation Seed, C-Certified seed

Crop-wise and State-wise categories of district having LALP (Low Area Low Productivity), LAHP (Low Area High Productivity), HALP (High Area High Productivity), HAHP (High Area High Productivity) than its average area and productivity.

1. Total pulses

States	LALP	LAHP	HALP	НАНР
Andhra Pradesh	Ananthapuram, Srikakulam,	East Godavari, Visakhapatnam	Kadapa	Kurnool, Prakasham,
	Vizianagaram, West Godavari,			Guntur, Krishna
	Nellore, Chittoor			
Assam	Kamrup (Rural), Chirang, Golaghat,	Hailakandi, Cachar, Tinsukia,	Barpeta, Jorhat, Goalpara,	Sonitpur, Dhubri,
	Dibrugarh, Dima, Hasao, Karimganj,	Udalguri, Morigaon, Sivasagar,	Kokrajhar, Karbi,	lakhimpur, Darrang, Baksa,
	Kamrup (Metro)	Nalbari, Dhemaji	Anglong	Nagaon, Bongaigaon
Bihar	Bhabhua, Darbhanga, Bhagalpur,	Jamui, Rohtash, E.Champaran,	Aurangabad, Muzaffarpur,	Patna, Gaya, Jahanabad,
	Vaishali, Nawada, Khagaria,	Lakhisarai, W.Champaran,	Supaul, Nalanda,	Bhojpur
	Sitamarhi, Arwal, Saran, Sheohar	Araria, Banka, Kisangunj,	Madhubani, Saharsa,	
		Katihar, Buxer, Shekhpura,	Madhepura, Samastipur	
		Purnia, Begusarai, Siwan,		
		Mungher, Gopalgunj		
Chhattisgarh	Koriya, Surguja, Balrampur, Korba,	Bilaspur, Raipur, Durg, Janjgir	Raigarh, Balod,	Rajnandgaon, Bemetara,
	Surajpur, Kondagaon, Mahasamund,		Gariaband, Baloda bazaar,	Kabirdham, Mungeli
	Jagdalpur, Sukama, Narayanpur,		Kanker, Dhamtari,	
	Bijapur, Dantewada		Jashpur	
Gujrat	Arvalli, Ahmedabad, Valsad,	Dahod , Panchmahal, Surat ,	Kutch, Patan, Mehesana,	Bharuch, Vadodara ,
	Surendranagar, Amreli, Jamnagar,	Junagadh , Mahisagar, Dang,	Banaskantha, Narmada	Chhotaudepur,
	Rajkot , Bhavnagar, Gandhinagar,	Anand		Sabarkantha, Tapi
	Morbi, Gir Somnath, Kheda,			
	Porbandar , Botad , Devbhumi			
	Dwarka, Navsari			
J& K	Reasi	Kishtwar, Rajouri, Poonch	Jammu, Kathua, Samba,	Doda
			Udhampur	

States	LALP	LAHP	HALP	НАНР
Jharkhand	Sahebganj, West Singhbum, Pakud, Bokaro, Lohardaga, Jaamtara	Khunti, Devghar, Giridih, Koderma, Ramgarh, Dhanbad	Garhwa, Ranchi, Hazaaribagh, East	Simdega, Latehaar, Saraikela, Chatra, Palamu,
			Singhbhum, Dumka, Godda	Gumla
Karnataka	Mysore, Mandya, Tumkur, Chamarajanagar, Chitradurga, Ballari, Ramanagaram, Hassan, Chickballapur, Chikmagalur, Davanagere, Kolar, Haveri, Bengaluru ó urban, Udupi, Uttara kannada, Shimoga, Dakshina kannada, Kodagu	Bengaluru - rural	Gadag, Bagalkote, Belgavi, Yadgir, Koppal, Dharwad	Gulbarga, Vijayapur, Bidar, Raichur
Madhya Pradesh	Datia, Mandsaur, Dindori, Betul, Shajapur, Alirajpur, Singroli, Aagar, Mandla, Anuppur, Umaria, Barwani, Shahdol	Sehore, Hoshangabad, Chhindwara, Dewas, UjjainKatni, Indore, Sidhi, Khargone, Balaghat, Morena, Khandwa, Bhopal, Bhind, Gwalior, Neemuch, Harda, Jhabua, Sheopurkala, Burhanpur	Sagar, Vidisha, Chattarpur Ashoknagar, Rewa, Tikamgarh, Rajgarh, Shivpuri, Guna, Seoni, Dhar, Ratlam	Damoh, Satna, Narsinghpur, Raisen, Panna, Jabalpur
Maharashtra	Aurangabad, Chandrapur, Nasik, Satara, Sangli, Pune, Dhule, Nandurbar, Bhandara, Gondia, Gadchiroli, Kolhapur, Raigad, Palghar, Ratnagiri, Thane, Sindhudurg	Wardha	Osmanabad, Buldhana, Solapur, Beed, Ahmednagar, Parbhani, Jalgaon, Washim, Hingoli	Latur, Yavatmal, Amravati, Nanded, Jalna, Akola, Nagpur
Odisha	Puri, Jagatsinghpur, Jajpur, Sambalpur, Khordha, Koraput, Deogarh, Balasore, Nabarangpur, Jharsuguda, Malkangiri	Mayurbhanj, Rayagada, Subarnapur, Boudh, Gajapati Kandhamal, Bhadrak	Bolangir, Nuapada, Angul, Bargarh, Cuttack, Kendrapara, Nayagarh, Dhenkanal	Kalahandi, Ganjam, Keonjhar, Sundargarh

States	LALP	LAHP	HALP	НАНР
Punjab	Tarntarn, Hoshiarpur, Mohali,	Bathinda, Shri Mukatsar Sahib,	Amritsar, Pathankot,	Fazilka, Patiala
	Gurdaspur, Barnala	Ferozpur, Sangrur, Jalandhar,	Ludhiana	
		Mansa, Moga, Faridkot, Ropar		
Rajasthan	Jalore	Bhilwara, Bundi, Jhunjhunu,	Churu, Jodhpur, Pali,	Nagaur, Bikaner, Ajmer
		Sikar, Jhalawar, Baran, Kota	Barmer, Hanumangarh	Jaipur, Jaisalmer, Tonk,
		Sawai madhopur, Pratapgarh,		Ganganagar
		Banswara, Dungarpur, Udaipur, Chittor, Sirohi,		
		Dausa, Alwar, Karoli,		
		Bharatpur, Rajsamand,		
		Dholpur		
Tamilnadu	Tiruppur, Virudhunagar, Karur	Theni, Thiruvallur, Namakkal,	Tuticorin, Thiruvarur,	Villupuram, Cuddalore,
		Madurai, Coimbatore, Trichy,	Nagapatinam, Tirunelveli,	Salem, T.V Malai,
		Ariyalur, Kancheepuram,	Dindigul	Krishnagiri, Dharmapuri,
		Pudukkotai, Erode, Ramanad,		Vellore, Thanjavur
		Sivagangai, Perambalur,		
		Kanyakumari, The Nilgris		
Telangana	Mahabubabad, Yadadri, Jangaon,	Nirmal, Wanaparthy, Siddipet,	Mahabubnagar,	Sangareddy, Adilabad,
	Nagarkurnool, Medak	Nizamabad, Rangareddy,	Vikarabad, Suryapet,	Kamareddy, Komaram
	Bhadradri, Jayashankar, Warangal	Mancherial, Jagtial, Peddapalli,	Jogulamba, Nalgonda,	bheem
	rural, Warangal urban	Rajanna,	Khammam	
Uttar Pradesh	Raebareli, Barabanki, Pratapgarh,	Karimnagar, Medchal malkajg Kanpur city, Ballia, Jaunpur,	Jhansi, Hamirpur,	Lalitpur, Jalaun, Fatehpur,
Ottai Frauesii	Sultanpur, Gonda, Aligarh, Amethi,	Ghazipur, Kaushambi,	Mahoba, Banda,	Allahabad, Kanpur dehat,
	Kheri, Shravasti, Chandauli, Sambal,	Azamgarh, Auraiya, Etawah,	Chitrakut, Budaun,	Mirzpur, Sonbhadra,
	B.shahar, Bareilly, Faizabad,	Varanasi, Farrukhabad,	Bahraich, Hardoi, Sitapur,	Shahjahanpur
	Rampur, Mahrajganj, Lucknow,	Gorakhpur, Mainpuri, Deoria,	Unnao, Balrampur	a constant of
	Siddharth nagar, Ambedkar nagar,	S. Ravi das ngr, Mau, Agra,	, 1	
	Etah, Sant kabir ngr, Hathras,	Kannauj, G.buddha ngr.,		
	Firozabad, Kashganj, Saharanpur,	Pilibhit, Ghaziabad		
	Moradabad, Mathura, Amroha, Kushi			
	nagar, Meerut, Hapur, Bijnor,			

States	LALP	LAHP	HALP	НАНР
	Muzaffarnagar, Bagpat, Shamli			
Uttrakhand	Almora, Uttarkashi, Nainital, Bageshwar, Nainital (Plain), Haridwar, Dehradun (Plain)	Udhamsinghnagar, Dehradun , Chamoli, Dehradun (Hill), Nainital (Hill), Champawat, Rudraprayag	Uttarakhand (Hill), Tehri Garhwal, Pauri Garhwal, Uttarakhand (Plain), Pithoragarh	
West Bengal	Coochbehar, Uttar Dinajpur, Alipurduar, Purba Bardhaman, Jalpaiguri, Darjeeling, Jhargram, Bankura, Dakshin, Dinajpur, Paschim Bardhaman	Paschim Medinipur, Howrah, Hooghly, Kalimpong	Nadia, South 24-Parganas, North 24-Parganas, Purulia	Murshidasbad, Birbhum, Malda, Purba Medinipur

2. Gram

States	LALP	LAHP	HALP	НАНР
Andhra Pradesh	Ananthapurum, Nellore, Chittor	Guntur, Krishna, East Godavari, Vizianagaram, Visakhapatnam, West Godavari	Kurnool, Kadapa	Prakasham
Bihar	Nawada,Shekhpura, Kisangunj, Muzaffarpur, Katihar, Purnia, Begusarai, W.Champaran, Vaishali, Madhubani	Buxer, Arwal, Mungher, Siwan, Araria, Samastipur, Darbhanga	Aurangabad, Bhabhua, Rohtash, Jamui, Bhagalpur	Bhojpur, Patna, Nalanda, Gaya, Lakhisarai, Banka, Jahanabad
Chhattisgarh	Korba, Bijapur, Dantewada, Gariaband, Balrampur, Surajpur, Narayanpur	Durg, Balod, Raipur, Koria Kaker, Baloda bazaar, Sarguja, Bilaspur, Raigarh, Kondagaon, Jaspur, Jagdalpur, Mahasamund, Janjgir	Kabir dham, Rajnandgaon	Bemetra, Mungeli, Dhamtari
Jharkhand	Koderma, Ranchi, Sahebganj, East Singhbhum, West Singhbum, Lohardaga, Jaamtara	Ramgarh, Devghar , Gumla Giridih, Bokaro, Dhanbad, Khunti	Garhwa, Latehaar, Godda Pakud	Chatra, Hazaaribagh, Dumka, Simdega, Palamu, Saraikela
Karnataka	Ballari, Chitradurga, Chikmagalur, Haveri, Hassan, Davanagere, Chamarajanagar, Bengaluru ó rural, Mandya Chickballapur, Tumkur, Uttara kannada	Yadgir, Mysore, Shimoga	Vijayapur, Gadag, Raichur, Belgavi, Bagalkote, Bidar, Dharwad Koppal	Gulbarga
Madhya Pradesh	Betul, Mandsaur, Aagar, Seoni, Harda, Datia,Bhopal, Neemuch, Mandla, Singroli, Dindori, Tikamgarh, Umaria, Anuppur, Alirajpur	Jabalpur, Katni, Balaghat Khargone, Sidhi, Khandwa Gwalior, Bhind, Jhabua, Sheopurkala, Morena, Burhanpur, Barwani	Sagar, Vidisha, Ashoknagar, Sehore, Ujjain, Rajgarh, Dhar, Ratlam, Chhindwara, Shajapur	Damoh, Satna, Raisen, Dewas, Narsinghpur, Panna, Chattarpur, Shivpuri, Guna, Indore, Hoshangabad Rewa
Maharashtra	Washim, Aurangabad, Solapur, Nasik, Pune, Dhule, Satara, Chandrapur, Sangli, Nandurbar, Bhandara, Gondia Gadchiroli, Palghar, Thane, Raigad	Akola, Jalna, Jalgaon, Wardha, Kolhapur	Yavatmal, Osmanabad, Beed, Ahmednagar, Parbhani, Hingoli	Latur, Nanded, Buldhana, Amravati, Nagpur

States	LALP	LAHP	HALP	НАНР
Odisha	Koraput, Jagatsinghpur, Bargarh Puri, Cuttack, Nayagarh, Khordha, Subarnapur, Sambalpur, Jharsuguda, Rayagada, Jajpur, Deogarh, Dhenkanal, Boudh	Bolangir, Ganjam, Kandhamal Balasore, Malkangiri, Kendrapara	Sundargarh, Nuapada, Nabarangpur, Angul	Kalahandi, Mayurbhanj, Keonjhar
Rajasthan	Nagaur, Sirohi	Sikar, Bhilwara, Jodhpur, Jhalawar, Jalore, Sawai madhopur, Dausa, Pratapgarh, Baran, Dungarpur, Banswara, Alwar Udaipur, Karoli, Kota, Bundi Chittor, Bharatpur, Barmer	Bikaner, Churu, Jaisalmer, Ajmer, Hanumangarh	Jaipur, Ganganagar, Pali, Jhunjhunu, Tonk
Telangana	Medak, Nagarkurnool, Mahabubnagar, Mancherial, Peddapalli, Nalgonda, Suryapet, Khammam, Mahabubabad	Komaram bheem, Rangareddy, Rajanna, Jangaon, Warangal rural, Jagtial, Warangal urban Medchal malkajg, Yadadri	Jogulamba, Nirmal	Kamareddy, Adilabad, Nizamabad, Sangareddy, Siddipet, Vikarabad
Uttar Pradesh	Raebareli, Pratapgarh, Unnao, Chandauli, Gonda, Lucknow, Basti, S. Ravi das ngr, Hardoi Sitapur, Gorakhpur, Balrampur, Sant kabir ngr, Kheri, Etah, Budaun, Deoria, B.shahar, Shahjahanpur, Bahraich, Kashganj, Shravasti, Aligarh, Saharanpur, Muzaffarnagar, Bijnor, Moradabad, Siddharth nagar, Mahrajganj, Pilibhit, Meerut, Kushi nagar	Ghazipur, Jaunpur, Auraiya Azamgarh, Ballia, Amethi, Varanasi, Agra, Sultanpur, Etawah, Faizabad, Mau, Ambedkar nagar, Kannauj, Mainpuri, Farrukhabad, Barabanki, Firozabad, Hathras, Shamli, Bareilly	Hamirpur, Banda, Chitrakut, Mahoba, Allahabad, Kanpur city, Mirzpur, Sonbhadra, Kaushambi	Jhansi, Fatehpur, Jalaun, Kanpur dehat, Lalitpur
West Bengal	North 24-Parganas, South 24-Parganas, Uttar Dinajpur, Dakshin Dinajpur, Darjeeling, Howrah	Purba Bardhaman, Bankura, Alipurduar, Purulia, Hooghly Paschim Medinipur, Paschim Bardhaman	Nadia, Murshidasbad, Malda	Birbhum

3. Arhar

States	LALP	LAHP	HALP	НАНР
Andhra Pradesh	Kadapa, East Godavari,	Krishna, Vizianagaram,	Kurnool, Ananthapurum,	Prakasham, Guntur
	Chittor	Visakhapatnam, West		
		Godavari, Nellore		
Bihar	Saran, Aurangabad,	Sitamarhi, Gopalgunj,	Bhabhua, Nawada,	Jamui, Begusarai, Gaya,
	Jahanabad, W.Champaran,	Muzaffarpur, Bhagalpur,	Rohtash, Banka,	E.Champaran, Siwan,
	Bhojpur, Nalanda, Purnia,	Buxer, Darbhanga,	Samastipur, Patna	Vaishali
	Araria, Kisangunj, Supaul,	Madhubani, Shekhpura,		
	Madhepura, Arwal	Khagaria, Sheohar, Mungher		
Chhattisgarh	Baloda bazaar, Bilaspur, Balod	Kondagaon, Durg, Janjgir,	Gariaband, Kabirdham,	Rajnandgaon, Balrampur,
	Dhamtari, Mahasamund,	Korba, Mungeli	Koria, Jaspur, Surajpur	Raigarh, Sarguja, Bemetra
	Sukama, Bijapur, Jagdalpur,			
	Raipur, Dantewada,			
	Narayanpur,			
Gujrat	Navsari , Arvalli,	Mahisagar, Kutch, Anand,	Bharuch, Narmada, Tapi	Vadodara, Chhotaudepur,
	Valsad, Botad, Kheda, Morbi,	Patan, Banaskantha , Gir		Panchmahal, Dahod,
	Porbandar,	Somnath		Surat, Sabarkantha
	Mehesana, Devbhumi Dwarka			
Jharkhand	Bokaro, Pakud, West	Saraikela, Khunti, Giridih,	Garhwa, Simdega,	Latehaar, Chatra
	Singhbum, Ramgarh, Devghar,	Dhanbad	Gumla, Dumka, Ranchi,	
	Koderma, Sahebganj,		Hazaaribagh, East	
	Lohardaga, Jaamtara, Palamu		Singhbhum	
Karnataka	Bagalkote, Koppal,	Bengaluru ó rural, Bengaluru	Vijayapur, Yadgir,	Gulbarga, Bidar
	Chitradurga, Tumkur,	ó urban, Chikmagalur,	Raichur	
	Chickballapur, Davanagere,	Shimoga, Uttara kannada		
	Ballari, Belgavi, Kolar, Gadag			
	Haveri, Ramanagaram,			
	Mysore, Hassan, Dharwad,			
	Mandya, Chamarajanagar			

States	LALP	LAHP	HALP	НАНР
Madhya Pradesh	Bhopal, Sagar, Umaria, Jabalpur, Seoni, Khandwa, Sehore, Anuppur, Katni, Chattarpur, Barwani, Dhar, Ujjain, Vidisha, Aagar,	Mandla, Jhabua, Bhind, Panna Khargone, Balaghat, Burhanpur	Satna, Rewa, Betul, Singroli	Narsinghpur, Raisen, Sidhi Chhindwara, Damoh, Morena, Hoshangabad
Maharashtra	Sheopurkala Ahmednagar, Jalgaon, Nandurbar, Nasik, Sangli, Bhandara, Dhule, Gondia, Pune, Satara, Kolhapur, Palghar, Raigad, Ratnagiri Thane, Sindhudurg	Chandrapur, Gadchiroli	Amravati, Osmanabad, Solapur, Buldhana, Wardha, Nanded, Beed Washim, Parbhani, Aurangabad, Hingoli	Yavatmal, Latur, Jalna, Akola, Nagpur
Odisha	Keonjhar, Gajapati, Dhenkanal Jajpur, Bargarh, Nabarangpur Deogarh, Malkangiri, Khordha Nayagarh, Cuttack, Kendrapara, Jagatsinghpur, Balasore	Sambalpur, Jharsuguda, Subarnapur, Kandhamal	Mayurbhanj, Nuapada, Koraput, Boudh	Sundargarh, Kalahandi, Rayagada, Ganjam, Bolangir, Angul
Rajasthan	Sawai madhopur, Jaipur, Karoli, Bharatpur, Jhalawar, Sirohi, Bundi, Pali, Dausa, Baran Kota, Bikaner, Chittor, Tonk, Jaisalmer, Sikar	Dholpur, Pratapgarh, Bhilwara	Banswara, Hanumangarh, Dungarpur	Ganganagar, Udaipur, Alwar
Tamilnadu	Erode, Namakkal, Villupuram, Thanjavur, Sivagangai, Kancheepuram, Tirunelveli, Tuticorin, Ramanad, Kanyakumari	T.V Malai, Dindigul, Perambalur, Pudukkotai, Thiruvallur, Cuddalore, Tiruppur	Vellore, Dharmapuri, Salem, Karur, Trichy, Theni	Krishnagiri, Madurai

States	LALP	LAHP	HALP	НАНР
Telangana	Yadadri, Nagarkurnool, Jangaon, Siddipet, Kamareddy, Khammam, Medak, Mahabubabad, Bhadradri, Warangal rural, Nizamabad, Warangal urban, Jayashankar	Peddapalli, Karimnagar, Medchal malkajg, Rangareddy, Nirmal, Mancherial, Rajanna, Jagtial	Mahabubnagar, Sangareddy, Jogulamba, Nalgonda, Suryapet, Wanaparthy	Vikarabad, Adilabad, Komaram bheem
Uttar Pradesh	Gorakhpur, Varanasi, Mahoba, Amethi, Bahraich, Mathura, Basti, Chandauli, Budaun, Sant kabir ngr, Siddharth nagar Bareilly, Ambedkar nagar, Etah, Unnao, Jhansi, Barabanki, Faizabad, Shravasti, Firozabad, Kheri, Hardoi, Kashganj, Agra, Mainpuri, Kushi nagar, Lucknow, Mahrajganj, Muzaffarnagar, Bijnor, Shahjahanpur	S. Ravi das ngr, Sitapur, G.buddha ngr., Auraiya, Etawah, Mau, Farrukhabad, Kannauj, Hapur, Ghaziabad, Meerut, Bagpat, Amroha, Sambal, Rampur	Aligarh, Hamirpur, Ballia, Sonbhadra, Jaunpur, B.shahar, Balrampur, Ghazipur, Hathras, Sultanpur, Gonda, Raebareli, Deoria	Fatehpur, Chitrakut, Banda, Mirzapur, Allahabad, Kanpur dehat, Azamgarh, Kaushambi, Pratapgarh, Jalaun, Kanpur city
Uttrakhand	Dehradun , Chamoli, Almora, Champawat, Nainital, Haridwar	Rudraprayag	Pauri Garhwal, Uttarkashi	Tehri Garhwal
West Bengal	Hooghly, Purba Bardhaman, Paschim Medinipur, Darjeeling, North 24-Parganas, Bankura, Paschim Bardhaman	Malda, Murshidasbad	Purulia, South 24- Parganas, Jhargram, Nadia, Birbhum	

4. Urd

States	LALP	LAHP	HALP	НАНР
Andhra Pradesh	Prakasham, Vizianagaram, West Godavari, Nellore, Chittoor	Kurnool, Kadapa, Visakhapatnam, Ananthapuram	Guntur, Srikakulam	Krishna, East Godavari
Bihar	Madhepura, Bhojpur, Darbhanga Nalanda, Arwal, Buxer	Gaya, Araria, W.Champaran, Kisangunj, Vaishali, Banka, Saran, Rohtash, Nawada, Gopalgunj, Lakhisarai, Purnia, Aurangabad, Saharsa, Jahanabad	Khagaria, Begusarai, Madhubani	Katihar, Bhagalpur, Samastipur, Supaul, Siwan
Chhattisgarh	Korba, Balod , Janjgir, Bilaspur, Mungeli	Baloda bazaar, Jagdalpur Kabir dham, Narayanpur, Bijapur, Sukama, Dhamtari, Raipur, Bemetra, Dantewada Durg	Raigarh, Mahasamund, Koria, Kaker, Surajpur	Jaspur, Kondagaon, Rajnandgaon, Gariaband, Sarguja, Balrampur
Gujrat	Devbhumi Dwarka, Mahisagar, Navsari	Valsad , Dang , Bharuch , Tapi Panchmahal, Surat, Porbandar , Narmada , Botad , Vadodara	Patan, Mehesana, Sabarkantha , anaskantha	Arvalli, Chhotaudepur , Dahod , Junagadh
Jharkhand	Godda, Latehaar, Ramgarh, Dumka, Bokaro, Garhwa, Pakud, Lohardaga, Devghar, Koderma, Dhanbad, Jaamtara	Chatra, Giridih	Palamu, Ranchi, Hazaaribagh, East Singhbhum, West Singhbum	Simdega, Saraikela, Sahebganj, Gumla, Khunti
Karnataka	Tumkur, Mandya, Bagalkote, Haveri, Gadag, Bengaluru ó urban, Dharwad, Hassan Yadgir, Vijayapur, Chikmagalur, Ballari	Udupi, Koppal, Dakshina kannada, Uttara kannada, Davanagere, Shimoga, Bengaluru - rural	Mysore, Belgavi, Chamarajanagar	Bidar, Gulbarga

States	LALP	LAHP	HALP	НАНР
Madhya Pradesh	Rewa, Singroli, Anuppur, Seoni, Dindori, Sidhi, Umaria Bhopal, Betul, Khargone, Katni	Ratlam, Panna, Narsinghpur Rajgarh, Shahdol, Neemuch Jhabua, Barwani, Khandwa, Gwalior, Hoshangabad, Dhar Aagar,Burhanpur, Bhind, Mandla, Ujjain, Balaghat	Sagar, Ashoknagar, Shivpuri, Guna, Satna, Raisen, Datia	Tikamgarh, Chattarpur, Damoh, Alirajpur, Mandsaur, Jabalpur
Maharashtra	Latur, Nandurbar, Parbhani, Yavatmal, Hingoli, Dhule, Pune, Kolhapur, Wardha, Nagpur, Chandrapur, Raigad, Sindhudurg, Ratnagiri Gadchiroli	Nasik, Aurangabad, Satara, Palghar, Thane	Osmanabad, Solapur, Nanded, Ahmednagar, Beed, Amravati	Jalgaon, Akola, Buldhana Jalna, Sangli, Washim
Odisha	Sambalpur, Bargarh, Nayagarh Rayagada, Mayurbhanj, Deogarh, Koraput, Boudh, Kandhamal, Nabarangpur, Malkangiri, Jharsuguda	Jagatsinghpur, Gajapati, Khordha, Balasore, Subarnapur, Bhadrak	Angul, Nuapada, Bolangir, Sundargarh	Kalahandi, Cuttack, Kendrapara, Ganjam, Puri, Jajpur
Rajasthan	Udaipur, Chittor, Sirohi, Ganganagar, Dholpur, Karoli, Bikaner, Jalore	Dungarpur, Banswara, Pratapgarh, Bharatpur, Hanumangarh, Jodhpur, Barmer, Churu, Jhunjhunu	Bundi, Bhilwara, Tonk	Kota, Baran, Jhalawar, Ajmer, Sawai madhopur
Tamilnadu	Virudhunagar, Ramanad, Tiruppur, Thiruvallur, Namakkal	Salem, Ariyalur, Pudukkotai, Kancheepuram, Vellore, Trichy, Dharmapuri, Theni, Krishnagiri, Madurai, Erode, Karur, Sivagangai, Coimbatore, Kanyakumari, Perambalur	Tuticorin, Nagapatinam, Tirunelveli, Dindigul	Villupuram, Cuddalore, T.V Malai, Thanjavur

States	LALP	LAHP	HALP	НАНР
Telangana	Medak, Rajanna, Mahabubabad Jogulamba, Komaram bheem, Jayashankar, Bhadradri, Khammam, Nalgonda, Mahabubnagar, Suryapet, Nagarkurnool, Warangal rural Peddapalli, Siddipet, Yadadri, Mancherial	Nizamabad, Karimnagar, Rangareddy, Warangal urban	Sangareddy, Nirmal, Adilabad, Wanaparthy, Jagtial	Vikarabad, Kamareddy
Uttar Pradesh	Amethi, Jalaun, Fatehpur, Lucknow, Faizabad, Banda, Allahabad, Aligarh, Balrampur, Kaushambi, Shravasti, Agra Siddharth nagar, Mathura, Deoria, Sant kabir ngr	Kanpur city, Pilibhit, Jaunpur Kanpur dehat, Sultanpur, Moradabad, Bareilly, Kheri, Saharanpur, Farrukhabad Amroha, Sonbhadra Firozabad, Bijnor, Ambedkar nagar, Varanasi, Auraiya, Rampur, Kashganj, Chitrakut, Mainpuri, Bahraich, Kannauj, Etawah, Gonda, B.shahar, Meerut, Hapur, Etah, Muzaffarnagar, Shamli, Ghazipur, Bagpat, Mirzpur, S. Ravi das ngr, Azamgarh, Basti, Ghaziabad, Hathras, Kushi nagar, Chandauli, G.buddha ngr., Mahrajganj, Gorakhpur Ballia, Mau	Lalitpur, Jhansi, Mahoba, Hamirpur, Sitapur, Raebareli, Barabanki	Budaun, Hardoi, Unnao, Sambal, Pratapgarh, Shahjahanpur
Uttrakhand	Uttarkashi, Pithoragarh, Dehradun , Udhamsinghnagar Bageshwar	Chamoli, Champawat, Rudraprayag, Haridwar	Pauri Garhwal, Tehri Garhwal, Nainital	Almora
West Bengal	Uttar Dinajpur, Dakshin Dinajpur, Purba Medinipur, Birbhum, Jhargram, Hooghly	Alipurduar, Jalpaiguri, South 24- Parganas, Darjeeling, Paschim Medinipur, Purba Bardhaman, Bankura, Kalimpong, Howrah	Purulia, Coochbehar	Murshidasbad, Nadia, Malda, North 24-Parganas

5. Mung bean

States	LALP	LAHP	HALP	НАНР
Andhra	West Godavari,	Krishna, Kurnool, Nellore,	Srikakulam, Guntur,	East Godavari, Prakasham
Pradesh	Kadapa, Ananthapuram	Chittoor	Vizianagaram	
Bihar	Sitamarhi, Nawada, Saran, Banka, Bhagalpur, E.Champaran, Sheohar,	Purnia, Gaya, Katihar, Nalanda, Khagaria, Jamui, Begusarai, Mungher, Jahanabad, Arwal,	Muzaffarpur, Supaul, Madhepura, Vaishali	Saharsa, Madhubani, Samastipur Darbhanga, Araria, Kisangunj
	Gopalgunj, Siwan, W.Champaran, Rohtash, Patna, Bhabhua, Bhojpur, Buxer	Shekhpura, Lakhisarai, Aurangabad		
Chhattisgarh	Sukama, Koria, Balod, Surajpur, Janjgir, Korba, Dhamtari, Kabirdham, Bilaspur, Sarguja, Dantewada, Jaspur, Raipur, Durg	Bijapur, Kondagaon, Baloda bazaar, Jagdalpur, Mungeli, Balrampur, Bemetra	Gariaband, Mahasamund	Raigarh, Rajnandgaon, Kaker
Gujrat	Tapi , Arvalli , Narmada Vadodara, Navsari	Gandhinagar , Bhavnagar , Sabarkantha , Porbandar , Mahisagar , Chhotaudepur Panchmahal	Kutch, Mehesana	Bharuch, Banaskantha, Patan
Jharkhand	Bokaro, Koderma, Devghar, Garhwa, Lohardaga, Pakud, Sahebganj	Palamu, Gumla, Giridih, Godda, Dumka, Latehaar, Chatra, Ramgarh, Dhanbad	East Singhbhum	Saraikela, West Singhbum, Ranchi, Simdega, Hazaaribagh, Khunti
Madhya Pradesh	Sagar, Chattarpur, Anuppur, Ujjain, Datia, Betul, Alirajpur, Balaghat	Panna, Mandsaur, Aagar, Bhind	Rewa, Barwani, Satna, Tikamgarh, Khargone, Damoh	Hoshangabad, Jabalpur, Dhar, Shivpuri, Narsinghpur, Sidhi

States	LALP	LAHP	HALP	НАНР
Karnataka	Vijayapur, Tumkur, Chamarajanagar, Chikmagalur, Haveri, Hassan, Chitradurga, Shimoga, Uttara kannada, Dakshina kannada, Davanagere Ballari, Udupi, Mysore, Bengaluru ó urban, Bengaluru ó rural	Raichur, Mandya	Gadag, Yadgir, Bagalkote, Dharwad, Koppal	Gulbarga, Belgavi, Bidar
Maharashtra	Latur, Nasik, Hingoli, Nandurbar, Yavatmal Nagpur, Wardha, Chandrapur, Palghar, Gadchiroli, Thane	Washim, Sangli, Kolhapur, Sindhudurg, Ratnagiri, Bhandara	Nanded, Amravati, Osmanabad, Dhule, Solapur, Aurangabad, Pune	Ahmednagar, Jalna, Parbhani, Jalgaon, Akola, Buldhana, Beed
Odisha	Subarnapur, Sambalpur, Jajpur, Deogarh, Sundargarh, Nabarangpur, Malkangiri, Rayagada, Jharsuguda, Koraput, Kandhamal	Keonjhar, Boudh, Mayurbhanj, Balasore, Bhadrak, Gajapati	Bargarh, Nayagarh, Nuapada, Jagatsinghpur, Kendrapara, Cuttack, Angul, Khordha, Dhenkanal, Puri	Ganjam, Kalahandi, Bolangir
Rajasthan	Hanumangarh, Jaisalmer, Sirohi, Chittor, Kota, Dausa, Pratapgarh	Jhunjhunu, Bikaner, Sikar, Bhilwara, Sawai madhopur, Udaipur, Jhalawar, Bharatpur, Karoli, Banswara, Dungarpur, Alwar	Pali, Jodhpur, Churu, Jalore, Barmer	Nagaur, Jaipur, Ajmer, Ganganagar, Tonk
Tamilnadu	Virudhunagar, Dindigul, Tiruppur, Krishnagiri, Ramanad	Madurai, Vellore, Theni, Villupuram, Thanjavur, Dharmapuri, Coimbatore, T.V Malai, Kancheepuram, Erode, Trichy, Karur, Sivagangai, Pudukkotai, Ariyalur, The Nilgris	Tuticorin, Thiruvarur, Nagapatinam, Tirunelveli	Salem, Thiruvallur, Cuddalore, Namakkal

States	LALP	LAHP	HALP	НАНР
Telangana	Nalgonda, Yadadri	Mahabubnagar, Komaram bheem, Medak, Jangaon, Jagtial, Adilabad, Nirmal, Jayashankar, Mancherial Rajanna, Bhadradri, Peddapalli, Warangal rural, Wanaparthy, Nizamabad, Siddipet, Karimnagar, Rangareddy, Warangal urban, Nagarkurnool, Jogulamba, Medchal malkajg	Suryapet, Khammam	Sangareddy, Mahabubabad, Vikarabad, Kamareddy
Uttar Pradesh	Jalaun, Kashganj, Agra, Hardoi, Varanasi, Meerut, Sambal, Mirzpur, Budaun, Sonbhadra, Amroha, Rampur, Ghaziabad	Kanpur city, Farrukhabad, Kanpur dehat, Kannauj, Raebareli, Kheri, Jaunpur, Amethi, Sultanpur, Kaushambi, Faizabad, Ghazipur, Gonda, Hapur, Ambedkar nagar, Shahjahanpur, Ballia, Kushi nagar, Lucknow, Mahrajganj, Chandauli, Gorakhpur, Deoria, Mau, Basti, Azamgarh, Bagpat, Barabanki, Shamli, S. Ravi das ngr, Muzaffarnagar, Pilibhit, Bahraich, Bijnor, Sitapur, Saharanpur, Moradabad	Mahoba, Jhansi, Hamirpur, Aligarh, Etah, Mathura, Chitrakut, Banda, Firozabad, Hathras, B.shahar	Fatehpur, Lalitpur, Etawah, Unnao, Allahabad, Mainpuri, Auraiya, G.buddha ngr., Pratapgarh

6. Pea

States	LALP	LAHP	HALP	НАНР
Bihar	Mungher, W.Champaran, Gaya, E.Champaran, Jamui, Muzaffarpur, Shekhpura, Nawada, Sheohar, Sitamarhi, Vaishali, Darbhanga, Araria	Rohtash, Supaul, Gopalgunj, Saharsa, Purnia, Katihar, Saran, Kisangunj, Madhubani, Banka	Begusarai, Khagaria, Jahanabad Lakhisarai, Arwal, Aurangabad	Nalanda, Siwan, Buxer, Patna, Bhabhua, Samastipur, Madhepura
Chhattisgarh	Janjgir, Rajnandgaon, Balod, Mungeli, Kabirdham, Bemetra, Korba, Narayanpur, Durg, Bijapur, Dantewada	Jagdalpur, Mahasamund, Raipur	Kaker, Jaspur	Raigarh, Baloda bazaar, Koria, Sarguja, Gariaband, Balrampur, Dhamtari, Kondagaon, Surajpur, Bilaspur
Jharkhand	Lohardaga, Saraikela, Palamu, Latehaar, Khunti, Sahebganj, Ramgarh, Dhanbad, Bokaro, West Singhbum, Jaamtara, Pakud	Devghar, Godda, Dumka, Giridih, Koderma	Ranchi, Garhwa, East Singhbhum, Hazaaribagh	Gumla, Simdega, Chatra
Madhya Pradesh	Tikamgarh, Vidisha, Raisen, Satna, Rajgarh, Hoshangabad, Balaghat, Anuppur, Betul, Ratlam, Umaria, Ashoknagar, Indore, Aagar	Katni, Bhind, Gwalior	Narsinghpur, Seoni, Dindori, Sagar, Damoh	Panna, Jabalpur, Datia, Chattarpur
Odisha	Nayagarh, Kandhamal, Dhenkanal, Subarnapur, Cuttack, Boudh, Jharsuguda, Nabarangpur, Malkangiri, Khordha, Jajpur, Jagatsinghpur, Balasore, Bhadrak, Deogarh	Sambalpur, Gajapati, Puri Kendrapara	Koraput, Ganjam, Nuapada, Mayurbhanj	Rayagada, Keonjhar, Kalahandi, Bargarh, Bolangir, Sundargarh, Angul

States	LALP	LAHP	HALP	НАНР
Rajasthan	Alwar, Tonk, Chittor,	Hanumangarh, Sikar,	Jaipur, Ajmer	Nagaur, Bundi
	Jhunjhunu, Bhilwara,	Bharatpur, Ganganagar,		
	Dholpur, Sawai madhopur,	Jhalawar, Baran, Rajsamand,		
	Pratapgarh, Udaipur, Kota,	Sirohi, Karoli,		
	Dausa, Bikaner, Banswara,	Jodhpur		
	Churu, Jalore, Jaisalmer			
Uttar Pradesh	Amethi, Basti, Pratapgarh,	Allahabad, Varanasi, Ballia,	Mahoba, Hamirpur,	Jalaun, Lalitpur, Jhansi,
	Sant kabir ngr, Mirzpur,	Kanpur city, Kanpur dehat,	Sultanpur	Azamgarh, Jaunpur
	Ghazipur, Sonbhadra, Banda,	Fatehpur, Mau, Mainpuri,	_	
	Siddharth nagar, Ambedkar	Mahrajganj, Farrukhabad,		
	nagar, Barabanki, Rampur,	Kannauj, Kushi nagar,		
	Sitapur, Raebareli, Faizabad,	Firozabad, Shravasti,		
	Chandauli	Etawah, Meerut, Aligarh,		
	Kaushambi, Chitrakut,	Agra, Saharanpur,		
	Hardoi, S. Ravi das ngr,	Ghaziabad, Auraiya,		
	Budaun, Lucknow, Unnao,	Muzaffarnagar, Bagpat,		
	Kheri, Sambal, Moradabad,	Mathura		
	Amroha, Bijnor, Hapur,			
	Hathras, Shamli, G.buddha			
	ngr.			

7. Lentil

States	LALP	LAHP	HALP	НАНР
Bihar	Nawada, Jamui, Sitamarhi, Bhagalpur, Darbhanga, Banka, Supaul, Muzaffarpur, Kisangunj, Begusarai, Madhepura, Sheohar, Khagaria, Siwan, Araria, Vaishali, Saran, Gopalgunj	Lakhisarai, Shekhpura, Rohtash, Buxer, Banka, Samastipur, Katihar, Mungher, Saharsa	Nalanda, Aurangabad, Madhubani, E.Champaran, Gaya, Jahanabad, Arwal	Patna, W.Champaran, Bhabhua, Bhojpur
Chhattisgarh	Balrampur, Surajpur, Gariaband, Kaker, Jaspur, Kondagaon, Korba, Jagdalpur, Narayanpur, Mahasamund	Dhamtari, Durg, Balod, Bilaspur	Rajnandgaon, Kabirdham, Koria	Bemetra, Sarguja, Raipur, Raigarh, Baloda bazaar, Mungeli
Jharkhand	Latehaar, East Singhbhum, Ranchi, Hazaaribagh, Bokaro	Devghar, Gumla, Khunti, Giridih	Garhwa, Palamu, Pakud, Godda, Sahebganj,Dumka	Chatra, Simdega, West Singhbum, Koderma,
Rajasthan	Chittor, Bharatpur, Ajmer, Baran, Kota, Jhunjhunu, Dholpur, Alwar, Bikaner, Sikar, Jaipur, Pali, Karoli, Dausa, Ganganagar	Sawai madhopur	Bundi	Bhilwara, Jhalawar, Pratapgarh, Tonk
Madhya Pradesh	Shajapur, Ashoknagar, Katni, Anuppur, Mandsaur, Sidhi, Bhind, Umaria, Singroli, Shahdol, Chhindwara, Sehore, Tikamgarh, Balaghat, Morena	Chattarpur, Shivpuri, Bhopal, Datia, Guna	Satna, Dindori, Rajgarh, Rewa, Damoh, Panna, Mandla, Seoni	Sagar, Narsinghpur, Vidisha, Raisen, Jabalpur
Uttar Pradesh	Mirzpur, Bareilly, Kushi nagar, Saharanpur, Lucknow, Gorakhpur, Farrukhabad, Fatehpur, Sambal, Bijnor, Kanpur city, Unnao, Moradabad, Kanpur dehat, Deoria, Muzaffarnagar, Kannauj, Pratapgarh, Hapur, Varanasi, Jaunpur, Kaushambi, Bagpat, Mainpuri, Shamli, Amroha, Raebareli, Mathura, Auraiya, Etawah	Sultanpur, Budaun, Faizabad Pilibhit, Etah, Aligarh, Amethi, Rampur, Kashganj, Ambedkar nagar, Hathras, MaU, Firozabad, Ghaziabad, S. Ravi das ngr	Bahraich, Banda, Hamirpur, Balrampur, Chitrakut, Sitapur, Mahoba, Jhansi, Shravasti, Kheri, Gonda, Allahabad, Chandauli, Mahrajganj, Sonbhadra	Jalaun, Shahjahanpur, Lalitpur, Ballia, Barabanki, Ghazipur, Hardoi
West Bengal	Alipurduar, Purba Bardhaman, Bankura, Coochbehar, Hooghly, Uttar Dinajpur, Jalpaiguri, Jhargram, Darjeeling, Paschim Bardhaman, Dakshin Dinajpur, Howrah, Purulia	Paschim Medinipur, South 24- Parganas, Purba Medinipur	Nadia, North 24-Parganas	Murshidasbad, Birbhum Malda

8. Lathyrus

States	LALP	LAHP	HALP	НАНР
Bihar	Bhabhua, Nawada,	Buxer, Samastipur,	Aurangabad, Jahanabad,	Patna, Rohtash, Bhojpur,
	Muzaffarpur, Gaya, Arwal,	Kisangunj, Katihar, Purnia,	Nalanda, Bhagalpur	Lakhisarai, Shekhpura,
	E.Champaran, Banka,	Araria, Madhepura, Khagaria,		Supaul
	Sheohar, Sitamarhi, Vaishali,	Madhubani, Mungher,		
	W.Champaran	Begusarai, Saran		
Chhattisgarh	Korba, Gariaband, Durg,	Kaker, Raigarh,	Balod , Baloda bazaar,	Mungeli, Bilaspur, Dhamtari,
_	Kabir dham, Jaspur, Koria,	Mahasamund, Surajpur,	Rajnandgaon, Raipur,	Janjgir
	Kondagaon, Narayanpur,	Balrampur	Bemetra	
	Jagdalpur, Sarguja			
West Bengal	Howrah, North 24-Parganas,	Paschim Medinipur,	South 24-Parganas,	Purba Medinipur, Malda
	Alipurduar, Coochbehar,	Jhargram, Bankura	Murshidasbad, Nadia,	
	Purba Bardhaman, Purulia,		Birbhum	
	Jalpaiguri, Darjeeling, Uttar			
	Dinajpur, Dakshin Dinajpur,			
	Hooghly, Paschim			
	Bardhaman			

9. Horse gram

States	LALP	LAHP	HALP	НАНР
Andhra Pradesh	Ananthapuram, Kadapa, Kurnool	Visakhapatnam, Krishna, Prakasham, Nellore, West Godavari, East Godavari	Vizianagaram, Chittoor	Srikakulam
Bihar	Begusarai, Nawada, Mungher	Madhubani, Saharsa, Katihar Purnia, W.Champaran, Buxer	Supaul, Jamui, Khagaria	Gaya, Araria, Banka, Kisangunj, Bhagalpur
Chhattisgarh	Jaspur, Mahasamund	Balrampur, Surajpur, Kondagaon, Narayanpur, Dantewada, Bijapur, Bilaspur, Sukama, Balod, Dhamtari, Janjgir, Baloda bazar	Raigarh, KakerSarguja, Jagdalpur	Koria, Gariaband, Rajnandgaon, Korba
Jharkhand	Saraikela, Gumla, Bokaro, Pakud, Godda, Koderma, Devghar, Palamu, Ramgarh, Dhanbad, Latehaar, Garhwa	Chatra	Hazaaribagh, West Singhbum, Lohardaga, East Singhbhum, Ranchi	Simdega
Karnataka	Haveri, Hassan, Gadag, Dharwad, Dakshina kannada, Shimoga, Udupi	Chitradurga, Belgavi, Ballari, Bengaluru ó rural, Chickballapur, Bagalkote, Davanagere, Uttara kannada Bengaluru ó urban, Gulbarga, Vijayapur Raichur Bidar, Yadgir	Mandya, Chamarajanagar, Ramanagaram, Mysore, Kolar	Tumkur, Koppal, Chikmagalur

States	LALP	LAHP	HALP	НАНР
Odisha	Puri, Khordha	Jajpur, Nayagarh, Jharsuguda, Subarnapur, Koraput, Jagatsinghpur, Sambalpur, Deogarh, Kendrapara, Bargarh, Malkangiri, Gajapati, Bhadrak, Balasore, Nabarangpur	Kandhamal, Dhenkanal	Bolangir, Sundargarh, Kalahandi, Keonjhar, Angul, Nuapada, Cuttack, Ganjam, Rayagada, Mayurbhanj, Boudh
Tamilnadu	Salem, Erode, Pudukkotai, Kancheepuram	Sivagangai, Villupuram, Virudhunagar, Madurai, Tuticorin, Perambalur, Ariyalur, Thiruvallur, Ramanad, Tirunelveli	Tiruppur, Dharmapuri, Dindigul	Krishnagiri, Vellore, T.V Malai
Telangana	Jangaon, Nagarkurnool, Komaram bheem, Mancherial	Siddipet, Medak, Wanaparthy, Mahabubnagar, angareddy, Vikarabad, Jayashankar	Rangareddy, Jogulamba, Suryapet	Nalgonda, Yadadri
Uttrakhand	Uttarkashi, Dehradun, Bageshwar	Pithoragarh, Nainital , Champawat, Chamoli, Rudraprayag	Pauri Garhwal, Almora	Tehri Garhwal

Crop-wise and State-wise districts of higher and lower National Yield.

Ind	lia	High yielding districts	Total d	istricts	Percentage
		312	58	33	53.51%
Crop/Season	National Yield		Yield	l in Districts	
Crop/Season	ivational lieu	Higher than Nati	onal		Lower than National
Total Pulses	786 kg/ha	AP (02)- Visakhapatnam, Praka	asham	` /	, Krishna, Guntur, East Godavari, ulam , Vizianagaram, Kadapa, West or, Ananthpuram
		Assam (09)- Darrang, Hailak	andi, Lakhimpur,	Assam (18)- I	Baksa, Barpeta, Bongaigaon, Cachar,
		Morigaon, Nalbari, Sivas Tinsukia, Udalguri		Chirang, Dhema Golaghat, Jorha	aji, Dibrugarh, Dima Hasao, Goalpara, t, Kamrup (Metro), Kamrup (Rural), Karimganj, Kokrajhar, Nagaon
		Bihar (31)- Lakhisarai, Buxer Bhojpur, Shekhpura, W.Cha Mungher, Gaya, Banka, Kisangunj, Siwan, Begu Gopalgunj, Nalanda, Jahanaba Bhabhua, Khagaria, Arwal, Dar Samastipur, Aurangabad, Madhubani, Saran CG (05)- Bemetara, Mungeli,	amparan, Jamui, Purnia, Araria, sarai, Katihar, d, E.Champaran, bhanga, Nawada, Bhagalpur,	Sheohar, Vaisha	1
		Kabirdham		Balodabazaar, Jagdalpur, Kor Surajpur, Kar Narayanpur, Dar	nandgaon, Bilaspur, Raipur, Janjgir, Kondagaon, Balrampur, Balod, riya, Sarguja, Gariyaband, Korba, nker, Jashpur, Sukma, Raigarh, ntewada, Mahasamund, Bijapur mreli, Porbandar, Botad, Jamnagar,
		Gujrat (15)-Mahisagar , Pan		• , ,	
		Vadodara, Anand, Dahod, Bharuch, Dang, Junagadh, Ta Narmada, Arvalli, Rajkot	- '	Gandhinagar, S	Gir Somnath, Bhavnagar, Ahmedabad, urendranagar, Navsari, Morbi, Patan, ka, Banaskantha, Kutch, Mehesana

Cyon/Soogan	National Yield	Yield in Dis	stricts
Crop/Season	National Yleid	Higher than National	Lower than National
Total Pulses	786 kg/ha	J&K (01)- Kishtwar	J&K (08)- Doda, Rajouri, Poonch, Reasi, Jammu, Udhampur, Kathua, Samba
		JH (21)- Chatra, Simdega, Saraikela, Gumla, Giridih, Dhanbad, Khoonti, Palamu, Devghar, Latehaar, Koderma, Ramgarh, Dumka, Hazaaribagh, Sahebganj, Ranchi, Godda, West Singhbum, Bokaro, Garhwa, Pakud	JH (03)- Jaamtara, Lohardaga, East Singhbhum
		Karnataka (03)-Gulbarga, Bidar, Bengaluru ó rural	Karnataka (27)- Vijayapur , Raichur, Yadgir, Davanagere, Bengaluru ó urban, Ramanagaram, Shimoga, Belgavi, Mysore, Udupi, Hassan, Ballari, Chickballapur, Bagalkote, Koppal, Dakshina kannada, Uttara kannada, Haveri, Chikmagalur, Kolar, Chitradurga, Dharwad, Tumkur Kodagu, Gadag, Mandya, Chamarajanagar
		MP (47)- Khargone, Gwalior, Khandwa, Chhindwara, Narsinghpur, Indore, Dewas, Jabalpur, Hoshangabad, Sheopurkala, Bhind, Jhabua, Katni, Harda, Raisen, Panna, Balaghat, Ujjain, Neemuch, Sehore, Burhanpur, Sidhi, Morena, Bhopal, Damoh, Satna, Mandla, Betul, Guna, Datia, Dhar, Chattarpur, Sagar, Shivpuri, Dindori, Rewa, Barwani, Seoni, Ratlam, Aagar, Tikamgarh, Singroli, Ashoknagar, Mandsaur, Umaria, Rajgarh, Vidish	MP (04)- Shajapur, Alirajpur, Shahdol, Anuppur
		MS (18)- Latur, Nagpur, Akola, Jalna, Yavatmal, Wardha, Nanded, Amravati, Chandrapur, Buldhana, Jalgaon, Hingoli, Parbhani, Gadchiroli, Aurangabad, Kolhapur, Beed, Osmanabad	MS (16)- Sangli, Ahmednagar, Bhandara, Pune, Dhule, Nandurbar, Raigad, Washim, Satara, Thane, Nasik, Ratnagiri, Palghar, Sindhudurg, Solapur, Gondia

Cuar /Caasar	Notional Viold	Yield in Dis	tricts
Crop/Season	National Yield	Higher than National	Lower than National
Total Pulses	786 kg/ha	Meghalaya (11)- East Khasi hills, South West Khasi Hills, West Khasi Hills, South garo hills, West Garo hills, South West garo Hills, Ri- Bhoi Distt, West Jaintia, North Garo Hills, East Jaintia, East Garo Hills Nagaland (11)- Kiphiri, Tuensang, Longleng, Wokha, Mon, Kohima, Mokokchung, Phek, Zunheboto, Peren,	
		Dimapur Punjab (14)- Patiala, Sangrur, Bathinda, Mukatsar, Faridkot, Ferozepur, Jalandhar, Mansa, Moga, Ropar, Fazilka, Mohali, Amritsar, Ludhiana	Punjab (05)- T.Taran, Barnala Gurdaspur, Pathankot, Hoshiarpur
		RJ (18)- Karoli, Dausa, Alwar, Pratapgarh, Bharatpur, Sawai madhopur, Dholpur, Baran, Kota, Sikar, Udaipur, Dungarpur, Jhunjhunu, Jaipur, Banswara, Jhalawar, Chittor, Ganganagar	RJ (15)- Ajmer, Bundi, Jaisalmer, Tonk, Rajsamand, Bhilwara, Bikaner, Nagaur, Pali, Hanumangarh, Sirohi, Jalore, Jodhpur, Churu, Barmer
		TN (01)- The Nilgiris	TN (30)- Perambalur, Vellore, Thanjavur, Thiruvallur, Salem, Namakkal, Madurai, Villupuram, Sivagangai, Cuddalore, Krishnagiri, Kancheepuram, Pudukkotai, Coimbatore, T.V Malai, Dharmapuri, Trichy, Erode, Kanyakumari, Ariyalur, Theni, Dindigul, Virudhunagar, Ramanad, Thiruvarur, Tiruppur, Nagapatinam, Karur, Tuticorin, Tirunelveli
		Telangana (14) - Nizamabad, Adilabad, Kamareddy, Komaram bheem, Mancherial, Nirmal, Jagtial, Siddipet, Medchal malkajg, Karimnagar, Peddapalli, Rangareddy, Rajanna, Sangareddy	Telangana (16)- Jayashankar, Mahabubabad, Vikarabad, Medak, Bhadradri, Wanaparthy, Warangal rural, Warangal urban, Yadadri, Jangaon, Nalgonda, Jogulamba, Nagarkurnool, Mahabubnagar, Suryapet, Khammam

Cyon/Saggar	National Viold	Yield in Distri	cts
Crop/Season	National Yield	Higher than National	Lower than National
Total Pulses	786 kg/ha	UP (58)- Azamgarh, S. Ravi das ngr, Mau, Etawah, Jalaun, Ballia, Kanpur dehat, Agra, Kaushambi, Jaunpur, Ghazipur, Mainpuri, Pilibhit, Auraiya, Varanasi, Kannauj, Kanpur city, Ghaziabad, Mirzpur, Fatehpur, Shahjahanpur, Allahabad, Farrukhabad, G.buddha ngr., Gorakhpur, Deoria, Sonbhadra, Basti, Lalitpur, Sant kabir ngr, Sambal, Rampur, Siddharth nagar, Jhansi, Bagpat, Chitrakut, Pratapgarh, Gonda, Meerut, Firozabad, Ambedkar nagar, Hapur, Hathras, Amroha, Barabanki, Sultanpur, Kushi nagar, Bareilly, Bahraich, Kheri, Banda, Hamirpur, Bijnor, Faizabad, Amethi, Mahrajganj, Balrampur, Aligarh UK (16)- Udhamsinghnagar, Chamoli, Dehradun (Hill),	UP (17) ó Etah, Chandauli, Sitapur, Shravasti, Muzaffarnagar, Kashganj, Saharanpur, Shamli, Hardoi, Budaun, Lucknow, Mahoba, Moradabad, Unnao, Raebareli, B.shahar, Mathura UK (03)-Haridwar, Nainital, Pauri Garhwal
		Dehradun , Champawat, Rudraprayag, Uttarakhand (Plain), Nainital (Hill), Tehri Garhwal, Bageshwar, Uttarakhand (Hill), Uttarkashi, Almora, Dehradun (Plain), Pithoragarh, Nainital WB (17)- Purba Medinipur, Kalimpong, Birbhum, Murshidasbad, Hooghly, Paschim Medinipur, Malda, Howrah, Purba Bardhaman, Bankura, Jalpaiguri, Jhargram, Nadia, North -Parganas, Darjeeling, Uttar Dinajpur,	WB (05)- South 24-Parganas, Paschim Bardhaman, Coochbehar, Dakshin Dinajpur, Purulia
		Alipurduar	Odisha (30)- Rayagada, Kalahandi, Mayurbhanj, Keonjhar, Sundargarh, Gajapati, Ganjam, Subarnapur, Bhadrak, Boudh, Kandhamal, Angul, Jharsuguda, Bolangir, Koraput, Nabarangpur, Sambalpur, Cuttack, Balasore, Bargarh, Jajpur, Nuapada, Kendrapara, Puri, Khordha, Malkangiri, Jagatsinghpur, Dhenkanal, Nayagarh, Deogarh

Crop/Season	National Yield	Yield i	n Districts
_		Higher than National	Lower than National
Pigeonpea/ Arhar/Tur	913 kg/ha	AP (02)- Guntur, Krishna	AP (11)- Vizianagaram, Nellore, West Godavari, Visakhapatnam, Srikakulam, Prakasham, Kurnool, East Godavari, Kadapa, Chittoor, Ananthapuram
		Bihar (33)-Darbhanga, E.Champaran, Sitamarhi, Sheohar, Vaishali, Muzaffarpur, Madhubani, Bhagalpur, Buxer, Jamui, Begusarai, Mungher, Shekhpura, Khagaria, Gopalgunj, Gaya, Siwan, Supaul, Madhepura, Purnia, Kisangunj, Araria, Aurangabad, Saran, Rohtash, Patna, Jahanabad, W.Champaran, Samastipur, Banka, Nalanda, Arwal, Bhojpur	Bihar(02)- Bhabhua, Nawada
			Chhattisgarh(27)-Balrampur, Raigarh, Mungeli, Durg, Rajnandgaon, Janjgir, Korba, Sarguja, Bemetra, Kondagaon, Kaker, Narayanpur, Surajpur, Jagdalpur, Balod, Dantewada, Sukama, Koria, Kabirdham, Jaspur, Bilaspur, Bijapur, Gariaband, Raipur, Dhamtari, Mahasamund, Baloda bazaar
		Gujarat(29)-Mahisagar, Panchmahal, Surat, Dahod, Banaskantha, Chhotaudepur, Sabarkantha, Vadodara, Patan, Anand, Kutch, Gir Somnath, Surendranagar, Gandhinagar, Bhavnagar, Junagadh, Amreli, Jamnagar, Ahmedabad, Rajkot, Dang, Devbhumi Dwarka, Botad, Kheda, Mehesana, Morbi, Porbandar, Bharuch, Arvalli	Gujarat(04)-Tapi, Narmada, Valsad, Navsari
		Jharkhand (11)-Saraikela, Latehaar, Simdega, Giridih, Gumla, Chatra, Dhanbad, Khunti, Godda, West Singhbum, Palamu	Jharkhand(13)-Ramgarh, Hazaaribagh, Ranchi, Jaamtara, Pakud, Dumka, Devghar, Lohardaga, Koderma, Bokaro, East Singhbhum, Garhwa, Sahebganj

Crop/Season	National Yield	Y	ield in Districts		
		Higher than National	Lower than National		
Pigeonpea/ Arhar/Tur	913 kg/ha	Karnataka (07)- Bengaluru ó rural, Gulbarga, Bidar, Bengaluru ó urban, Chikmagalur, Shimoga, Uttara kannada	Karnataka (20)-Vijayapur, Yadgir, Raichur, Bagalkote, Belgavi, Haveri, Davanagere, Ballari, Hassan, Dharwad, Koppal, Gadag, Chickballapur, Mysore, Mandya, Ramanagaram, Chitradurga, Kolar, Tumkur, Chamarajanagar		
		Maharashtra (20)-Latur, Nagpur, Akola,	Maharashtra (14)-Buldhana, Beed, Gondia, Raigad, Nandurbar,		
		Chandrapur, Gadchiroli, Jalna, Yavatmal, Parbhani, Amravati, Wardha, Bhandara, Hingoli, Dhule,	Satara, Sangli, Kolhapur, Thane, Washim, Solapur, Ratnagiri, Sindhudurg,		
		Nanded, Jalgaon, Aurangabad, Pune, Osmanabad, Nasik, Ahmednagar	Palghar		
		Meghalaya (04)-South west Garo Hills, West Garo Hills, North Garo Hills, East Garo Hills	Meghalaya (01)- South Garo Hills		
		Nagaland (02)-Phek, Peren	Nagaland(09)-Mon, Longleng, Wokha, Dimapur, Kohima, Mokokchung, Kiphiri, Zunheboto, Tuensang		
		Odisha (07)-Kalahandi, Jharsuguda, Sundargarh, Rayagada, Angul, Bolangir, Ganjam	Odisha(21)-Subarnapur, Sambalpur, Kandhamal, Keonjhar, Nuapada, Bargarh, Cuttack, Dhenkanal, Mayurbhanj, Gajapati, Jagatsinghpur, Nabarangpur, Jajpur, Deogarh, Khordha, Kendrapara, Boudh, Koraput, Nayagarh, Balasore, Malkangiri		
		Rajasthan (14)- Bhilwara, Ganganagar, Dholpur, Pratapgarh, Udaipur, Alwar, Baran, Bikaner, Jaisalmer, Kota, Sikar, Tonk, Karoli, Bharatpur	Rajasthan (11)-Banswara, Dungarpur, Sawai madhopur, Dausa, Bundi, Sirohi, Pali, Hanumangarh, Chittor, Jaipur, Jhalawar		
		Tamilnadu (02)-T.V Malai, Vellore	Tamilnadu (27)-Perambalur, Madurai, Dindigul, Krishnagiri, Tiruppur, Thiruvallur, Cuddalore, Pudukkotai, Virudhunagar, Thiruvarur, Coimbatore, Ariyalur, Thanjavur, Kancheepuram, Sivagangai, Tirunelveli, Tuticorin, Dharmapuri, Trichy, Ramanad, Kanyakumari, Salem, Theni, Erode, Namakkal, Villupuram, Karur		

Crop/Season	National Yield	Yield	in Districts		
		Higher than National	Lower than National		
Pigeonpea/	913 kg/ha	Telangana(08)-Adilabad, Komaram bheem,	Telangana(22)-Medchal malkajg, Rangareddy,		
Arhar/Tur		Mancherial, Nirmal, Jagtial, Peddapalli, Karimnagar,	Vikarabad, Sangareddy, Medak, Siddipet, Yadadri,		
		Rajanna	Nalgonda, Suryapet, Nizamabad, Kamareddy,		
			Jayashankar, Mahabubabad, Warangal rural, Warangal		
			urban, Jangaon, Mahabubnagar, Jogulamba, Wanaparthy,		
			Nagarkurnool, Bhadradri, Khammam		
		Uttar Pradesh(42)-Etawah, Kanpur dehat, S. Ravi	Uttar Pradesh(30)-Mainpuri, Agra, Firozabad,		
		das ngr, Kaushambi, Sitapur, Azamgarh,	Kashganj, Etah, Aligarh, Ghazipur, Ambedkar nagar,		
		Farrukhabad, Auraiya, Kannauj, Mau, Allahabad,	Amethi, Faizabad, Sultanpur, Barabanki, Jaunpur, Unnao,		
		Pratapgarh, Ghaziabad, G.buddha ngr., Hapur,	Shahjahanpur, Budaun, Bareilly, Varanasi,		
		Bagpat, Meerut, Mirzpur, Jalaun, Kanpur city,	Muzaffarnagar, Chandauli, Mathura, Sant kabir ngr,		
		Banda, Chitrakut, Sambal, Fatehpur, Rampur,	Basti, Siddharth nagar, Raebareli,		
		Amroha, Bijnor, Ballia, Mahoba, Pilibhit,	Bahraich, Gonda, Balrampur, Shravasti, B.shahar		
		Moradabad, Sonbhadra, Lucknow, Hardoi, Kheri,			
		Hathras, Jhansi, Hamirpur, Deoria, Kushi nagar,			
		Mahrajganj, Gorakhpur			
		Uttrakhand (02)-Tehri Garhwal, Rudraprayag	Uttrakhand (08)-Uttarkashi, Dehradun, Pauri Garhwal,		
			Chamoli, Haridwar, Almora, hampawat, Nainital		
		West Bengal (14)-Malda, Murshidasbad, Hooghly,			
		South 24-Parganas, Nadia, Birbhum, Purba,			
		Bardhaman, Darjeeling, North -Parganas, Paschim			
		Bardhaman, Paschim Medinipur, Jhargram, Purulia,			
		Bankura			

Cron/Soccor	National Yield	Yield in Districts				
Crop/Season	National Yleid	Higher than National	Lower than National			
Mungbean	500 kg/ha	Andhra Pradesh(07)-Kurnool, Visakhapatnam,	Andhra Pradesh (06)- Vizianagaram, Srikakulam,			
		Krishna, East Godavari, Prakasham, Chittoor, Nellore	Guntur, Kadapa, West Godavari, Ananthapuram			
		Bihar (32)-Gaya, Purnia, Kisangunj, Araria, Katihar,	Bihar(06)-Sheohar, E.Champaran, Bhagalpur,			
		Jahanabad, Aurangabad, Shekhpura, Arwal	W.Champaran, Vaishali, Muzaffarpur			
		Khagaria, Begusarai, Lakhisarai, Jamui, Mungher,				
		Samastipur, Darbhanga, Saharsa, Madhubani, Nalanda,				
		Bhabhua, Rohtash, Patna, Bhojpur, Supaul, Buxer,				
		Banka, Gopalgunj, Madhepura, Nawada, Siwan,				
		Sitamarhi, Saran				
		Chhattisgarh (02)-Mungeli, Jagdalpur	Chhattisgarh(24)-Balrampur, Raigarh, Kaker,			
			Kondagaon, Rajnandgaon, Bemetra, Baloda bazaar,			
			Bijapur, Sukama, Dantewada, Dhamtari, Durg,			
			Janjgir, Balod, Jaspur, Koria, Raipur, Gariaband,			
			Kabirmatha, Mahasamund, Korba, Sarguja, Bilaspur,			
			Surajpur			
		Gujarat (05)-Bharuch, Porbandar, Sabarkantha,	Gujarat (24)-Panchmahal, Banaskantha, Patan,			
		Gandhinagar, Bhavnagar	Chhotaudepur, Mahisagar, Botad, Surat, Dahod,			
			Junagadh, Devbhumi Dwarka, Surendranagar,			
			Jamnagar, Kheda, Amreli, Morbi, Rajkot, Gir			
			Somnath, Tapi, Arvalli, Narmada, Vadodara,			
			Navsari, Kutch, Mehesana			
		Jharkhand(20)-Giridih, Khunti, Godda, Simdega,	Jharkhand(03)-Lohardaga, Garhwa, Bokaro			
		Saraikela, Gumla, Dhanbad, Hazaaribagh, Ramgarh,				
		Dumka, West Singhbum, Latehaar, Palamu, Chatra,				
		Ranchi, Koderma, Pakud, Devghar, East Singhbhum,				
		Sahebganj				

C/C	National Yield	Yield in Districts					
Crop/Season		Higher than National	Lower than National				
Mungbean	500 kg/ha	Karnataka (01)-Bidar	Karnataka(25)-Gulbarga, Raichur, Belgavi, Mandya, Bengaluru ó urban, Bengaluru ó rural, Mysore, Dharwad, Yadgir, Haveri, Vijayapur, Chitradurga, Chikmagalur, Bagalkote, Davanagere Tumkur, Ballari, Dakshina kannada, Chamarajanagar, Hassan, Gadag, Uttara kannada, Shimoga, Udupi, Koppal				
		Madhya Pradesh (16) - Sidhi, Bhind, Mandsaur, Narsinghpur, Aagar, Shivpuri, Panna, Hoshangabad Jabalpur, Dhar, Barwani, Damoh, Sagar, Khargone Tikamgarh, Balaghat	Madhya Pradesh (08) - Rewa, Anuppur, Satna, Ujjain, Chattarpur, Betul, Alirajpur, Datia				
		Maharashtra (20)-Akola, Sangli, Buldhana, Bhandara, Kolhapur, Jalna, Jalgaon, Beed, Sindhudurg, Washim, Satara, Ratnagiri, Parbhani, Ahmednagar, Amravati, Aurangabad, Latur, Solapur, Nanded, Hingoli	Maharashtra (12)-Thane, Nagpur, Nandurbar, Yavatmal, Palghar, Nasik, Osmanabad, Gadchiroli Pune, Wardha, Dhule, Chandrapur				
		Odisha(01)-Bhadrak	Odisha (29)-Gajapati, Kalahandi, Ganjam, Boudh Mayurbhanj, Bolangir, Balasore, Keonjhar, Bargarh, Sambalpur, Cuttack, Subarnapur, Khordha, Jagatsinghpur, Nuapada, Malkangiri, Nabarangpur, Jharsuguda, Kendrapara, Angul, Puri, Nayagarh, Jajpur, Deogarh, Koraput, Dhenkanal, Rayagada, Sundargarh, Kandhamal				
			Nagaland (09)-Dimapur, Kohima, Phek, Mokokchung, Tuensang, Mon, Wokha, Zunheboto, Peren				
		Rajasthan (14)-Sikar, Tonk, Ajmer, Bikaner, Jaipur, Nagaur, Ganganagar, Jhunjhunu, Bhilwara, Banswara, Bharatpur, Dungarpur, Karoli, Alwar	Rajasthan (19)-Udaipur, Jhalawar, Sawai madhopur, Dholpur, Bundi, Rajsamand, Baran, Chittor, Kota, Dausa, Hanumangarh, Pratapgarh, Jodhpur, Pali, Sirohi, Jalore, Jaisalmer, Churu, Barmer				
		Tamilnadu (13)- The Nilgris, Salem, Erode, Namakkal, Sivagangai, Thiruvallur, Kancheepuram, T.V Malai, Cuddalore, Pudukkotai, Coimbatore, Karur, Ariyalur	Tamilnadu (16)-Madurai, Trichy, Dharmapuri, Vellore, Villupuram, Thanjavur, Theni, Krishnagiri, Virudhunagar, Tiruppur, Ramanad, Thiruvarur, Nagapatinam, Tuticorin, Dindigul, Tirunelveli				

Cuan/Saasan	National World	Yield in Districts					
Crop/Season	National Yield	Higher than National	Lower than National				
Mungbean	500 kg/ha	Telangana(26)-Nizamabad, Jayashankar, Kamareddy, Warangal rural, Warangal urban, Mahabubabad, Jangaon, Bhadradri, Jagtial, Komaram bheem, Mancherial, Jogulamba, Medchal malkajg, Siddipet, Rangareddy, Vikarabad, Medak, Sangareddy, Karimnagar, Wanaparthy, Peddapalli, Nagarkurnool, Nirmal, Rajanna, Adilabad, Mahabubnagar	Telangana(04)-Khammam, Nalgonda, Yadadri Suryapet				
		Uttar Pradesh (52)-Kannauj, Etawah, Mainpuri, Jaunpur, Farrukhabad, Ghazipur, Chandauli, Bijnor, Saharanpur, Auraiya, Kanpur dehat, S. Ravi das ngr, Muzaffarnagar, Kushi nagar, Gonda, Mahrajganj, Gorakhpur, Ballia, Basti, Deoria, Pilibhit, Bahraich, Azamgarh, Shamli, Mau, Allahabad, Kaushambi, Kanpur city, Fatehpur, Moradabad, Barabanki, Faizabad, Sultanpur, Ambedkar nagar, Amethi, G.buddha ngr., Lalitpur, Pratapgarh, Bagpat, Raebareli, Kheri, Unnao, Sitapur, Hapur, Lucknow, Hardoi, B.shahar, Meerut, Etah, Shahjahanpur, Varanasi, Ghaziabad	Uttar Pradesh(18)-Hathras, Budaun, Firozabad, Kashganj, Agra, Sonbhadra, Mirzpur, Sambal, Hamirpur, Jalaun, Amroha, Rampur, Aligarh, Jhansi, Chitrakut, Banda, Mahoba, Mathura				
		West Bengal (19) -Bankura, Birbhum, Hooghly, Uttar Dinajpur, Jalpaiguri, Purba Bardhaman, Alipurduar, Darjeeling, Paschim Medinipur, Coochbehar, Howrah, Murshidasbad, Purba Medinipur, Dakshin Dinajpur, Nadia, North -Parganas, South 24-Parganas, Purulia, Jhargram	West Bengal(01)- Malda				

Cuon/Cooson	National Wield	Yield in	n Districts		
Crop/Season	National Yield	Higher than National	Lower than National		
Urd	632 kg/ha	Andhra Pradesh (06)-Kurnool, Ananthapuram,	Andhra Pradesh(07)-Chittoor, Guntur, Srikakulam,		
		Visakhapatnam, Krishna, East Godavari, Kadapa	Vizianagaram, West Godavari, Prakasham, Nellore		
		Bihar (29)-Purnia, Jahanabad, Aurangabad, ,			
		Saharsa, Lakhisarai, Kisangunj, Araria, Katihar,			
		Gopalgunj, Bhagalpur, Banka, Siwan, W.Champaran,			
		Vaishali, Nawada, Saran, Supaul, Rohtash, Gaya,			
		Samastipur, Bhojpur, Arwal, Madhubani,			
		Madhepura, Darbhanga, Nalanda, Begusarai,			
		Khagaria, Buxer			
			Chhattisgarh(27)-Kondagaon, Rajnandgaon, Sukama,		
			Dantewada, Bijapur, Gariaband, Jagdalpur, Balrampur,		
			Dhamtari, Raipur, Jaspur, Bemetra, Narayanpur, Sarguja,		
			Kabirmatha, Durg Baloda bazaar, Janjgir, Koria, Mungeli,		
			Surajpur, Mahasamund, Balod, Bilaspur, Kaker, Raigarh,		
			Korba		
		Gujarat(12)-Surat, Junagadh, Dang, Panchmahal,	Gujarat(20)-Vadodara, Kutch, Rajkot, Gir Somnath,		
		Bharuch, Porbandar, Narmada, Tapi, Arvalli, Dahod,	Gandhinagar, Surendranagar, Bhavnagar, Botad,		
		Chhotaudepur, Valsad	Ahmedabad, Jamnagar, Amreli Morbi, Kheda,		
			Mahisagar, Devbhumi Dwarka, Navsari, Sabarkantha,		
		H H 1/40/011 ' 1/4 ' 0' 1	Patan, Banaskantha, Mehesana		
		Jharkhand(19)-Sahebganj, Khunti, Simdega,	Jharkhand(05)-Jaamtara, Garhwa, East Singhbhum,		
		Saraikela, Chatra, Giridih, Gumla, Pakud, Ranchi	Lohardaga, Bokaro		
		West Singhbum, Palamu, Koderma, Godda, Dumka			
		Hazaaribagh, Ramgarh, Latehaar, Dhanbad, Devghar,			

G/G	N - 42 1 X/2 - 1 J	Yield in Districts				
Crop/Season	National Yield	Higher than National	Lower than National			
Urd	632 kg/ha	Madhya Pradesh(20)-Mandsaur, Neemuch, Dhar, Narsinghpur, Burhanpur, Jabalpur, Jhabua, Tikamgarh, Khandwa, Ujjain, Barwani, Damoh, Rajgarh, Ratlam, Gwalior, Balaghat, Mandla, Shahdol, Aagar, Alirajpur, Panna, Hoshangabad, Chattarpur, Bhind	Madhya Pradesh(18)-Bhopal, Umaria, Khargone, Seoni, Satna, Dindori, Rewa, Raisen, Sidhi, Shivpuri, Betul, Singroli, Anuppur, Ashoknagar, Guna, Sagar, Katni, Datia			
			Karnataka(24)-Bidar, Gulbarga, Dakshina kannada, Koppal, Shimoga, Davanagere, Uttara kannada, Bengaluru ó rural, Udupi, Ballari, Mandya, Gadag, Dharwad, Tumkur, Bengaluru ó urban, Yadgir, Belgavi, Mysore, Bagalkote, Chikmagalur, Vijayapur, Hassan, Haveri, Chamarajanagar			
		Maharashtra(12)-Akola, Jalna, Buldhana, Thane Sangli, Washim, Aurangabad, Palghar, Sindhudurg, Jalgaon, Ratnagiri, Nasik	Maharashtra(20)-Satara, Solapur, Latur, Pune Nandurbar, Nagpur, Amravati, Hingoli, Raigad Yavatmal, Kolhapur, Ahmednagar, Beed, Gadchiroli, Parbhani, Osmanabad, Dhule, Wardha Chandrapur, Nanded Odisha(30)-Kalahandi, Kendrapara, Puri,			
			Jagatsinghpur, Cuttack, Bhadrak, Jajpur, Gajapati, Subarnapur, Khordha, Ganjam, Balasore, Keonjhar, Dhenkanal, Angul, Malkangiri, Boudh, Sundargarh, Mayurbhanj, Sambalpur, Nuapada, Nayagarh, Bargarh, Bolangir, Rayagada, Koraput, Jharsuguda, Nabarangpur, Deogarh, Kandhamal			
		Rajasthan (20)-Churu, Jhunjhunu, Pratapgarh, KotaBaran, Dungarpur, Sawai madhopur, Ajmer, Banswara, Jhalawar, Barmer, Jodhpur, Bharatpur, Hanumangarh, Pali, Dausa, Rajsamand, Jaipur, Dholpur, Karoli	Rajasthan(09)-Bundi, Bikaner, Jalore, Tonk, Bhilwara, Sirohi, Udaipur, Chittor, Ganganagar			

Cwan/Saasan	National Yield	Yield in	Districts
Crop/Season	National Yleid	Higher than National	Lower than National
Urd	632 kg/ha	Tamilnadu (07)- Erode, Thanjavur, Perambalur, Salem, Krishnagiri, Vellore, Madurai	Tamilnadu(23)-Karur, Sivagangai, Villupuram, Coimbatore, Kancheepuram, Cuddalore, Pudukkotai, Trichy, Kanyakumari, Dharmapuri, T.V Malai, Ariyalur, Theni, Thiruvarur, Thiruvallur, Namakkal, Virudhunagar, Dindigul, Tiruppur, Ramanad, Nagapatinam, Tuticorin, Tirunelveli
		Telangana(29)-Nizamabad, Kamareddy, Rangareddy, Vikarabad, Warangal urban, Karimnagar, Jangaon, Khammam, Mahabubnagar, Suryapet, Jagtial, Peddapalli, Nalgonda, Wanaparthy, Jogulamba, Mahabubabad, Bhadradri, Rajanna, Jayashankar, Nagarkurnool, Warangal rural, Yadadri, Siddipet, Sangareddy, Medak, Komaram bheem, Nirmal, Adilabad, Mancherial	
		Uttar Pradesh(27)-Varanasi, Jaunpur, Shahjahanpur, Sambal, Kannauj, Amroha, Bijnor, Kanpur dehat, Farrukhabad, Auraiya, Rampur, Chandauli, Muzaffarnagar, Ghazipur, Etawah, Kheri, Saharanpur, Shamli, Ballia, G.buddha ngr., Pilibhit, Mahrajganj, Mau, Bareilly, Ambedkar nagar, Kanpur city, Kushi nagar	Uttar Pradesh(48)-Moradabad, Pratapgarh, Gorakhpur, Meerut, Bagpat, Firozabad, Hapur, Gonda, Budaun, B.shahar, Unnao, Mainpuri, Ghaziabad, Basti, Azamgarh, Etah, Kashganj, Hathras, Bahraich, Hardoi, S. Ravi das ngr, Sonbhadra, Mirzpur, Sultanpur, Aligarh, Lalitpur, Shravasti, Balrampur, Agra, Lucknow, Mathura, Siddharth nagar, Allahabad, Deoria, Sant kabir ngr, Kaushambi, Amethi, Chitrakut, Mahoba, Hamirpur, Sitapur, Barabanki, Jalaun, Faizabad, Jhansi, Fatehpur, Raebareli, Banda

Annexure-VIII

Crop-wise major Potential Pulse State/Districts for the proposed pulse export zone

1. Arhar

(Area-Lakh ha, Production-Lakh Tonnes)

State	District	Area	%	to	Production	% t	
			India	State		India	State
Maharashtra	Latur	1.18	2.21	8.22	3.98	8.18	26.62
	Yavatmal	1.74	3.25	12.12	2.34	4.80	15.65
	Amravati	1.34	2.52	9.33	1.55	3.19	10.37
	Nagpur	0.58	1.09	4.04	1.46	3.00	9.77
	Akola	0.6	1.12	4.18	1.33	2.72	8.90
	Osmanabad	1.17	2.20	8.15	1.16	2.37	7.76
Total Above		6.61	12.39	46.03	11.82	24.26	79.06
Total State		14.36	26.90		14.95		
Karnataka	Gulbarga	3.89	7.29	32.04	5.42	11.12	59.63
	Vijayapur	3.89	7.29	32.04	3.30	6.78	36.30
Total Above		7.78	14.58	64.09	8.72	17.90	95.93
Total State		12.14	22.74		9.09		
Madhya Pradesh	Narsinghpur	0.66	1.24	9.57	1.29	2.64	16.50
Total State	Total State		12.93		7.82		
Gujarat	Bharuch	1.06	1.98	30.55	1.14	2.33	28.43
Total State		3.47	6.50		4.01		
All India		53.38			48.73		

Source: All India and State- DES, GoI and District- SDAs 2016-17

2. Gram

(Area-Lakh ha, Production-Lakh Tonnes)

State	District	Area	%	to	Production	% to		
			India	State		India	State	
Madhya	Sagar	1.78	1.85	5.52	2.47	2.63	6.96	
Pradesh	Damoh	1.74	1.80	5.38	2.43	2.59	6.85	
	Satna	1.20	1.25	3.72	2.10	2.24	5.93	
	Vidisha	1.72	1.79	5.35	1.99	2.13	5.62	
	Narsinghpur	0.98	1.02	3.05	1.88	2.00	5.30	
	Raisen	1.20	1.25	3.72	1.79	1.91	5.04	
Total Above		8.62	8.96	26.75	12.65	13.49	35.70	
Total State		32.22	33.47		35.44	37.79		
Karnataka	Gulbarga	1.85	1.92	18.45	2.17	2.32	36.71	
Total State		10.03	10.42		5.92	6.31		
Maharashtra	Latur	2.10	2.18	10.87	2.50	2.66	14.52	
	Nanded	1.14	1.19	5.93	1.76	1.87	10.21	
Total Above		3.24	3.37	16.80	4.25	4.53	24.73	
Total State		19.30	20.05		17.19	18.33		
Rajasthan	Rajasthan Bikaner		2.35	14.64	1.94	2.07	13.74	
Total State	Total State		16.08		14.09	15.02		
All India		96.26			93.78			

Source: All India and State- DES, GoI and District- SDAs 2016-17

3. Mungbean

(Area-Lakh ha, Production-Lakh Tonnes)

State	District	Area	% to Proc		Production	%	to
			India	State		India	State
Rajasthan	Nagaur	5.58	12.90	34.09	3.36	15.52	41.48
	Jodhpur	2.77	6.40	16.92	1.10	5.08	13.58
	Pali	2.79	6.45	17.04	1.07	4.94	13.21
	Ajmer	1.05	2.43	6.41	0.74	3.42	9.14
	Jaipur	1.07	2.47	6.54	0.73	3.37	9.01
	Churu	1.76	4.07	10.75	0.50	2.31	6.17
Total Above	Total Above		34.71	91.75	7.50	34.64	92.59
Total States		16.37	37.83		8.10	37.41	
Maharashtra	Jalna	0.42	0.97	9.46	0.29	1.34	11.15
	Ahmednagar	0.48	1.11	10.81	0.29	1.34	11.15
	Akola	0.31	0.72	6.98	0.27	1.25	10.38
Total Above	Total Above		2.80	27.25	0.85	3.93	32.69
Total States		4.44	10.26		2.60	12.01	
Odisha	Ganjam	1.08	2.50	33.13	0.53	2.45	39.55
Total State		3.26	7.53		1.34	6.19	
All India		43.27			21.65		

Source: All India and State- DES, GoI and District- SDAs 2016-17

4. Urdbean

(Area-Lakh ha, Production-Lakh Tonnes)

State	District	Area	% to		Production	% to	
			India	State		India	State
Andhra Pradesh	Krishna	1.59	3.55	31.80	1.23	4.35	37.45
	Guntur	1.00	2.23	20.00	0.51	1.81	15.56
	Kurnool	0.34	0.76	6.82	0.49	1.74	14.95
Total Above		2.93	6.55	58.62	2.24	7.90	67.96
Total State		5.00	11.17		3.29	11.62	
Madhya Pradesh	Tikamgarh	1.16	2.59	9.64	1.03	3.63	12.57
	Chattarpur	1.00	2.22	8.28	0.64	2.27	7.85
	Damoh	0.65	1.46	5.44	0.52	1.85	6.40
	Mandsaur	0.40	0.90	3.34	0.42	1.49	5.18
Total Above		3.21	7.17	26.70	2.61	9.23	32.00
Total State		12.03	26.86		8.17	28.85	
Rajasthan	Bundi	0.91	2.03	19.08	0.57	2.01	18.63
Total State		4.77	10.65		3.06	10.81	
Tamilnadu	Villupuram	0.95	2.13	22.12	0.53	1.88	19.38
Total State		4.30	9.60		2.74	9.68	
Uttar Pradesh	Lalitpur	1.61	3.59	24.98	0.88	3.10	24.84
Total State	Total State		14.38		3.53	12.46	
All India		44.78			28.32		

Source: All India and State- DES, GoI and District- SDAs 2016-17

5. Lentil

(Area-Lakh ha, Production-Lakh Tonnes)

State	District	Area	% to		Production	% to	
			India	State		India	State
Madhya	Satna	0.54	3.87	10.21	0.58	5.39	12.98
Pradesh	Sagar	0.46	3.32	8.70	0.56	5.18	12.53
	Narsinghpur	0.42	3.02	7.94	0.51	4.71	11.41
	Raisen	0.30	2.16	5.67	0.50	4.62	11.19
	Vidisha	0.39	2.80	7.37	0.50	4.61	11.19
	Dindori	0.37	2.65	6.99	0.37	3.40	8.28
	Rajgarh	0.34	2.47	6.43	0.36	3.35	8.05
Total Above		2.82	20.29	53.31	3.38	31.26	75.62
Total State		5.29	37.95		4.47	41.50	
Uttar Pradesh	Bahraich	0.51	3.64	10.43	0.43	4.03	11.56
	Jalaun	0.30	2.16	6.13	0.40	3.75	10.75
	Banda	0.48	3.46	9.82	0.38	3.49	10.22
Total Above		1.29	9.26	26.38	1.21	11.27	32.53
Total State		4.89	35.08		3.72	34.54	
All India		13.94			10.77		

Source: All India and State- DES, GoI and District- SDAs 2016-17

6. Field pea

(Area-Lakh ha, Production-Lakh Tonnes)

State	District	Area	% to		Production	% to	
			India	State		India	State
Uttar	Jalaun	0.79	8.50	20.57	1.40	15.72	33.49
Pradesh	Lalitpur	0.70	7.46	18.23	1.08	12.11	25.84
	Jhansi	0.66	7.06	17.19	0.92	10.40	22.01
	Mahoba	0.42	4.52	10.94	0.34	3.78	8.13
Total Above		2.57	27.54	66.93	3.74	42.01	89.47
Total Sta	Total State		36.29		4.18	41.35	
Madhya	Panna	0.52	5.57	11.18	0.87	9.76	24.44
Pradesh	Jabalpur	0.52	5.57	11.18	0.81	9.15	22.75
	Datia	0.35	3.75	7.53	0.58	6.51	16.29
	Narsinghpur	0.39	4.18	8.39	0.48	5.39	13.48
	Chattarpur	0.25	2.68	5.38	0.34	3.79	9.55
	Dindori	0.28	3.00	6.02	0.32	3.57	8.99
Total Above		2.31	24.75	49.68	3.40	38.17	95.51
Total State		4.65	43.95		3.56	35.21	
All India		10.58			10.11		

Source: All India and State- DES, GoI and District- SDAs 2016-17

CONTACT DETAILS

i) DAC&FW: CROPS DIVISION

Designation	Name	Contact number	Email	
Agriculture Commissioner				
Agriculture Commissioner	Dr. S.K. Malhotra	9968978191(M) 011-23383549 (O)	ag.comm@nic.in	
PPS to AC	R.N. Sharma	011-23381012(O)	rns119@yahoo.co.in	
Joint Secretary (Crops)				
Joint Secretary (Crops)	Dr. B. Rajender	9599428809 (M) 011-23381176 (O)	b.rajender@ias.nic.in	
PA to JS (Crops)	Shri R.N. Bansal	9810863107 (M) 011-23381176 (O)	rnbansal164@gmail.com	
Additional Commissioner (Crops)				
Additional Comm. (NFSM)/CC	Dr. A.P. Singh	9811137009 (M) 011-23388579	anilpratapsingh1962@yahoo .co.in	
Additional Comm. (NFSM)/Crops	Dr. S.S.Tomar	08859245361 (M) 011-23387175	drtomarss@yahoo.com	
Additional Comm. (Oilseeds)	Dr. Anupam Barik,	011 -23384618	adcoilseeds-agri@gov.in	
Additional Comm. I/C Director (Crops)	Sh. Tarun Johri	011 -23383980	addrty.crops.asri@gov.in	
DC (Crops)	Dr. M.N. Singh (NFSM & BGREI)	9811572591(M) 011-23385781	mnsingh1959@gmail.com	
NFSM Cell	<u> </u>	<u>.</u>		
AD (Crops), Pulses	Shri. Ashok Kumar Singh	9650669923(M)	aksadcrops@yahoo.in	
AD (Crops),(Coordination & CC)	Dr. Manoj Kumar	9891090571(M)	mksadcrops@gmail.com	
AD (Crops), Research Projects	Smt. G. Sandhya	9873319305(M)	gsrani71@gmail.com	
Sr. Programmer(NFSM-Website)	Sh. Suraj Prakash	8826836683(M)	suraj0786@gmail.com	
Technical Support Group (TSG Ce	II)			
Consultant (NFSM-Rice)	Sh. Shamsher Singh	9810284556 (M)	shamsher123@yahoo.com	
Consultant (NFSM-Wheat)	Dr. P K Saha	09953124101 (M)	pksaha58@gmail.com	
Consultant (NFSM-Coarse Cereals)	Dr. Virender Singh Pahil	9466114433/ 08076047919	dr.vspahil@gmail.com	
Consultant (NFSM-Oilseeds)	Dr. M Dutta,	011-23383476	manoranjandutta@rediffma.co m	
Consultant (NFSM-Oilseeds)	SH. S K Dalal	011-23383476	skdalal@gmail.com	

ii) CROP DEVELOPMENT DIRECTORATES

Dr. Virendra Singh, Director	Dr. M N Singh, Director		
Directorate of Rice Development,	Directorate of Wheat Development,		
DAC&FW, Ministry of Agri.& Farmers Welfare,	C.G.O.Complex ó I, 3rd Floor,, Kamla Nehru Nagar,		
5th Floor, 'G' Wing, Karpuri Thakur Sadan,	Ghaziabad-201002 (U.P.)		
Kendriya Karmachari Parisar (GPOA),	Tele: 0120 - 2711380, 2710897		
Ashiana Digha Road,	Fax: 0120 - 2711380		
Patna ó 800 025 Bihar Tel: 0612-2262720, 2262843	E-mail:, dwd.wheat@gmail.com		
Fax: 0612-2262720			
E-mail: drdpatna@nic.in			
Dr. A. K.Tiwari, Director	DrK. Manoharan, Director		
Directorate of Pulses Development,	Directorate of Jute Development,		
6th Floor, Vindhyachal Bhavan,	Nizam Palace Campus ,234/4, Acharya Jagdish Bose		
Bhopal-462004 (M.P.)	Road, Kolkata- 700020 (West Bengal)		
Tel: 0755-2550353, 2572313	Tel. no: 033-22879465/9337/9189		
Telefax: 0755-2571678	Fax :033-22879521		
E-mail: dpd.mp@nic.in	E-mail: djd@nic.in		
Dr. Man Singh, Director	Dr. Subhash Chandra, Director-Incharge		
Directorate of Sugarcane Development,	Directorate of Millets Development,		
8th Floor, Hall No.3, KendriyaBhavan, Aliganj,	Kendriya Sadan Room No. 210, 2nd Floor, Block-		
Lucknow - 226024 (U.P)	A,Sector-10,Vidyadhar Nagar, Jaipur-302023		
Tel. no.: 0522-2323913	Tel: 0141-2233004		
Fax: 0522- 2323913;	Fax: 0141-2235631		
E-mail: dsd@nic.in;	E-mail: dmdrj00@nic.in		
Shri R. P Singh, Director	Dr. B. K. Shirivastv, Director-Incharge		
Directorate of Cotton Development,	Directorate of Oilseeds Development,		
Bhoomi Sarvekshan Bhavan,	Telhan Bhavan, Himayat Nagar,		
Near Centre Point School,	Hyderabad-500029 (A.P.)		
Katol Road, Nagpur-440013 (Maharashtra)	Tel. No.:040-23224381;		
Tel. No 0712-2585831	Fax :040-23224381		
Fax: 0712-2595505	E-mail: dod@nic.in;		
E-mail: directordocd@rediffmail.com			

iii) STATES/UTS: DIRECTOR (AGRICULTURE)

State	Name of the officer	Contact No. &Fax	Email
Andhra	Shri Dhananjaya	09849908692 (M)	comag@nic.in
Pradesh	Reddy	0863-2216461(O)	comagr.ap@gmail.com
		0863-2216463 (F)	
Arunachal	Shri Hage Kano	09436252188 (M)	rkvyita@rediffmail.com
Pradesh		0360-2244252(O)2244252 (F)	
Assam	Shri B.N. Faikia	07670004458 (M)	agridept@nic.in
		0361-2332215/2333877 (O)	
		0361-2332796 (F)	
Bihar	Shri Himanshu	09431818704 (M)	diragribih@nic.in
	Kumar Rai	0612-2204245/2215895 (O)	diragri.bih@gmail.com
		0612-2204245 (F)	
Chhattisgarh	Shri M.S.	09425542364 (M)	diragricg@nic.in
	Kerketta	0771-2442015 (O) 2442036 (F)	
Gujarat	Dr. B.R. Shah	09978405999 (M)	diragr@gujarat.gov.in
U		079-23256073 (O)	
		079-23256227/ 23256159 (F)	
Goa	Shri Ulhas	09422457328 (M)	diragri.goa@nic.in
	Pai Kakode	0832-2465443/2465840 (O)	
		0832-2465441 (F)	
Haryana	Shri	09468296051 (M)	agriharyana2009@gmail.com
	Bhupinder	0172-2570662 (O)	agriculture@hry.nic.in
	Singh	0172-2563242 (F)	
Himachal	Shri Somraj	09418031268 (M)	krishinidesh@yahoo.com
Pradesh	Kalia	0177-2830162/ 2830618/	
		2830174 (O)	
		0177-2830612 (F)	
Jammu &	Shri Showkat	09419036454 (M)	diragrikmr@gmail.com
Kashmir	Ahmad Beigh	0194-2310675 (O) 2311569 (F)	
Jharkhand	Jatashankar	09431115706 (M)	directoragriculture@gmail.com
Shri	Choudhary	0651-6550124 (O) 2233549 (F)	
Karnataka	Shri B.Y. Srinivas	07259004007 (M)	agridir@kar.nic.in
		080-22242746 (O)22212818 (F)	
Kerala	Biju Brabhakar,	09447063200 (M)	krishidirector@gmail.com
	IAS	0471-2304480 (O) 2304230 (F)	_
Madhya	Shri M.L.	09425036356 (M)	mpkrishi@mp.gov.in
Pradesh	Meena	0755-2551336 (O) 2572468 (F)	diragri@mp.gov.in
Maharashtra	Shri Vikas	09420490003 (M)	commagricell@gmail.com
	Deshmukh	020-26123648/ 26126150 (O)	com.pune@agri.maharashtra.gov.i
	Commissioner of	020-26127707 (F)	<u>n</u>
	Agriculture		
Manipur	Shri Louis	09436032021 (M)	I.ngasainao@gmail.com
r ··	Ngasainao	0385-2452019 (O)	amdmn@nic.in
Meghalaya	Ms.	09436104524 (M)	directoratejdashillong@gmaill.com
ogiminju	D.Syiemiong	0364-2222460 (O) 2223228 (F)	s ocoracojado inforigegridan.com
Mizoram	Dr. C.	09436142745 (M)	Zara.agril@gmail.com
MIZOIGIII	Lalzarliana	0389-2322437 (O) 2322511(F)	<u> zara.agin eginan.com</u>
Nagaland	Smt. Rongsdninla	09436001048 (M)	agrilanngl@gov.in
ragaiailu	Sint. Kongsumma	0370-2243116 (O) 2243116 (F)	agmanngregov.m
		0370-22 1 3110 (O) 22 1 3110 (F)	_

State	Name of the officer	Contact No. &Fax	Email
Punjab	Shri J.S.	09915938757 (M)	director.agri@punjab.gov.in
-	Bains	0172-2970602(O) 2970609 (F)	
Rajasthan Shri Ambrish		09829084089 (M)	dir_agri@rediffmail.com
	Kumar, IAS	0141-2227089 (O)2227088 (F)	
Sikkim	Shri Kiran	09832017868 (M)	kiran_pradhan_2000@yahoo.com
	Pradhan	03592-232942 (O) 231892 (F)	
Uttarakhand	Shri C. Gauri	09410996912 (M)	dir.agri.uttarakhand@gmail.com
	Shankar	0135-2772677 (O) 2771881(F)	
Tamil Nadu	M. Rajendran,	09751534567 (M)	diragri@tn.nic.in
	IAS	044-28524894 (O) 28521998 (F)	
Telangana	Smt. G.D.	09866894624 (M)	agriculture.telangana@gmail.com
_	Priyadarshini	040-23232107 (O) 23383520 (F)	
Tripura	Dr. D.P.	09436124922 (M)	krishibhawantripura@gmail.com
_	Sarkar	0381-2323778 (O)2323883 (F)	tmcelltripura@gmail.com
Uttar Pradesh	Shri Mukesh	09235629301 (M)	dirag@nic.in
	Kumar Srivastava	0522-2205869 (O) 2206582 (F)	
West Bengal	Dr. Paritosh	09433829300 (M)	direct6oragriculturewb@gmail.com
•	Bhattacharyya	033-22627713 (O) 22145307 (F)	
Delhi	Ms Sakshi	9910182340 (M)	dagri@sansad.nic.in
	Mittal	011-23818601(O)23818298 (F)	
Puducherry	Dr.A.Ramamurthy	09443235329 (M)	
,		0413-2336977 (O) 2337121(F)	
Andaman &	Shri Anjan Kumar	09434272016 (M)	diragri@gmail.com
Nicobar	Das	03192-233257 (F)	
Daman &	Shri Debendra Dalai	08980090090 (M)	ddir-agri-dd@nic.in
Diu		0260-2230856 (O)	
Chandigarh	Shri. Bhupinder	09468296051 (M)	agriharyana2009@gmail.com
J	Singh	0172-2570662 (O)	agriculture@hry.nic.in





Directorate of Pulses Development Ministry of Agriculture & Farmers Welfare Department of Agriculture, Cooperation & Farmers Welfare Government of India

6th Floor, Vindhyachal Bhavan Bhopal – 462004 (M.P.)

Phone: 0755-2550353/ 2572313

Telefax: 0755-2571678 E-mail: dpd.mp@nic.in Website: www.dpd.gov.in







https//:www.agricoop.nic.in https//:www.nfsm.gov.in

M-kisan portal - https://mkisan.gov.in Farmers portal - https://farmer.gov.in Kisan Call Centre & Toll Free 1800-180-1551